


SOLUTIONS MANUAL




Chapter 2

Linear and Quadratic Functions

Section 2.1

- From the equation $y = 2x - 3$, we see that the y-intercept is -3 . Thus, the point $(0, -3)$ is on the graph. We can obtain a second point by choosing a value for x and finding the corresponding value for y . Let $x = 1$, then $y = 2(1) - 3 = -1$. Thus, the point $(1, -1)$ is also on the graph. Plotting the two points and connecting with a line yields the graph below.



- $$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - 5}{-1 - 2} = \frac{-2}{-3} = \frac{2}{3}$$
- $$f(2) = 3(2)^2 - 2 = 10$$

$$f(4) = 3(4)^2 - 2 = 46$$

$$\frac{\Delta y}{\Delta x} = \frac{f(4) - f(2)}{4 - 2} = \frac{46 - 10}{4 - 2} = \frac{36}{2} = 18$$
- $$60x - 900 = -15x + 2850$$

$$75x - 900 = 2850$$

$$75x = 3750$$

$$x = 50$$

The solution set is $\{50\}$.
 - $$5x - 4 \leq 3x + 6$$

$$5x - 3x = 6 + 4$$


$$2x = 10$$

$$x = 5$$


The solution set is $(-\infty, 5]$
- $$f(0) = -5(0) - 14 = 0 - 14 = -14$$
- True

- slope; y-intercept
- $-4; 3$
- positive
- True
- False. If x increases by 3, then y increases by 2.
- False. The y-intercept is 8. The average rate of change is 2 (the slope).

- $$f(x) = 2x + 3$$
 - Slope = 2; y-intercept = 3
 - Plot the point $(0, 3)$. Use the slope to find an additional point by moving 1 unit to the right and 2 units up.



- Domain: $(-\infty, \infty)$; Range: $(-\infty, \infty)$
 - average rate of change = 2
 - increasing
- $$g(x) = 5x - 4$$
 - Slope = 5; y-intercept = -4
 - Plot the point $(0, -4)$. Use the slope to find an additional point by moving 1 unit to the right and 5 units up.




Section 2.1: Properties of Linear Functions and Linear Models

- c. Domain: $(-\infty, \infty)$; Range: $(-\infty, \infty)$
- d. average rate of change = 5
- e. increasing

15. $h(x) = -3x + 4$


- a. Slope = -3 ; y-intercept = 4
- b. Plot the point $(0, 4)$. Use the slope to find an additional point by moving 1 unit to the right and 3 units down.



- c. Domain: $(-\infty, \infty)$; Range: $(-\infty, \infty)$
- d. average rate of change = -3
- e. decreasing

16. $p(x) = -x + 6$

- a. Slope = -1 ; y-intercept = 6
- b. Plot the point $(0, 6)$. Use the slope to find an additional point by moving 1 unit to the right and 1 unit down.




- c. Domain: $(-\infty, \infty)$; Range: $(-\infty, \infty)$
- d. average rate of change = -1
- e. decreasing

17. $f(x) = \frac{1}{4}x - 3$

- a. Slope = $\frac{1}{4}$; y-intercept = -3


- b. Plot the point $(0, -3)$. Use the slope to find an additional point by moving 4 units to the right and 1 unit up.



- c. Domain: $(-\infty, \infty)$; Range: $(-\infty, \infty)$
- d. average rate of change = $\frac{1}{4}$
- e. increasing

18. $h(x) = -\frac{2}{3}x + 4$


- a. Slope = $-\frac{2}{3}$; y-intercept = 4
- b. Plot the point $(0, 4)$. Use the slope to find an additional point by moving 3 units to the right and 2 units down.



- c. Domain: $(-\infty, \infty)$; Range: $(-\infty, \infty)$
- d. average rate of change = $-\frac{2}{3}$
- e. decreasing

19. $F(x) = 4$

- a. Slope = 0; y-intercept = 4
- b. Plot the point $(0, 4)$ and draw a horizontal line through it.




Chapter 2: Linear and Quadratic Functions

- c. Domain: $(-\infty, \infty)$; Range: $\{y \mid y = 4\}$
- d. average rate of change = 0
- e. constant

20. $G(x) = -2$

- a. Slope = 0; y-intercept = -2
- b. Plot the point $(0, -2)$ and draw a horizontal line through it.




- c. Domain: $(-\infty, \infty)$; Range: $\{y \mid y = -2\}$
- d. average rate of change = 0
- e. constant

21. $g(x) = 2x - 8$

- a. $2x - 8 = 0$
 $2x = 8$
 $x = 4$


- b. y-intercept = -8



22. $g(x) = 3x + 12$

- a. $3x + 12 = 0$
 $3x = -12$
 $x = -4$


- b. y-intercept = 12



23. $f(x) = -5x + 10$

- a. $-5x + 10 = 0$
 $-5x = -10$
 $x = 2$


- b. y-intercept = 10



24. $f(x) = -6x + 12$

- a. $-6x + 12 = 0$
 $-6x = -12$
 $x = 2$

- b. y-intercept = 12




25. $H(x) = -\frac{1}{2}x + 4$

- a. $-\frac{1}{2}x + 4 = 0$
 $-\frac{1}{2}x = -4$
 $x = 8$

Section 2.1: Properties of Linear Functions and Linear Models

- b. y-intercept = 4




26. $G(x) = \frac{1}{3}x - 4$

a. $\frac{1}{3}x - 4 = 0$

$$\frac{1}{3}x = 4$$

$$x = 12$$

- b. y-intercept = -4



27.

x	y	Avg. rate of change = $\frac{\Delta y}{\Delta x}$
-2	4	
-1	1	$\frac{1-4}{-1-(-2)} = \frac{-3}{1} = -3$
0	-2	$\frac{-2-1}{0-(-1)} = \frac{-3}{1} = -3$
1	-5	$\frac{-5-(-2)}{1-0} = \frac{-3}{1} = -3$
2	-8	$\frac{-8-(-5)}{2-1} = \frac{-3}{1} = -3$

Since the average rate of change is constant at -3, this is a linear function with slope = -3. The y-intercept is (0, -2), so the equation of the line is $y = -3x - 2$.

28.

x	y	Avg. rate of change = $\frac{\Delta y}{\Delta x}$
-2	$\frac{1}{4}$	
-1	$\frac{1}{2}$	$\frac{(\frac{1}{2}-\frac{1}{4})}{-1-(-2)} = \frac{\frac{1}{4}}{1} = \frac{1}{4}$
0	1	$\frac{(1-\frac{1}{2})}{0-(-1)} = \frac{\frac{1}{2}}{1} = \frac{1}{2}$
1	2	
2	4	

Since the average rate of change is not constant, this is not a linear function.

29.

x	y	Avg. rate of change = $\frac{\Delta y}{\Delta x}$
-20	-8	
-10	-3	$\frac{-3-(-8)}{-10-(-20)} = \frac{5}{10} = \frac{1}{2}$
0	0	$\frac{0-(-3)}{0-(-10)} = \frac{3}{10}$
10	1	
20	0	

Since the average rate of change is not constant, this is not a linear function.

30.

x	y	Avg. rate of change = $\frac{\Delta y}{\Delta x}$
-20	-40	
-10	0	$\frac{0-(-40)}{-10-(-20)} = \frac{40}{10} = 4$
0	40	$\frac{40-0}{0-(-10)} = \frac{40}{10} = 4$
10	80	$\frac{80-40}{10-0} = \frac{40}{10} = 4$
20	120	$\frac{120-80}{20-10} = \frac{40}{10} = 4$

Since the average rate of change is constant at 4, this is a linear function with slope = 4. The y-intercept is (0, 40), so the equation of the line is $y = 4x + 40$.

Chapter 2: Linear and Quadratic Functions

31.

x	y	Avg. rate of change = $\frac{\Delta y}{\Delta x}$
-2	-26	
-1	-4	$\frac{-4 - (-26)}{-1 - (-2)} = \frac{22}{1} = 22$
0	2	$\frac{2 - (-4)}{0 - (-1)} = \frac{6}{1} = 6$
1	-2	
2	-10	

Since the average rate of change is not constant, this is not a linear function.

32.

x	y	Avg. rate of change = $\frac{\Delta y}{\Delta x}$
-2	-4	
-1	-3.5	$\frac{-3.5 - (-4)}{-1 - (-2)} = \frac{0.5}{1} = 0.5$
0	-3	$\frac{-3 - (-3.5)}{0 - (-1)} = \frac{0.5}{1} = 0.5$
1	-2.5	$\frac{-2.5 - (-3)}{1 - 0} = \frac{0.5}{1} = 0.5$
2	-2	$\frac{-2 - (-2.5)}{2 - 1} = \frac{0.5}{1} = 0.5$

Since the average rate of change is constant at 0.5, this is a linear function with slope = 0.5. The y-intercept is (0, -3), so the equation of the line is $y = 0.5x - 3$.

33.

x	y	Avg. rate of change = $\frac{\Delta y}{\Delta x}$
-8	41	
-5	29	$\frac{29 - 41}{-5 - (-8)} = \frac{-12}{3} = -4$
0	9	$\frac{9 - 29}{0 - (-5)} = \frac{-20}{5} = -4$
2	1	$\frac{1 - 9}{2 - 0} = \frac{-8}{2} = -4$
6	-15	$\frac{-15 - 1}{6 - 2} = \frac{-16}{4} = -4$

Since the average rate of change is constant at -4, this is a linear function with slope = -4. The y-intercept is (0, 9), so the equation of the line is $y = -4x + 9$ or $y = 8$.

34.

x	y	Avg. rate of change = $\frac{\Delta y}{\Delta x}$
-10	0	
-5	2	$\frac{2 - 0}{-5 - (-10)} = \frac{1}{5}$
0	4	$\frac{4 - 2}{0 - (-5)} = \frac{2}{5}$
3	9	
8	16	

Since the average rate of change is not constant, this is not a linear function.

35. $f(x) = 4x - 1$; $g(x) = -2x + 5$

a. $f(x) = 0$

$$4x - 1 = 0$$

$$x = \frac{1}{4}$$

b. $f(x) > 0$

$$4x - 1 > 0$$

$$x > \frac{1}{4}$$

The solution set is $\left\{x \mid x > \frac{1}{4}\right\}$ or $\left(\frac{1}{4}, \infty\right)$.

c. $f(x) = g(x)$

$$4x - 1 = -2x + 5$$

$$6x = 6$$

$$x = 1$$

d. $f(x) \leq g(x)$


$$4x - 1 \leq -2x + 5$$

$$6x \leq 6$$

$$x \leq 1$$

The solution set is $\{x \mid x \leq 1\}$ or $(-\infty, 1]$.

e.



Section 2.1: Properties of Linear Functions and Linear Models

36. $f(x) = 3x + 5$; $g(x) = -2x + 15$

a. $f(x) = 0$

$$3x + 5 = 0$$

$$x = -\frac{5}{3}$$

b. $f(x) < 0$

$$3x + 5 < 0$$

$$x < -\frac{5}{3}$$

The solution set is $\left\{x \mid x < -\frac{5}{3}\right\}$ or $\left(-\infty, -\frac{5}{3}\right)$.

c. $f(x) = g(x)$

$$3x + 5 = -2x + 15$$

$$5x = 10$$

$$x = 2$$

d. $f(x) \geq g(x)$


$$3x + 5 \geq -2x + 15$$

$$5x \geq 10$$

$$x \geq 2$$

The solution set is $\{x \mid x \geq 2\}$ or $[2, \infty)$.

e.



37. a. The point (40, 50) is on the graph of $y = f(x)$, so the solution to $f(x) = 50$ is $x = 40$.

b. The point (88, 80) is on the graph of $y = f(x)$, so the solution to $f(x) = 80$ is $x = 88$.

c. The point (-40, 0) is on the graph of $y = f(x)$, so the solution to $f(x) = 0$ is $x = -40$.

d. The y-coordinates of the graph of $y = f(x)$ are above 50 when the x-coordinates are larger

than 40. Thus, the solution to $f(x) > 50$ is $\{x \mid x > 40\}$ or $(40, \infty)$.

e. The y-coordinates of the graph of $y = f(x)$ are below 80 when the x-coordinates are smaller than 88. Thus, the solution to $f(x) \leq 80$ is $\{x \mid x \leq 88\}$ or $(-\infty, 88]$.

f. The y-coordinates of the graph of $y = f(x)$ are between 0 and 80 when the x-coordinates are between -40 and 88. Thus, the solution to $0 < f(x) < 80$ is $\{x \mid -40 < x < 88\}$ or $(-40, 88)$.

38. a. The point (5, 20) is on the graph of $y = g(x)$, so the solution to $g(x) = 20$ is $x = 5$.

b. The point (-15, 60) is on the graph of $y = g(x)$, so the solution to $g(x) = 60$ is $x = -15$.

c. The point (15, 0) is on the graph of $y = g(x)$, so the solution to $g(x) = 0$ is $x = 15$.

d. The y-coordinates of the graph of $y = g(x)$ are above 20 when the x-coordinates are smaller than 5. Thus, the solution to $g(x) > 20$ is $\{x \mid x < 5\}$ or $(-\infty, 5)$.

e. The y-coordinates of the graph of $y = f(x)$ are below 60 when the x-coordinates are larger than -15. Thus, the solution to $g(x) \leq 60$ is $\{x \mid x \geq -15\}$ or $[-15, \infty)$.

f. The y-coordinates of the graph of $y = f(x)$ are between 0 and 60 when the x-coordinates are between -15 and 15. Thus, the solution to $0 < f(x) < 60$ is $\{x \mid -15 < x < 15\}$ or $(-15, 15)$.

39. a. $f(x) = g(x)$ when their graphs intersect. Thus, $x = -4$.

b. $f(x) \leq g(x)$ when the graph of f is above the graph of g . Thus, the solution is $\{x \mid x < -4\}$ or $(-\infty, -4)$.

40. a. $f(x) = g(x)$ when their graphs intersect. Thus, $x = 2$.

Chapter 2: Linear and Quadratic Functions

- b. $f(x) \leq g(x)$ when the graph of f is below or intersects the graph of g . Thus, the solution is $\{x \mid x \leq 2\}$ or $(-\infty, 2]$.
41. a. $f(x) = g(x)$ when their graphs intersect. Thus, $x = -6$.
- b. $g(x) \leq f(x) < h(x)$ when the graph of f is above or intersects the graph of g and below the graph of h . Thus, the solution is $\{x \mid -6 \leq x < 5\}$ or $[-6, 5)$.
42. a. $f(x) = g(x)$ when their graphs intersect. Thus, $x = 7$.
- b. $g(x) \leq f(x) < h(x)$ when the graph of f is above or intersects the graph of g and below the graph of h . Thus, the solution is $\{x \mid -4 \leq x < 7\}$ or $[-4, 7)$.
43. $C(x) = 0.25x + 35$
- a. $C(40) = 0.25(40) + 35 = \45 .
- b. Solve $C(x) = 0.25x + 35 = 80$
 $0.25x + 35 = 80$
 $0.25x = 45$
 $x = \frac{45}{0.25} = 180$ miles
- c. Solve $C(x) = 0.25x + 35 \leq 100$
 $0.25x + 35 \leq 100$
 $0.25x \leq 65$
 $x \leq \frac{65}{0.25} = 260$ miles
- d. The number of mile driven cannot be negative, so the implied domain for C is $\{x \mid x \geq 0\}$ or $[0, \infty)$.
44. $C(x) = 0.38x + 5$
- a. $C(50) = 0.38(50) + 5 = \$24$.
- b. Solve $C(x) = 0.38x + 5 = 29.32$
 $0.38x + 5 = 29.32$
 $0.38x = 24.32$
 $x = \frac{24.32}{0.38} = 64$ minutes
- c. Solve $C(x) = 0.38x + 5 \leq 60$
 $0.38x + 5 \leq 60$
 $0.38x \leq 55$
 $x \leq \frac{55}{0.38} \approx 144$ minutes
- d. The number of minutes cannot be negative, so $x \geq 0$. If there are 30 days in the month, then the number of minutes can be at most $30 \cdot 24 \cdot 60 = 43,200$. Thus, the implied domain for C is $\{x \mid 0 \leq x \leq 43,200\}$ or $[0, 43200]$.
45. $S(p) = -200 + 50p$; $D(p) = 1000 - 25p$
- a. Solve $S(p) = D(p)$.
 $-200 + 50p = 1000 - 25p$
 $75p = 1200$
 $p = \frac{1200}{75} = 16$
 $S(16) = -200 + 50(16) = 600$
 Thus, the equilibrium price is \$16, and the equilibrium quantity is 600 T-shirts.
- b. Solve $D(p) > S(p)$.
 $1000 - 25p > -200 + 50p$
 $1200 > 75p$
 $\frac{1200}{75} > p$
 $16 > p$
 The demand will exceed supply when the price is less than \$16 (but still greater than \$0). That is, $\$0 < p < \16 .
- c. The price will eventually be increased.
46. $S(p) = -2000 + 3000p$; $D(p) = 10000 - 1000p$
- a. Solve $S(p) = D(p)$.
 $-2000 + 3000p = 10000 - 1000p$
 $4000p = 12000$
 $p = \frac{12000}{4000} = 3$
 $S(3) = -2000 + 3000(3) = 7000$
 Thus, the equilibrium price is \$3, and the equilibrium quantity is 7000 hot dogs.

- b. Solve $D(p) < S(p)$.

$$10000 - 1000p < -2000 + 3000p$$

$$12000 < 4000p$$

$$\frac{12000}{4000} < p$$

$$3 < p$$

The demand will be less than the supply when the price is greater than \$3.

- c. The price will eventually be decreased.

47. a. We are told that the tax function T is for adjusted gross incomes x between \$8,350 and \$33,950, inclusive. Thus, the domain is $\{x \mid 8,350 \leq x \leq 33,950\}$ or $[8350, 33950]$.

- b. $T(20,000) = 0.15(20,000 - 8350) + 835$
 $= 2582.50$

If a single filer's adjusted gross income is \$20,000, then his or her tax bill will be \$2582.50.

- c. The independent variable is adjusted gross income, x . The dependent variable is the tax bill, T .

- d. Evaluate T at $x = 8350$, 20000, and 33950.

$$T(8350) = 0.15(8350 - 8350) + 835$$

$$= 835$$

$$T(20,000) = 0.15(20,000 - 8350) + 835$$

$$= 2582.50$$


$$T(33,950) = 0.15(33,950 - 8350) + 835$$

$$= 4675$$

Thus, the points $(8350, 835)$,

$(20000, 2582.5)$, and $(33950, 4675)$

are on the graph.



- e. We must solve $T(x) = 3707.50$.

$$0.15(x - 8350) + 835 = 3707.50$$

$$0.15x - 1252.5 + 835 = 3707.50$$

$$0.15x - 417.5 = 3707.50$$

$$0.15x = 4125$$

$$x = 27,500$$

A single filer with an adjusted gross income of \$27,500 will have a tax bill of \$3707.50.

48. a. The independent variable is payroll, p . The payroll tax only applies if the payroll exceeds \$136.5 million. Thus, the domain of T is $\{p \mid p > 136.5\}$ or $(136.5, \infty)$.

- b. $T(171.1) = 0.40(171.1 - 136.5) = 13.84$

The luxury tax for the New York Yankees was \$13.84 million.


- c. Evaluate T at $p = 136.5$, 171.1, and 300 million.

$$T(136.5) = 0.40(136.5 - 136.5) = 0 \text{ million}$$

$$T(171.1) = 0.40(171.1 - 136.5) = 13.84 \text{ million}$$

$$T(300) = 0.40(300 - 136.5) = 65.4 \text{ million}$$

Thus, the points $(136.5 \text{ million}, 0 \text{ million})$, $(171.1 \text{ million}, 13.84 \text{ million})$, and $(300 \text{ million}, 65.4 \text{ million})$ are on the graph.



- d. We must solve $T(p) = 11.7$.

$$0.40(p - 136.5) = 11.7$$

$$0.40p - 54.6 = 11.7$$

$$0.40p = 66.3$$

$$p = 165.75$$

If the luxury tax is \$11.7 million, then the payroll of the team is \$165.75 million.

Chapter 2: Linear and Quadratic Functions

49. $R(x) = 8x$; $C(x) = 4.5x + 17,500$

a. Solve $R(x) = C(x)$.

$$8x = 4.5x + 17,500$$

$$3.5x = 17,500$$

$$x = 5000$$

The break-even point occurs when the company sells 5000 units.

b. Solve $R(x) > C(x)$

$$8x > 4.5x + 17,500$$

$$3.5x > 17,500$$

$$x > 5000$$

The company makes a profit if it sells more than 5000 units.

50. $R(x) = 12x$; $C(x) = 10x + 15,000$

a. Solve $R(x) = C(x)$

$$12x = 10x + 15,000$$

$$2x = 15,000$$

$$x = 7500$$

The break-even point occurs when the company sells 7500 units.

b. Solve $R(x) > C(x)$

$$12x > 10x + 15,000$$

$$2x > 15,000$$

$$x > 7500$$

The company makes a profit if it sells more than 7500 units.

51. a. Consider the data points (x, y) , where x = the age in years of the computer and y = the value in dollars of the computer. So we have the points $(0, 3000)$ and $(3, 0)$. The slope formula yields:

$$m = \frac{\Delta y}{\Delta x} = \frac{0 - 3000}{3 - 0} = \frac{-3000}{3} = -1000$$


The y-intercept is $(0, 3000)$, so $b = 3000$.

Therefore, the linear function is

$$V(x) = mx + b = -1000x + 3000.$$

- b. The age of the computer cannot be negative, and the book value of the computer will be \$0 after 3 years. Thus, the implied domain for V is $\{x \mid 0 \leq x \leq 3\}$ or $[0, 3]$.

- c. The graph of $V(x) = -1000x + 3000$



d. $V(2) = -1000(2) + 3000 = 1000$

The computer's book value after 2 years will be \$1000.

e. Solve $V(x) = 2000$

$$-1000x + 3000 = 2000$$

$$-1000x = -1000$$

$$x = 1$$

The computer will have a book value of \$2000 after 1 year.

52. a. Consider the data points (x, y) , where x = the age in years of the machine and y = the value in dollars of the machine. So we have the points $(0, 120000)$ and $(10, 0)$. The slope formula yields:

$$m = \frac{\Delta y}{\Delta x} = \frac{0 - 120000}{10 - 0} = \frac{-120000}{10} = -12000$$

The y-intercept is $(0, 120000)$, so


$$b = 120,000.$$

Therefore, the linear function is

$$V(x) = mx + b = -12,000x + 120,000.$$

- b. The age of the machine cannot be negative, and the book value of the machine will be \$0 after 10 years. Thus, the implied domain for V is $\{x \mid 0 \leq x \leq 10\}$ or $[0, 10]$.

- c. The graph of $V(x) = -12,000x + 120,000$



Section 2.1: Properties of Linear Functions and Linear Models

d. $V(4) = -12000(4) + 120000 = 72000$

The machine's value after 4 years is given by \$72,000.

e. Solve $V(x) = 72000$.

$$-12000x + 120000 = 72000$$


$$-12000x = -48000$$

$$x = 4$$

The machine will be worth \$72,000 after 4 years.

53. a. Let x = the number of bicycles manufactured. We can use the cost function $C(x) = mx + b$, with $m = 90$ and $b = 1800$. Therefore $C(x) = 90x + 1800$

- b. The graph of $C(x) = 90x + 1800$



- c. The cost of manufacturing 14 bicycles is given by $C(14) = 90(14) + 1800 = \3060 .

- d. Solve $C(x) = 90x + 1800 = 3780$

$$90x + 1800 = 3780$$

$$90x = 1980$$

$$x = 22$$


So 22 bicycles could be manufactured for \$3780.

54. a. The new daily fixed cost is

$$1800 + \frac{100}{20} = \$1805$$

- b. Let x = the number of bicycles manufactured. We can use the cost function $C(x) = mx + b$, with $m = 90$ and $b = 1805$. Therefore $C(x) = 90x + 1805$

- c. The graph of $C(x) = 90x + 1805$



- d. The cost of manufacturing 14 bicycles is given by $C(14) = 90(14) + 1805 = \3065 .

- e. Solve $C(x) = 90x + 1805 = 3780$

$$90x + 1805 = 3780$$

$$90x = 1975$$

$$x \approx 21.94$$

So approximately 21 bicycles could be manufactured for \$3780.

55. a. Let x = number of miles driven, and let C = cost in dollars. Total cost = (cost per mile)(number of miles) + fixed cost $C(x) = 0.07x + 29$

- b. $C(110) = (0.07)(110) + 29 = \36.70


$$C(230) = (0.07)(230) + 29 = \$45.10$$

56. a. Let x = number of minutes used, and let C = cost in dollars. Total cost = (cost per minute)(number of minutes) + fixed cost $C(x) = 0.05x + 5$

- b. $C(105) = (0.05)(105) + 5 = \10.25

$$C(180) = (0.05)(180) + 5 = \$14$$

57. a.



Chapter 2: Linear and Quadratic Functions

b.

p	q	Avg. rate of change = $\frac{\Delta q}{\Delta p}$
150	100	
200	80	$\frac{80-100}{200-150} = \frac{-20}{50} = -0.4$
250	60	$\frac{60-80}{250-200} = \frac{-20}{50} = -0.4$
300	40	$\frac{40-60}{300-250} = \frac{-20}{50} = -0.4$

Since each input (price) corresponds to a single output (quantity demanded), we know that the quantity demanded is a function of price. Also, because the average rate of change is constant at $-\$0.4$ per MP3 player, the function is linear.


- c.** From part (b), we know $m = -0.4$. Using $(p_1, q_1) = (150, 100)$, we get the equation:

$$\begin{aligned} q - q_1 &= m(p - p_1) \\ q - 100 &= -0.4(p - 150) \\ q - 100 &= -0.4p + 60 \\ q &= -0.4p + 160 \end{aligned}$$

Using function notation, we have $q(p) = -0.4p + 160$.


- d.** The price cannot be negative, so $p \geq 0$. Likewise, the quantity cannot be negative, so, $q(p) \geq 0$.
 $-0.4p + 160 \geq 0$
 $-0.4p \geq -160$
 $p \leq 400$

Thus, the implied domain for $q(p)$ is $\{p \mid 0 \leq p \leq 400\}$ or $[0, 400]$.



- f.** If the price increases by \$1, then the quantity demanded of MP3 players decreases by 0.4 player.
- g.** p -intercept: If the price is \$0, then 160 MP3 players will be demanded.
 q -intercept: There will be 0 MP3 players demanded when the price is \$400.

58. a.



b.

s	h	Avg. rate of change = $\frac{\Delta h}{\Delta s}$
20	0	
15	3	$\frac{3-0}{15-20} = \frac{3}{-5} = -0.6$
10	6	$\frac{6-3}{10-15} = \frac{3}{-5} = -0.6$
5	9	$\frac{9-6}{5-10} = \frac{3}{-5} = -0.6$

Since each input (soda) corresponds to a single output (hot dogs), we know that number of hot dogs purchased is a function of number of sodas purchased. Also, because the average rate of change is constant at -0.6 hot dogs per soda, the function is linear.


- c.** From part (b), we know $m = -0.6$. Using $(s_1, h_1) = (20, 0)$, we get the equation:

$$\begin{aligned} h - h_1 &= m(s - s_1) \\ h - 0 &= -0.6(s - 20) \\ h &= -0.6s + 12 \end{aligned}$$

Using function notation, we have $h(s) = -0.6s + 12$.

- d.** The number of sodas cannot be negative, so $s \geq 0$. Likewise, the number of hot dogs cannot be negative, so, $h(s) \geq 0$.
 $-0.6s + 12 \geq 0$
 $-0.6s \geq -12$
 $s \leq 20$

Thus, the implied domain for $h(s)$ is $\{s \mid 0 \leq s \leq 20\}$ or $[0, 20]$.



- f. If the number of hot dogs purchased increases by \$1, then the number of sodas purchased decreases by 0.6.
- g. s -intercept: If 0 hot dogs are purchased, then 20 sodas can be purchased.
 h -intercept: If 0 sodas are purchased, then 12 hot dogs may be purchased.

59. The graph shown has a positive slope and a positive y -intercept. Therefore, the function from (d) and (e) might have the graph shown.

60. The graph shown has a negative slope and a positive y -intercept. Therefore, the function from (b) and (e) might have the graph shown.

61. A linear function $f(x) = mx + b$ will be odd provided $f(-x) = -f(x)$.

That is, provided $m(-x) + b = -(mx + b)$.

$$-mx + b = -mx - b$$

$$b = -b$$

$$2b = 0$$

$$b = 0$$

So a linear function $f(x) = mx + b$ will be odd provided $b = 0$.

A linear function $f(x) = mx + b$ will be even provided $f(-x) = f(x)$.

That is, provided $m(-x) + b = mx + b$.

$$-mx + b = mx + b$$

$$-mx = mx$$


$$0 = 2mx$$

$$m = 0$$

So, yes, a linear function $f(x) = mx + b$ can be even provided $m = 0$.

62. If you solve the linear function $f(x) = mx + b$ for 0 you are actually finding the x -intercept. Therefore using x -intercept of the graph of $f(x) = mx + b$ would be same x -value as solving $mx + b > 0$ for x . Then the appropriate interval could be determined.

Section 2.2



No, the relation is not a function because an input, 1, corresponds to two different outputs, 5 and 12.

2. Let $(x_1, y_1) = (1, 4)$ and $(x_2, y_2) = (3, 8)$.

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{8 - 4}{3 - 1} = \frac{4}{2} = 2$$

$$y - y_1 = m(x - x_1)$$

$$y - 4 = 2(x - 1)$$

$$y - 4 = 2x - 2$$

$$y = 2x + 2$$

3. scatter diagram

4. True

5. Linear relation, $m > 0$


6. Nonlinear relation

7. Linear relation, $m < 0$

8. Linear relation, $m > 0$

9. Nonlinear relation

10. Nonlinear relation



Chapter 2: Linear and Quadratic Functions

- b. Answers will vary. We select (4, 6) and (8, 14). The slope of the line containing these points is:

$$m = \frac{14-6}{8-4} = \frac{8}{4} = 2$$


The equation of the line is:

$$y - y_1 = m(x - x_1)$$



$$y - 6 = 2(x - 4)$$

$$y - 6 = 2x - 8$$

$$y = 2x - 2$$



- d. Using the LINear REGression program, the line of best fit is:
 $y = 2.0357x - 2.3571$



- b. Answers will vary. We select (5, 2) and (11, 9). The slope of the line containing these points is: $m = \frac{9-2}{11-5} = \frac{7}{6}$


The equation of the line is:

$$y - y_1 = m(x - x_1)$$



$$y - 2 = \frac{7}{6}(x - 5)$$

$$y - 2 = \frac{7}{6}x - \frac{35}{6}$$

$$y = \frac{7}{6}x - \frac{23}{6}$$



- d. Using the LINear REGression program, the line of best fit is:
 $y = 1.1286x - 3.8619$



- b. Answers will vary. We select (-2, -4) and (2, 5). The slope of the line containing these points is: $m = \frac{5-(-4)}{2-(-2)} = \frac{9}{4}$


The equation of the line is:

$$y - y_1 = m(x - x_1)$$



$$y - (-4) = \frac{9}{4}(x - (-2))$$

$$y + 4 = \frac{9}{4}x + \frac{9}{2}$$

$$y = \frac{9}{4}x + \frac{1}{2}$$




- e. Using the LINear REGression program, the line of best fit is:
 $y = 2.2x + 1.2$





- b. Answers will vary. We select $(-2, 7)$ and $(2, 0)$. The slope of the line containing these points is: $m = \frac{0-7}{2-(-2)} = \frac{-7}{4} = -\frac{7}{4}$.

The equation of the line is:

$$\begin{aligned} y - y_1 &= m(x - x_1) \\ y - 7 &= -\frac{7}{4}(x - (-2)) \\ y - 7 &= -\frac{7}{4}x - \frac{7}{2} \\ y &= -\frac{7}{4}x + \frac{7}{2} \end{aligned}$$



- d. Using the LINear REGression program, the line of best fit is:
 $y = -1.8x + 3.6$




- b. Answers will vary. We select $(-20, 100)$ and $(-10, 140)$. The slope of the line containing these points is:



$$m = \frac{140-100}{-10-(-20)} = \frac{40}{10} = 4$$

The equation of the line is:

$$\begin{aligned} y - y_1 &= m(x - x_1) \\ y - 100 &= 4(x - (-20)) \\ y - 100 &= 4x + 80 \\ y &= 4x + 180 \end{aligned}$$



- d. Using the LINear REGression program, the line of best fit is:
 $y = 3.8613x + 180.2920$



- b. Selection of points will vary. We select $(-30, 10)$ and $(-14, 18)$. The slope of the line containing these points is:

$$m = \frac{18-10}{-14-(-30)} = \frac{8}{16} = \frac{1}{2}$$

The equation of the line is:

Chapter 2: Linear and Quadratic Functions


$$y - y_1 = m(x - x_1)$$

$$y - 10 = \frac{1}{2}(x - (-30))$$

$$y - 10 = \frac{1}{2}x + 15$$

$$y = \frac{1}{2}x + 25$$


c.




d. Using the LINEAR REGression program, the line of best fit is:

$$y = 0.4421x + 23.4559$$

e.



17. a.



b. Linear.

c. Answers will vary. We will use the points (39.52, 210) and (66.45, 280).


$$m = \frac{280 - 210}{66.45 - 39.52} = \frac{70}{26.93} \approx 2.5993316$$

$$y - 210 = 2.5993316(x - 39.52)$$

$$y - 210 = 2.5993316x - 102.7255848$$

$$y = 2.599x + 107.274$$

d.




e. $x = 62.3$: $y = 2.599(62.3) + 107.274 \approx 269$

We predict that a candy bar weighing 62.3 grams will contain 269 calories.

f. If the weight of a candy bar is increased by one gram, then the number of calories will increase by 2.599.

18. a.



b. Linear with positive slope.

c. Answers will vary. We will use the points (42.3, 82) and (42.8, 93).

$$m = \frac{93 - 82}{42.8 - 42.3} = \frac{11}{0.5} = 22$$


$$N - N_1 = m(w - w_1)$$

$$N - 82 = 22(w - 42.3)$$

$$N - 82 = 22w - 930.6$$

$$N = 22w - 848.6$$

d.



Section 2.2: Building Linear Models from Data


e. $N(42.5) = 22(42.5) - 848.6 = 86.4$

We predict that approximately 86 raisins will be in a box weighing 42.5 grams.

- f. If the weight is increased by one gram, then the number of raisins will increase by 22.

19. a. The independent variable is the number of hours spent playing video games and cumulative grade-point average is the dependent variable because we are using number of hours playing video games to predict (or explain) cumulative grade-point average.

b.



- c. Using the LINear REGression program, the line of best fit is: $G(h) = -0.0942h + 3.2763$

- d. If the number of hours playing video games in a week increases by 1 hour, the cumulative grade-point average decreases 0.09, on average.

e. $G(8) = -0.0942(8) + 3.2763 = 2.52$

We predict a grade-point average of approximately 2.52 for a student who plays 8 hours of video games each week.

f. $2.40 = -0.0942(h) + 3.2763$


$$2.40 - 3.2763 = -0.0942h$$

$$-0.8763 = -0.0942h$$

$$9.3 = h$$

A student who has a grade-point average of 2.40 will have played approximately 9.3 hours of video games.

20. a.



- b. Using the LINear REGression program, the

line of best fit is: $C(H) = 0.3734H + 7.3268$

- c. If height increases by one inch, the head circumference increases by 0.3734 inch.

d. $C(26) = 0.3734(26) + 7.3268 \approx 17.0$ inches

- e. To find the height, we solve the following equation:

$$17.4 = 0.3734H + 7.3268$$


$$10.0732 = 0.3734H$$

$$26.98 \approx H$$

A child with a head circumference of 17.4 inches would have a height of about 26.98 inches.

21. a. The relation is not a function because 23 is paired with both 56 and 53.

b.



- c. Using the LINear REGression program, the line of best fit is: $D = -1.3355p + 86.1974$. The correlation coefficient is: $r \approx -0.9491$.

- d. If the price of the jeans increases by \$1, the demand for the jeans decreases by about 1.34 pairs per day.

e. $D(p) = -1.3355p + 86.1974$

f. Domain: $\{p \mid 0 < p \leq 64\}$

Note that the p -intercept is roughly 64.54 and that the number of pairs of jeans in demand cannot be negative.


g. $D(28) = -1.3355(28) + 86.1974 \approx 48.8034$

Demand is about 49 pairs.

Chapter 2: Linear and Quadratic Functions


22. a. The relation is not a function because 24 is paired with both 343 and 341.

b.



- c. Using the LINear REGression program, the line of best fit is: $S = 2.0667A + 292.8869$. The correlation coefficient is: $r \approx 0.9833$.
- d. As the advertising expenditure increases by \$1000, the sales increase by about \$2067.
- e. $S(A) = 2.0667A + 292.8869$
- f. Domain: $\{A \mid A \geq 0\}$
- g. $S(25) = 2.0667(25) + 292.8869 \approx 345$
Sales are about \$345 thousand.

23.



The data do not follow a linear pattern so it would not make sense to find the line of best fit.

24. Using the ordered pairs (1, 5) and (3, 8), the line of best fit is $y = 1.5x + 3.5$.

```
LinReg
y=ax+b
a=1.5
b=3.5
r=1
r=1
```

The correlation coefficient is $r = 1$. This is reasonable because two points determine a line.

25. A correlation coefficient of 0 implies that the data do not have a linear relationship.

26. The y-intercept would be the calories of a candy bar with weight 0 which would not be meaningful in this problem.

27. $G(0) = -0.0942(0) + 3.2763 = 3.2763$. The approximate grade-point average of a student who plays 0 hours of video games per week would be 3.28.

Section 2.3

1. a. $x^2 - 5x - 6 = (x - 6)(x + 1)$

b. $2x^2 - x - 3 = (2x - 3)(x + 1)$

2. $\sqrt{8^2 - 4 \cdot 2 \cdot 3} = \sqrt{64 - 24}$
 $= \sqrt{40} = \sqrt{4 \cdot 10} = 2\sqrt{10}$

3. $(x - 3)(3x + 5) = 0$
 $x - 3 = 0$ or $3x + 5 = 0$
 $x = 3$ or $3x = -5$

$$x = -\frac{5}{3}$$

The solution set is $\left\{-\frac{5}{3}, 3\right\}$.

4. add; $\left(\frac{1}{2} \cdot 6\right)^2 = 9$

5. If $f(4) = 10$, then the point (4, 10) is on the graph of f .

6. $f(-3) = (-3)^2 + 4(-3) + 3$
 $= 9 - 12 + 3 = 0$
 -3 is a zero of $f(x)$.

7. repeated; multiplicity 2

8. discriminant; negative

9. A quadratic functions can have either 0, 1 or 2 real zeros.

10. $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

Section 2.3: Quadratic Functions and Their Zeros

11. $f(x) = 0$

$$x^2 - 9x = 0$$

$$x(x-9) = 0$$

$$x = 0 \quad \text{or} \quad x - 9 = 0$$

$$x = 9$$

The zeros of $f(x) = x^2 - 9x$ are 0 and 9. The x -intercepts of the graph of f are 0 and 9.

12. $f(x) = 0$

$$x^2 + 4x = 0$$

$$x(x+4) = 0$$

$$x = 0 \quad \text{or} \quad x + 4 = 0$$

$$x = -4$$

The zeros of $f(x) = x^2 + 4x$ are -4 and 0. The x -intercepts of the graph of f are -4 and 0.

13. $g(x) = 0$

$$x^2 - 25 = 0$$

$$(x+5)(x-5) = 0$$

$$x + 5 = 0 \quad \text{or} \quad x - 5 = 0$$

$$x = -5 \quad x = 5$$

The zeros of $g(x) = x^2 - 25$ are -5 and 5. The x -intercepts of the graph of g are -5 and 5.

14. $G(x) = 0$

$$x^2 - 9 = 0$$

$$(x+3)(x-3) = 0$$

$$x + 3 = 0 \quad \text{or} \quad x - 3 = 0$$

$$x = -3 \quad x = 3$$

The zeros of $G(x) = x^2 - 9$ are -3 and 3. The x -intercepts of the graph of G are -3 and 3.

15. $F(x) = 0$

$$x^2 + x - 6 = 0$$

$$(x+3)(x-2) = 0$$

$$x + 3 = 0 \quad \text{or} \quad x - 2 = 0$$

$$x = -3 \quad x = 2$$

The zeros of $F(x) = x^2 + x - 6$ are -3 and 2.

The x -intercepts of the graph of F are -3 and 2.

16. $H(x) = 0$

$$x^2 + 7x + 6 = 0$$

$$(x+6)(x+1) = 0$$

$$x + 6 = 0 \quad \text{or} \quad x + 1 = 0$$

$$x = -6 \quad x = -1$$

The zeros of $H(x) = x^2 + 7x + 6$ are -6 and -1 .

The x -intercepts of the graph of H are -6 and -1 .

17. $g(x) = 0$

$$2x^2 - 5x - 3 = 0$$

$$(2x+1)(x-3) = 0$$

$$2x + 1 = 0 \quad \text{or} \quad x - 3 = 0$$

$$x = -\frac{1}{2} \quad x = 3$$

The zeros of $g(x) = 2x^2 - 5x - 3$ are $-\frac{1}{2}$ and 3.

The x -intercepts of the graph of g are $-\frac{1}{2}$ and 3.

18. $f(x) = 0$

$$3x^2 + 5x + 2 = 0$$

$$(3x+2)(x+1) = 0$$

$$3x + 2 = 0 \quad \text{or} \quad x + 1 = 0$$

$$x = -\frac{2}{3} \quad x = -1$$

The zeros of $f(x) = 3x^2 + 5x + 2$ are -1 and

$-\frac{2}{3}$. The x -intercepts of the graph of f are -1

and $-\frac{2}{3}$.

19. $P(x) = 0$

$$3x^2 - 48 = 0$$

$$3(x^2 - 16) = 0$$

$$3(x+4)(x-4) = 0$$

$$t + 4 = 0 \quad \text{or} \quad t - 4 = 0$$

$$t = -4 \quad t = 4$$

The zeros of $P(x) = 3x^2 - 48$ are -4 and 4.

The x -intercepts of the graph of P are -4 and 4.

Chapter 2: Linear and Quadratic Functions

20. $H(x) = 0$

$$2x^2 - 50 = 0$$

$$2(x^2 - 25) = 0$$

$$2(x+5)(x-5) = 0$$

$$y+5=0 \quad \text{or} \quad y-5=0$$

$$y = -5 \quad y = 5$$

The zeros of $H(x) = 2x^2 - 50$ are -5 and 5 .

The x -intercepts of the graph of H are -5 and 5 .

21. $g(x) = 0$

$$x(x+8)+12=0$$

$$x^2 + 8x + 12 = 0$$

$$(x+6)(x+2) = 0$$

$$x = -6 \quad \text{or} \quad x = -2$$

The zeros of $g(x) = x(x+8)+12$ are -6 and -2 .

The x -intercepts of the graph of g are -6 and -2 .

22. $f(x) = 0$

$$x(x-4)-12=0$$

$$x^2 - 4x - 12 = 0$$

$$(x-6)(x+2) = 0$$

$$x = -2 \quad \text{or} \quad x = 6$$

The zeros of $f(x) = x(x-4)-12$ are -2 and 6 .

The x -intercepts of the graph of f are -2 and 6 .

23. $G(x) = 0$

$$4x^2 + 9 - 12x = 0$$

$$4x^2 - 12x + 9 = 0$$

$$(2x-3)(2x-3) = 0$$

$$2x-3=0 \quad \text{or} \quad 2x-3=0$$

$$x = \frac{3}{2} \quad x = \frac{3}{2}$$

The only zero of $G(x) = 4x^2 + 9 - 12x$ is $\frac{3}{2}$.

The only x -intercept of the graph of G is $\frac{3}{2}$.

24. $F(x) = 0$

$$25x^2 + 16 - 40x = 0$$

$$25x^2 - 40x + 16 = 0$$

$$(5x-4)(5x-4) = 0$$

$$5x-4=0 \quad \text{or} \quad 5x-4=0$$

$$x = \frac{4}{5} \quad x = \frac{4}{5}$$

The only zero of $F(x) = 25x^2 + 16 - 40x$ is $\frac{4}{5}$.

The only x -intercept of the graph of F is $\frac{4}{5}$.

25. $f(x) = 0$

$$x^2 - 8 = 0$$

$$x^2 = 8$$

$$x = \pm\sqrt{8} = \pm 2\sqrt{2}$$

The zeros of $f(x) = x^2 - 8$ are $-2\sqrt{2}$ and $2\sqrt{2}$.

The x -intercepts of the graph of f are $-2\sqrt{2}$ and $2\sqrt{2}$.

26. $g(x) = 0$

$$x^2 - 18 = 0$$

$$x^2 = 18$$

$$x = \pm\sqrt{18} = \pm 3\sqrt{3}$$

The zeros of $g(x) = x^2 - 18$ are $-3\sqrt{3}$ and

$3\sqrt{3}$. The x -intercepts of the graph of g are $-3\sqrt{3}$ and $3\sqrt{3}$.

27. $g(x) = 0$

$$(x-1)^2 - 4 = 0$$

$$(x-1)^2 = 4$$

$$x-1 = \pm\sqrt{4}$$

$$x-1 = \pm 2$$

$$x-1=2 \quad \text{or} \quad x-1=-2$$

$$x=3 \quad x=-1$$

The zeros of $g(x) = (x-1)^2 - 4$ are -1 and 3 .

The x -intercepts of the graph of g are -1 and 3 .

Section 2.3: Quadratic Functions and Their Zeros

28. $G(x) = 0$

$$(x+2)^2 - 1 = 0$$

$$(x+2)^2 = 1$$

$$x+2 = \pm\sqrt{1}$$

$$x+2 = \pm 1$$

$$x+2 = 1 \quad \text{or} \quad x+2 = -1$$

$$x = -1 \quad x = -3$$

The zeros of $G(x) = (x+2)^2 - 1$ are -3 and -1 .

The x -intercepts of the graph of G are -3 and -1 .

29. $F(x) = 0$

$$(2x+3)^2 - 32 = 0$$

$$(2x+3)^2 = 32$$

$$2x+3 = \pm\sqrt{32}$$

$$2x+3 = \pm 4\sqrt{2}$$

$$2x = -3 \pm 4\sqrt{2}$$

$$x = \frac{-3 \pm 4\sqrt{2}}{2}$$

The zeros of $F(x) = (2x+3)^2 - 32$ are

$\frac{-3+4\sqrt{2}}{2}$ and $\frac{-3-4\sqrt{2}}{2}$. The x -intercepts of

the graph of F are $\frac{-3+4\sqrt{2}}{2}$ and $\frac{-3-4\sqrt{2}}{2}$.

30. $F(x) = 0$

$$(3x-2)^2 - 75 = 0$$

$$(3x-2)^2 = 75$$

$$3x-2 = \pm\sqrt{75}$$

$$3x-2 = \pm 5\sqrt{3}$$

$$3x = 2 \pm 5\sqrt{3}$$

$$x = \frac{2 \pm 5\sqrt{3}}{3}$$

The zeros of $G(x) = (3x-2)^2 - 75$ are $\frac{2+5\sqrt{3}}{3}$

and $\frac{2-5\sqrt{3}}{3}$. The x -intercepts of the graph of G

are $\frac{2-5\sqrt{3}}{3}$ and $\frac{2+5\sqrt{3}}{3}$.

31. $f(x) = 0$

$$x^2 + 4x - 8 = 0$$

$$x^2 + 4x = 8$$

$$x^2 + 4x + 4 = 8 + 4$$

$$(x+2)^2 = 12$$

$$x+2 = \pm\sqrt{12}$$

$$x+2 = \pm 2\sqrt{3}$$

$$x = -2 \pm 2\sqrt{3}$$

$$x = -2 + 2\sqrt{3} \quad \text{or} \quad x = -2 - 2\sqrt{3}$$

The zeros of $f(x) = x^2 + 4x - 8$ are $-2 + 2\sqrt{3}$

and $-2 - 2\sqrt{3}$. The x -intercepts of the graph of f are $-2 + 2\sqrt{3}$ and $-2 - 2\sqrt{3}$.

32. $f(x) = 0$

$$x^2 - 6x - 9 = 0$$

$$x^2 - 6x + 9 = 9 + 9$$

$$(x-3)^2 = 18$$

$$x-3 = \pm\sqrt{18}$$

$$x = 3 \pm 3\sqrt{2}$$

The zeros of $f(x) = x^2 - 6x - 9$ are $3 - 3\sqrt{2}$

and $3 + 3\sqrt{2}$. The x -intercepts of the graph of f are $3 - 3\sqrt{2}$ and $3 + 3\sqrt{2}$.

33. $g(x) = 0$

$$x^2 - \frac{1}{2}x - \frac{3}{16} = 0$$

$$x^2 - \frac{1}{2}x = \frac{3}{16}$$

$$x^2 - \frac{1}{2}x + \frac{1}{16} = \frac{3}{16} + \frac{1}{16}$$

$$\left(x - \frac{1}{4}\right)^2 = \frac{1}{4}$$

$$x - \frac{1}{4} = \pm\sqrt{\frac{1}{4}} = \pm\frac{1}{2}$$

$$x = \frac{1}{4} \pm \frac{1}{2}$$

$$x = \frac{3}{4} \quad \text{or} \quad x = -\frac{1}{4}$$

The zeros of $g(x) = x^2 - \frac{1}{2}x - \frac{3}{16}$ are $-\frac{1}{4}$ and $\frac{3}{4}$.

The x -intercepts of the graph of g are $-\frac{1}{4}$ and $\frac{3}{4}$.

Chapter 2: Linear and Quadratic Functions

34. $g(x) = 0$

$$x^2 + \frac{2}{3}x - \frac{1}{3} = 0$$

$$x^2 + \frac{2}{3}x = \frac{1}{3}$$

$$x^2 + \frac{2}{3}x + \frac{1}{9} = \frac{1}{3} + \frac{1}{9}$$

$$\left(x + \frac{1}{3}\right)^2 = \frac{4}{9}$$

$$x + \frac{1}{3} = \pm\sqrt{\frac{4}{9}} = \pm\frac{2}{3}$$

$$x = -\frac{1}{3} \pm \frac{2}{3}$$

$$x = \frac{1}{3} \text{ or } x = -1$$

The zeros of $g(x) = x^2 + \frac{2}{3}x - \frac{1}{3}$ are -1 and $\frac{1}{3}$.

The x -intercepts of the graph of g are -1 and $\frac{1}{3}$.

35. $F(x) = 0$

$$3x^2 + x - \frac{1}{2} = 0$$

$$x^2 + \frac{1}{3}x - \frac{1}{6} = 0$$

$$x^2 + \frac{1}{3}x = \frac{1}{6}$$

$$x^2 + \frac{1}{3}x + \frac{1}{36} = \frac{1}{6} + \frac{1}{36}$$

$$\left(x + \frac{1}{6}\right)^2 = \frac{7}{36}$$

$$x + \frac{1}{6} = \pm\sqrt{\frac{7}{36}} = \pm\frac{\sqrt{7}}{6}$$

$$x = \frac{-1 \pm \sqrt{7}}{6}$$

The zeros of $F(x) = 3x^2 + x - \frac{1}{2}$ are $\frac{-1-\sqrt{7}}{6}$ and

$\frac{-1+\sqrt{7}}{6}$. The x -intercepts of the graph of F are

$\frac{-1-\sqrt{7}}{6}$ and $\frac{-1+\sqrt{7}}{6}$.

36. $G(x) = 0$

$$2x^2 - 3x - 1 = 0$$

$$x^2 - \frac{3}{2}x - \frac{1}{2} = 0$$

$$x^2 - \frac{3}{2}x = \frac{1}{2}$$

$$x^2 - \frac{3}{2}x + \frac{9}{16} = \frac{1}{2} + \frac{9}{16}$$

$$\left(x - \frac{3}{4}\right)^2 = \frac{17}{16}$$

$$x - \frac{3}{4} = \pm\sqrt{\frac{17}{16}} = \pm\frac{\sqrt{17}}{4}$$

$$x = \frac{3 \pm \sqrt{17}}{4}$$

The zeros of $G(x) = 2x^2 - 3x - 1$ are $\frac{3-\sqrt{17}}{4}$ and

$\frac{3+\sqrt{17}}{4}$. The x -intercepts of the graph of G are

$\frac{3-\sqrt{17}}{4}$ and $\frac{3+\sqrt{17}}{4}$.

37. $f(x) = 0$

$$x^2 - 4x + 2 = 0$$

$$a = 1, \quad b = -4, \quad c = 2$$

$$x = \frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(2)}}{2(1)} = \frac{4 \pm \sqrt{16-8}}{2}$$

$$= \frac{4 \pm \sqrt{8}}{2} = \frac{4 \pm 2\sqrt{2}}{2} = 2 \pm \sqrt{2}$$

The zeros of $f(x) = x^2 - 4x + 2$ are $2 - \sqrt{2}$ and

$2 + \sqrt{2}$. The x -intercepts of the graph of f are

$2 - \sqrt{2}$ and $2 + \sqrt{2}$.

38. $f(x) = 0$

$$x^2 + 4x + 2 = 0$$

$$a = 1, \quad b = 4, \quad c = 2$$

$$x = \frac{-4 \pm \sqrt{4^2 - 4(1)(2)}}{2(1)} = \frac{-4 \pm \sqrt{16-8}}{2}$$

$$= \frac{-4 \pm \sqrt{8}}{2} = \frac{-4 \pm 2\sqrt{2}}{2} = -2 \pm \sqrt{2}$$

The zeros of $f(x) = x^2 + 4x + 2$ are $-2 - \sqrt{2}$

and $-2 + \sqrt{2}$. The x -intercepts of the graph of f

are $-2 - \sqrt{2}$ and $-2 + \sqrt{2}$.

39. $g(x) = 0$

$$x^2 - 4x - 1 = 0$$

$$a = 1, \quad b = -4, \quad c = -1$$

$$x = \frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(-1)}}{2(1)} = \frac{4 \pm \sqrt{16+4}}{2}$$

$$= \frac{4 \pm \sqrt{20}}{2} = \frac{4 \pm 2\sqrt{5}}{2} = 2 \pm \sqrt{5}$$

The zeros of $g(x) = x^2 - 4x - 1$ are $2 - \sqrt{5}$ and $2 + \sqrt{5}$. The x -intercepts of the graph of g are $2 - \sqrt{5}$ and $2 + \sqrt{5}$.

40. $g(x) = 0$

$$x^2 + 6x + 1 = 0$$

$$a = 1, \quad b = 6, \quad c = 1$$

$$x = \frac{-6 \pm \sqrt{6^2 - 4(1)(1)}}{2(1)} = \frac{-6 \pm \sqrt{36-4}}{2}$$

$$= \frac{-6 \pm \sqrt{32}}{2} = \frac{-6 \pm 4\sqrt{2}}{2} = -3 \pm 2\sqrt{2}$$

The zeros of $g(x) = x^2 + 6x + 1$ are $-3 - 2\sqrt{2}$ and $-3 + 2\sqrt{2}$. The x -intercepts of the graph of g are $-3 - 2\sqrt{2}$ and $-3 + 2\sqrt{2}$.

41. $F(x) = 0$

$$2x^2 - 5x + 3 = 0$$

$$a = 2, \quad b = -5, \quad c = 3$$

$$x = \frac{-(-5) \pm \sqrt{(-5)^2 - 4(2)(3)}}{2(2)} = \frac{5 \pm \sqrt{25-24}}{4}$$

$$= \frac{5 \pm 1}{4} = \frac{3}{4} \text{ or } 1$$

The zeros of $F(x) = 2x^2 - 5x + 3$ are 1 and $\frac{3}{4}$.

The x -intercepts of the graph of F are 1 and $\frac{3}{4}$.

42. $g(x) = 0$

$$2x^2 + 5x + 3 = 0$$

$$a = 2, \quad b = 5, \quad c = 3$$

$$x = \frac{-5 \pm \sqrt{5^2 - 4(2)(3)}}{2(2)} = \frac{-5 \pm \sqrt{25-24}}{4}$$

$$= \frac{-5 \pm 1}{4} = -1 \text{ or } -\frac{3}{2}$$

The zeros of $g(x) = 2x^2 + 5x + 3$ are $-\frac{3}{2}$ and -1 .

The x -intercepts of the graph of g are $-\frac{3}{2}$ and -1 .

43. $P(x) = 0$

$$4x^2 - x + 2 = 0$$

$$a = 4, \quad b = -1, \quad c = 2$$

$$x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(4)(2)}}{2(4)} = \frac{1 \pm \sqrt{1-32}}{8}$$

$$= \frac{1 \pm \sqrt{-31}}{8} = \text{not real}$$

The function $P(x) = 4x^2 - x + 2$ has no real zeros, and the graph of P has no x -intercepts.

44. $H(x) = 0$

$$4x^2 + x + 1 = 0$$

$$a = 4, \quad b = 1, \quad c = 1$$

$$x = \frac{-1 \pm \sqrt{1^2 - 4(4)(1)}}{2(4)} = \frac{-1 \pm \sqrt{1-16}}{8}$$

$$= \frac{-1 \pm \sqrt{-15}}{8} = \text{not real}$$

The function $H(x) = 4x^2 + x + 1$ has no real zeros, and the graph of H has no x -intercepts.

45. $f(x) = 0$

$$4x^2 - 1 + 2x = 0$$

$$4x^2 + 2x - 1 = 0$$

$$a = 4, \quad b = 2, \quad c = -1$$

$$x = \frac{-2 \pm \sqrt{2^2 - 4(4)(-1)}}{2(4)} = \frac{-2 \pm \sqrt{4+16}}{8}$$

$$= \frac{-2 \pm \sqrt{20}}{8} = \frac{-2 \pm 2\sqrt{5}}{8} = \frac{-1 \pm \sqrt{5}}{4}$$

The zeros of $f(x) = 4x^2 - 1 + 2x$ are $\frac{-1 - \sqrt{5}}{4}$

and $\frac{-1 + \sqrt{5}}{4}$. The x -intercepts of the graph of f

are $\frac{-1 - \sqrt{5}}{4}$ and $\frac{-1 + \sqrt{5}}{4}$.

46. $f(x) = 0$

$$2x^2 - 1 + 2x = 0$$

$$2x^2 + 2x - 1 = 0$$

$$a = 2, \quad b = 2, \quad c = -1$$

$$x = \frac{-2 \pm \sqrt{2^2 - 4(2)(-1)}}{2(2)} = \frac{-2 \pm \sqrt{4+8}}{4}$$

$$= \frac{-2 \pm \sqrt{12}}{4} = \frac{-2 \pm 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2}$$

Chapter 2: Linear and Quadratic Functions

The zeros of $f(x) = 2x^2 - 1 + 2x$ are $\frac{-1-\sqrt{3}}{2}$ and $\frac{-1+\sqrt{3}}{2}$. The x -intercepts of the graph of f are $\frac{-1-\sqrt{3}}{2}$ and $\frac{-1+\sqrt{3}}{2}$.

47. $G(x) = 0$

$$2x(x+2) - 3 = 0$$

$$2x^2 + 4x - 3 = 0$$

$$a = 2, \quad b = 4, \quad c = -3$$

$$x = \frac{-(4) \pm \sqrt{(4)^2 - 4(2)(-3)}}{2(2)} = \frac{-4 \pm \sqrt{16 + 24}}{4}$$

$$= \frac{-4 \pm \sqrt{40}}{4} = \frac{-4 \pm 2\sqrt{10}}{4} = \frac{-2 \pm \sqrt{10}}{2}$$

The zeros of $G(x) = 2x(x+2) - 3$ are $\frac{-2+\sqrt{10}}{2}$ and $\frac{-2-\sqrt{10}}{2}$. The x -intercepts of the graph of G are $\frac{-2+\sqrt{10}}{2}$ and $\frac{-2-\sqrt{10}}{2}$.

48. $F(x) = 0$

$$3x(x+2) - 1 = 0 \Rightarrow 3x^2 + 6x - 1 = 0$$

$$a = 3, \quad b = 6, \quad c = -1$$

$$x = \frac{-(6) \pm \sqrt{(6)^2 - 4(3)(-1)}}{2(3)} = \frac{-6 \pm \sqrt{36 + 12}}{6}$$

$$= \frac{-6 \pm \sqrt{48}}{6} = \frac{-6 \pm 4\sqrt{3}}{6} = \frac{-3 \pm 2\sqrt{3}}{3}$$

The zeros of $F(x) = 3x(x+2) - 1$ are $\frac{-3+2\sqrt{3}}{3}$ and $\frac{-3-2\sqrt{3}}{3}$. The x -intercepts of the graph of G are $\frac{-3+2\sqrt{3}}{3}$ and $\frac{-3-2\sqrt{3}}{3}$.

49. $p(x) = 0$

$$9x^2 - 6x + 1 = 0$$

$$a = 9, \quad b = -6, \quad c = 1$$

$$x = \frac{-(-6) \pm \sqrt{(-6)^2 - 4(9)(1)}}{2(9)} = \frac{6 \pm \sqrt{36 - 36}}{18}$$

$$= \frac{6 \pm 0}{18} = \frac{1}{3}$$

The only real zero of $p(x) = 9x^2 - 6x + 1$ is $\frac{1}{3}$.

The only x -intercept of the graph of g is $\frac{1}{3}$.

50. $q(x) = 0$

$$4x^2 + 20x + 25 = 0$$

$$a = 4, \quad b = 20, \quad c = 25$$

$$x = \frac{-20 \pm \sqrt{(20)^2 - 4(4)(25)}}{2(4)} = \frac{-20 \pm \sqrt{400 - 400}}{8}$$

$$= \frac{-20 \pm 0}{8} = -\frac{20}{8} = -\frac{5}{2}$$

The only real zero of $q(x) = 4x^2 + 20x + 25$ is $-\frac{5}{2}$. The only x -intercept of the graph of F is $-\frac{5}{2}$.

51. $f(x) = g(x)$

$$x^2 + 6x + 3 = 3$$

$$x^2 + 6x = 0 \Rightarrow x(x+6) = 0$$

$$x = 0 \quad \text{or} \quad x + 6 = 0$$

$$x = -6$$

The x -coordinates of the points of intersection are -6 and 0 . The y -coordinates are $g(-6) = 3$ and $g(0) = 3$. The graphs of the f and g intersect at the points $(-6, 3)$ and $(0, 3)$.

52. $f(x) = g(x)$

$$x^2 - 4x + 3 = 3$$

$$x^2 - 4x = 0$$

$$x(x-4) = 0$$

$$x = 0 \quad \text{or} \quad x - 4 = 0$$

$$x = 4$$

The x -coordinates of the points of intersection are 0 and 4 . The y -coordinates are $g(0) = 3$ and $g(4) = 3$. The graphs of the f and g intersect at the points $(0, 3)$ and $(4, 3)$.

53. $f(x) = g(x)$

$$-2x^2 + 1 = 3x + 2$$

$$0 = 2x^2 + 3x + 1$$

$$0 = (2x+1)(x+1)$$

$$2x+1 = 0 \quad \text{or} \quad x+1 = 0$$

$$x = -\frac{1}{2} \quad \text{or} \quad x = -1$$

The x -coordinates of the points of intersection are -1 and $-\frac{1}{2}$. The y -coordinates are $g(-1) = 3(-1) + 2 = -3 + 2 = -1$ and

Section 2.3: Quadratic Functions and Their Zeros

$$g\left(-\frac{1}{2}\right) = 3\left(-\frac{1}{2}\right) + 2 = -\frac{3}{2} + 2 = \frac{1}{2}.$$

The graphs of the f and g intersect at the points $(-1, -1)$ and $\left(-\frac{1}{2}, \frac{1}{2}\right)$.

$$\begin{aligned} 54. \quad f(x) &= g(x) \\ 3x^2 - 7 &= 10x + 1 \\ 3x^2 - 10x - 8 &= 0 \\ (3x + 2)(x - 4) &= 0 \\ 3x + 2 = 0 \quad \text{or} \quad x - 4 &= 0 \\ x = -\frac{2}{3} \quad x &= 4 \end{aligned}$$

The x -coordinates of the points of intersection are $-\frac{2}{3}$ and 4. The y -coordinates are

$$g\left(-\frac{2}{3}\right) = 10\left(-\frac{2}{3}\right) + 1 = -\frac{20}{3} + 1 = -\frac{17}{3} \quad \text{and}$$

$$g(4) = 10(4) + 1 = 40 + 1 = 41.$$

The graphs of the f and g intersect at the points $\left(-\frac{2}{3}, -\frac{17}{3}\right)$ and $(4, 41)$.

$$\begin{aligned} 55. \quad f(x) &= g(x) \\ x^2 - x + 1 &= 2x^2 - 3x - 14 \\ 0 &= x^2 - 2x - 15 \\ 0 &= (x + 3)(x - 5) \\ x + 3 = 0 \quad \text{or} \quad x - 5 &= 0 \\ x = -3 \quad x &= 5 \end{aligned}$$

The x -coordinates of the points of intersection are -3 and 5. The y -coordinates are

$$f(-3) = (-3)^2 - (-3) + 1 = 9 + 3 + 1 = 13 \quad \text{and}$$

$$f(5) = 5^2 - 5 + 1 = 25 - 5 + 1 = 21.$$

The graphs of the f and g intersect at the points $(-3, 13)$ and $(5, 21)$.

$$\begin{aligned} 56. \quad f(x) &= g(x) \\ x^2 + 5x - 3 &= 2x^2 + 7x - 27 \\ 0 &= x^2 + 2x - 24 \\ 0 &= (x + 6)(x - 4) \\ x + 6 = 0 \quad \text{or} \quad x - 4 &= 0 \\ x = -6 \quad x &= 4 \end{aligned}$$

The x -coordinates of the points of intersection are -6 and 4. The y -coordinates are

$$f(-6) = (-6)^2 + 5(-6) - 3 = 36 - 30 - 3 = 3 \quad \text{and}$$

$$f(4) = 4^2 + 5(4) - 3 = 16 + 20 - 3 = 33.$$

The graphs of the f and g intersect at the points $(-6, 3)$ and $(4, 33)$.

$$\begin{aligned} 57. \quad P(x) &= 0 \\ x^4 - 6x^2 - 16 &= 0 \\ (x^2 + 2)(x^2 - 8) &= 0 \\ x^2 + 2 = 0 \quad \text{or} \quad x^2 - 8 &= 0 \\ x^2 = -2 \quad x^2 &= 8 \\ x = \pm\sqrt{-2} \quad x &= \pm\sqrt{8} \\ &= \text{not real} \quad = \pm 2\sqrt{2} \end{aligned}$$

The zeros of $P(x) = x^4 - 6x^2 - 16$ are $-2\sqrt{2}$ and $2\sqrt{2}$. The x -intercepts of the graph of P are $-2\sqrt{2}$ and $2\sqrt{2}$.

$$\begin{aligned} 58. \quad H(x) &= 0 \\ x^4 - 3x^2 - 4 &= 0 \\ (x^2 + 1)(x^2 - 4) &= 0 \\ x^2 + 1 = 0 \quad \text{or} \quad x^2 - 4 &= 0 \\ x^2 = -1 \quad x^2 &= 4 \\ x = \pm\sqrt{-1} \quad x &= \pm\sqrt{4} \\ &= \text{not real} \quad = \pm 2 \end{aligned}$$

The zeros of $H(x) = x^4 - 3x^2 - 4$ are -2 and 2 . The x -intercepts of the graph of H are -2 and 2 .

$$\begin{aligned} 59. \quad f(x) &= 0 \\ x^4 - 5x^2 + 4 &= 0 \\ (x^2 - 4)(x^2 - 1) &= 0 \\ x^2 - 4 = 0 \quad \text{or} \quad x^2 - 1 &= 0 \\ x = \pm 2 \quad \text{or} \quad x &= \pm 1 \end{aligned}$$

The zeros of $f(x) = x^4 - 5x^2 + 4$ are -2 , -1 , 1 , and 2 . The x -intercepts of the graph of f are -2 , -1 , 1 , and 2 .

Chapter 2: Linear and Quadratic Functions

60. $f(x) = 0$

$$x^4 - 10x^2 + 24 = 0$$

$$(x^2 - 4)(x^2 - 6) = 0$$

$$x^2 - 4 = 0 \quad \text{or} \quad x^2 - 6 = 0$$

$$x^2 = 4 \quad x^2 = 6$$

$$x = \pm 2 \quad x = \pm\sqrt{6}$$

The zeros of $f(x) = x^4 - 10x^2 + 24$ are $-\sqrt{6}$, $\sqrt{6}$, 2 and -2 . The x -intercepts of the graph of f are $-\sqrt{6}$, $\sqrt{6}$, 2 and -2 .

61. $G(x) = 0$

$$3x^4 - 2x^2 - 1 = 0$$

$$(3x^2 + 1)(x^2 - 1) = 0$$

$$3x^2 + 1 = 0 \quad \text{or} \quad x^2 - 1 = 0$$

$$x^2 = -\frac{1}{3} \quad x^2 = 1$$

$$x = \pm\sqrt{-\frac{1}{3}} \quad x = \pm\sqrt{1}$$

$$x = \text{not real} \quad = \pm 1$$

The zeros of $G(x) = 3x^4 - 2x^2 - 1$ are -1 and 1 .
The x -intercepts of the graph of G are -1 and 1 .

62. $F(x) = 0$

$$2x^4 - 5x^2 - 12 = 0$$

$$(2x^2 + 3)(x^2 - 4) = 0$$

$$2x^2 + 3 = 0 \quad \text{or} \quad x^2 - 4 = 0$$

$$x^2 = -\frac{3}{2} \quad x^2 = 4$$

$$x = \pm\sqrt{-\frac{3}{2}} \quad x = \pm\sqrt{4}$$

$$= \text{not real} \quad = \pm 2$$

The zeros of $F(x) = 2x^4 - 5x^2 - 12$ are -2 and 2 .
The x -intercepts of the graph of F are -2 and 2 .

63. $g(x) = 0$

$$x^6 + 7x^3 - 8 = 0$$

$$(x^3 + 8)(x^3 - 1) = 0$$

$$x^3 + 8 = 0 \quad \text{or} \quad x^3 - 1 = 0$$

$$x^3 = -8 \quad x^3 = 1$$

$$x = -2 \quad x = 1$$

The zeros of $g(x) = x^6 + 7x^3 - 8$ are -2 and 1 .

The x -intercepts of the graph of g are -2 and 1 .

64. $g(x) = 0$

$$x^6 - 7x^3 - 8 = 0$$

$$(x^3 - 8)(x^3 + 1) = 0$$

$$x^3 - 8 = 0 \quad \text{or} \quad x^3 + 1 = 0$$

$$x^3 = 8 \quad x^3 = -1$$

$$x = 2 \quad x = -1$$

The zeros of $g(x) = x^6 - 7x^3 - 8$ are -1 and 2 .

The x -intercepts of the graph of g are -1 and 2 .

65. $G(x) = 0$

$$(x+2)^2 + 7(x+2) + 12 = 0$$

$$\text{Let } u = x+2 \rightarrow u^2 = (x+2)^2$$

$$u^2 + 7u + 12 = 0$$

$$(u+3)(u+4) = 0$$

$$u+3 = 0 \quad \text{or} \quad u+4 = 0$$

$$u = -3 \quad u = -4$$

$$x+2 = -3 \quad x+2 = -4$$

$$x = -5 \quad x = -6$$

The zeros of $G(x) = (x+2)^2 + 7(x+2) + 12$ are -6 and -5 . The x -intercepts of the graph of G are -6 and -5 .

66. $f(x) = 0$

$$(2x+5)^2 - (2x+5) - 6 = 0$$

$$\text{Let } u = 2x+5 \rightarrow u^2 = (2x+5)^2$$

$$u^2 - u - 6 = 0$$

$$(u-3)(u+2) = 0$$

$$u-3 = 0 \quad \text{or} \quad u+2 = 0$$

$$u = 3 \quad u = -2$$

$$2x+5 = 3 \quad 2x+5 = -2$$

$$x = -1 \quad x = -\frac{7}{2}$$

The zeros of $f(x) = (2x+5)^2 - (2x+5) - 6$ are $-\frac{7}{2}$ and -1 . The x -intercepts of the graph of f are $-\frac{7}{2}$ and -1 .

Section 2.3: Quadratic Functions and Their Zeros

67. $f(x) = 0$

$$(3x+4)^2 - 6(3x+4) + 9 = 0$$

$$\text{Let } u = 3x+4 \rightarrow u^2 = (3x+4)^2$$

$$u^2 - 6u + 9 = 0$$

$$(u-3)^2 = 0$$

$$u - 3 = 0$$

$$u = 3$$

$$3x+4 = 3$$

$$x = -\frac{1}{3}$$

The only zero of $f(x) = (3x+4)^2 - 6(3x+4) + 9$

is $-\frac{1}{3}$. The x -intercept of the graph of f is $-\frac{1}{3}$.

68. $H(x) = 0$

$$(2-x)^2 + (2-x) - 20 = 0$$

$$\text{Let } u = 2-x \rightarrow u^2 = (2-x)^2$$

$$u^2 + u - 20 = 0$$

$$(u+5)(u-4) = 0$$

$$u+5 = 0 \quad \text{or} \quad u-4 = 0$$

$$u = -5 \quad u = 4$$

$$2-x = -5 \quad 2-x = 4$$

$$x = 7 \quad x = -2$$

The zeros of $H(x) = (2-x)^2 + (2-x) - 20$ are -2 and 7 . The x -intercepts of the graph of H are -2 and 7 .

69. $P(x) = 0$

$$2(x+1)^2 - 5(x+1) - 3 = 0$$

$$\text{Let } u = x+1 \rightarrow u^2 = (x+1)^2$$

$$2u^2 - 5u - 3 = 0$$

$$(2u+1)(u-3) = 0$$

$$2u+1 = 0 \quad \text{or} \quad u-3 = 0$$

$$u = -\frac{1}{2} \quad u = 3$$

$$x+1 = -\frac{1}{2} \quad x = 2$$

$$x = -\frac{3}{2}$$

The zeros of $P(x) = 2(x+1)^2 - 5(x+1) - 3$ are

$-\frac{3}{2}$ and 2 . The x -intercepts of the graph of P are $-\frac{3}{2}$ and 2 .

70. $H(x) = 0$

$$3(1-x)^2 + 5(1-x) + 2 = 0$$

$$\text{Let } u = 1-x \rightarrow u^2 = (1-x)^2$$

$$3u^2 + 5u + 2 = 0$$

$$(3u+2)(u+1) = 0$$

$$3u+2 = 0 \quad \text{or} \quad u+1 = 0$$

$$u = -\frac{2}{3} \quad u = -1$$

$$1-x = -1$$

$$1-x = -\frac{2}{3} \quad x = 2$$

$$x = \frac{5}{3}$$

The zeros of $H(x) = 3(1-x)^2 + 5(1-x) + 2$ are

$\frac{5}{3}$ and 2 . The x -intercepts of the graph of H are

$\frac{5}{3}$ and 2 .

71. $G(x) = 0$

$$x - 4\sqrt{x} = 0$$

$$\text{Let } u = \sqrt{x} \rightarrow u^2 = x$$

$$u^2 - 4u = 0$$

$$u(u-4) = 0$$

$$u = 0 \quad \text{or} \quad u-4 = 0$$

$$u = 4$$

$$\sqrt{x} = 0 \quad \sqrt{x} = 4$$

$$x = 0^2 = 0 \quad x = 4^2 = 16$$

Check:

$$G(0) = 0 - 4\sqrt{0} = 0$$

$$G(16) = 16 - 4\sqrt{16} = 16 - 16 = 0$$

The zeros of $G(x) = x - 4\sqrt{x}$ are 0 and 16 . The x -intercepts of the graph of G are 0 and 16 .

72. $f(x) = 0$

$$x + 8\sqrt{x} = 0$$

$$\text{Let } u = \sqrt{x} \rightarrow u^2 = x$$

Chapter 2: Linear and Quadratic Functions

$$u^2 + 8u = 0$$

$$u(u+8) = 0$$

$$u = 0 \quad \text{or} \quad u + 8 = 0$$

$$u = -8$$

$$\sqrt{x} = 0$$

$$\sqrt{x} = -8$$

$$x = 0^2 = 0$$

$$x = \text{not real}$$

$$\text{Check: } f(0) = 0 + 8\sqrt{0} = 0$$

The only zero of $f(x) = x + 8\sqrt{x}$ is 0. The only x -intercept of the graph of f is 0.

$$73. \quad g(x) = 0$$

$$x + \sqrt{x} - 20 = 0$$

$$\text{Let } u = \sqrt{x} \rightarrow u^2 = x$$

$$u^2 + u - 20 = 0$$

$$(u+5)(u-4) = 0$$

$$u + 5 = 0 \quad \text{or} \quad u - 4 = 0$$

$$u = -5$$

$$u = 4$$

$$\sqrt{x} = -5$$

$$\sqrt{x} = 4$$

$$x = \text{not real}$$

$$x = 4^2 = 16$$

$$\text{Check: } g(16) = 16 + \sqrt{16} - 20 = 16 + 4 - 20 = 0$$

The only zero of $g(x) = x + \sqrt{x} - 20$ is 16. The only x -intercept of the graph of g is 16.

$$74. \quad f(x) = 0$$

$$x + \sqrt{x} - 2 = 0$$

$$\text{Let } u = \sqrt{x} \rightarrow u^2 = x$$

$$u^2 + u - 2 = 0$$

$$(u-1)(u+2) = 0$$

$$u - 1 = 0 \quad \text{or} \quad u + 2 = 0$$

$$u = 1$$

$$u = -2$$

$$\sqrt{x} = 1$$

$$\sqrt{x} = -2$$

$$x = 1^2 = 1$$

$$x = \text{not real}$$

$$\text{Check: } f(1) = 1 + \sqrt{1} - 2 = 1 + 1 - 2 = 0$$

The only zero of $f(x) = x + \sqrt{x} - 2$ is 1. The only x -intercept of the graph of f is 1.

$$75. \quad f(x) = 0$$

$$x^2 - 50 = 0$$

$$x^2 = 50 \Rightarrow x = \pm\sqrt{50} = \pm 5\sqrt{2}$$

The zeros of $f(x) = x^2 - 50$ are $-5\sqrt{2}$ and $5\sqrt{2}$. The x -intercepts of the graph of f are $-5\sqrt{2}$ and $5\sqrt{2}$.

$$76. \quad f(x) = 0$$

$$x^2 - 20 = 0$$

$$x^2 = 20 \Rightarrow x = \pm\sqrt{20} = \pm 2\sqrt{5}$$

The zeros of $f(x) = x^2 - 20$ are $-2\sqrt{5}$ and $2\sqrt{5}$. The x -intercepts of the graph of f are $-2\sqrt{5}$ and $2\sqrt{5}$.

$$77. \quad g(x) = 0$$

$$16x^2 - 8x + 1 = 0$$

$$(4x-1)^2 = 0$$

$$4x - 1 = 0 \Rightarrow x = \frac{1}{4}$$

The only real zero of $g(x) = 16x^2 - 8x + 1$ is $\frac{1}{4}$.

The only x -intercept of the graph of g is $\frac{1}{4}$.

$$78. \quad F(x) = 0$$

$$4x^2 - 12x + 9 = 0$$

$$(2x-3)^2 = 0$$

$$2x - 3 = 0 \Rightarrow x = \frac{3}{2}$$

The only real zero of $F(x) = 4x^2 - 12x + 9$ is $\frac{3}{2}$.

The only x -intercept of the graph of F is $\frac{3}{2}$.

$$79. \quad G(x) = 0$$

$$10x^2 - 19x - 15 = 0$$

$$(5x+3)(2x-5) = 0$$

$$5x + 3 = 0 \quad \text{or} \quad 2x - 5 = 0$$

$$x = -\frac{3}{5}$$

$$x = \frac{5}{2}$$

The zeros of $G(x) = 10x^2 - 19x - 15$ are $-\frac{3}{5}$ and $\frac{5}{2}$.

$\frac{5}{2}$. The x -intercepts of the graph of G are $-\frac{3}{5}$ and $\frac{5}{2}$.

80. $f(x) = 0$

$$6x^2 + 7x - 20 = 0$$

$$(3x - 4)(2x + 5) = 0$$

$$3x - 4 = 0 \quad \text{or} \quad 2x + 5 = 0$$

$$x = \frac{4}{3} \quad x = -\frac{5}{2}$$

The zeros of $f(x) = 6x^2 + 7x - 20$ are $-\frac{5}{2}$ and $\frac{4}{3}$.

The x -intercepts of the graph of f are $-\frac{5}{2}$ and $\frac{4}{3}$.

81. $P(x) = 0$

$$6x^2 - x - 2 = 0$$

$$(3x - 2)(2x + 1) = 0$$

$$3x - 2 = 0 \quad \text{or} \quad 2x + 1 = 0$$

$$x = \frac{2}{3} \quad x = -\frac{1}{2}$$

The zeros of $P(x) = 6x^2 - x - 2$ are $-\frac{1}{2}$ and $\frac{2}{3}$.

The x -intercepts of the graph of P are $-\frac{1}{2}$ and $\frac{2}{3}$.

82. $H(x) = 0$

$$6x^2 + x - 2 = 0$$

$$(3x + 2)(2x - 1) = 0$$

$$3x + 2 = 0 \quad \text{or} \quad 2x - 1 = 0$$

$$x = -\frac{2}{3} \quad x = \frac{1}{2}$$

The zeros of $H(x) = 6x^2 + x - 2$ are $-\frac{2}{3}$ and $\frac{1}{2}$.

The x -intercepts of the graph of H are $-\frac{2}{3}$ and $\frac{1}{2}$.

83. $G(x) = 0$

$$x^2 + \sqrt{2}x - \frac{1}{2} = 0$$

$$2\left(x^2 + \sqrt{2}x - \frac{1}{2}\right) = (0)(2)$$

$$2x^2 + 2\sqrt{2}x - 1 = 0$$

$$a = 2, \quad b = 2\sqrt{2}, \quad c = -1$$

$$\begin{aligned} x &= \frac{-(2\sqrt{2}) \pm \sqrt{(2\sqrt{2})^2 - 4(2)(-1)}}{2(2)} \\ &= \frac{-2\sqrt{2} \pm \sqrt{8+8}}{4} = \frac{-2\sqrt{2} \pm \sqrt{16}}{4} \\ &= \frac{-2\sqrt{2} \pm 4}{4} = \frac{-\sqrt{2} \pm 2}{2} \end{aligned}$$

The zeros of $G(x) = x^2 + \sqrt{2}x - \frac{1}{2}$ are $\frac{-\sqrt{2}-2}{2}$

and $\frac{-\sqrt{2}+2}{2}$. The x -intercepts of the graph of G

are $\frac{-\sqrt{2}-2}{2}$ and $\frac{-\sqrt{2}+2}{2}$.

84. $F(x) = 0$

$$\frac{1}{2}x^2 - \sqrt{2}x - 1 = 0$$

$$2\left(\frac{1}{2}x^2 - \sqrt{2}x - 1\right) = (0)(2)$$

$$x^2 - 2\sqrt{2}x - 2 = 0$$

$$a = 1, \quad b = -2\sqrt{2}, \quad c = -2$$

$$\begin{aligned} x &= \frac{-(-2\sqrt{2}) \pm \sqrt{(-2\sqrt{2})^2 - 4(1)(-2)}}{2(1)} \\ &= \frac{2\sqrt{2} \pm \sqrt{16}}{2} = \frac{2\sqrt{2} \pm 4}{2} = \frac{\sqrt{2} \pm 2}{1} \end{aligned}$$

The zeros of $F(x) = \frac{1}{2}x^2 - \sqrt{2}x - 1$ are $\sqrt{2} - 2$

and $\sqrt{2} + 2$. The x -intercepts of the graph of F are $\sqrt{2} - 2$ and $\sqrt{2} + 2$.

85. $f(x) = 0$

$$x^2 + x - 4 = 0$$

$$a = 1, \quad b = 1, \quad c = -4$$

$$\begin{aligned} x &= \frac{-(1) \pm \sqrt{(1)^2 - 4(1)(-4)}}{2(1)} \\ &= \frac{-1 \pm \sqrt{1+16}}{2} = \frac{-1 \pm \sqrt{17}}{2} \end{aligned}$$

The zeros of $f(x) = x^2 + x - 4$ are $\frac{-1-\sqrt{17}}{2}$ and

$\frac{-1+\sqrt{17}}{2}$. The x -intercepts of the graph of f are

$\frac{-1-\sqrt{17}}{2}$ and $\frac{-1+\sqrt{17}}{2}$.

Chapter 2: Linear and Quadratic Functions

86. $g(x) = 0$

$$x^2 + x - 1 = 0$$

$$a = 1, \quad b = 1, \quad c = -1$$

$$x = \frac{-(1) \pm \sqrt{(1)^2 - 4(1)(-1)}}{2(1)} = \frac{-1 \pm \sqrt{5}}{2}$$


The zeros of $g(x) = x^2 + x - 1$ are $\frac{-1 - \sqrt{5}}{2}$ and

$\frac{-1 + \sqrt{5}}{2}$. The x -intercepts of the graph of g are

$\frac{-1 - \sqrt{5}}{2}$ and $\frac{-1 + \sqrt{5}}{2}$.

87. a. $g(x) = (x-1)^2 - 4$

Using the graph of $y = x^2$, horizontally shift to the right 1 unit, and then vertically shift downward 4 units.



b. $g(x) = 0$

$$(x-1)^2 - 4 = 0$$


$$x^2 - 2x + 1 - 4 = 0$$

$$x^2 - 2x - 3 = 0$$

$$(x+1)(x-3) = 0 \Rightarrow x = -1 \text{ or } x = 3$$

88. a. $F(x) = (x+3)^2 - 9$

Using the graph of $y = x^2$, horizontally shift to the left 3 units, and then vertically shift downward 9 units.



b. $F(x) = 0$

$$(x+3)^2 - 9 = 0$$


$$x^2 + 6x + 9 - 9 = 0$$

$$x^2 + 6x = 0$$

$$x(x+6) = 0 \Rightarrow x = 0 \text{ or } x = -6$$

89. a. $f(x) = 2(x+4)^2 - 8$

Using the graph of $y = x^2$, horizontally shift to the left 4 units, vertically stretch by a factor of 2, and then vertically shift downward 8 units.



b. $f(x) = 0$

$$2(x+4)^2 - 8 = 0$$

$$2(x^2 + 8x + 16) - 8 = 0$$


$$2x^2 + 16x + 32 - 8 = 0$$

$$2x^2 + 16x + 24 = 0$$

$$2(x+2)(x+6) = 0 \Rightarrow x = -2 \text{ or } x = -6$$

90. a. $h(x) = 3(x-2)^2 - 12$

Using the graph of $y = x^2$, horizontally shift to the right 2 units, vertically stretch by a factor of 3, and then vertically shift downward 12 units.



Section 2.3: Quadratic Functions and Their Zeros

b. $h(x) = 0$

$$3(x-2)^2 - 12 = 0$$

$$3(x^2 - 4x + 4) - 12 = 0$$


$$3x^2 - 12x + 12 - 12 = 0$$

$$3x^2 - 12x = 0$$

$$3x(x-4) = 0 \Rightarrow x = 0 \text{ or } x = 4$$

91. a. $H(x) = -3(x-3)^2 + 6$

Using the graph of $y = x^2$, horizontally shift to the right 3 units, vertically stretch by a factor of 3, reflect about the x-axis, and then vertically shift upward 6 units.



b. $H(x) = 0$

$$-3(x-3)^2 + 6 = 0$$

$$-3(x^2 - 6x + 9) + 6 = 0$$

$$-3x^2 + 18x - 27 + 6 = 0$$

$$-3x^2 + 18x - 21 = 0$$

$$-3(x^2 - 6x + 7) = 0$$

$$a = 1, \quad b = -6, \quad c = 7$$


$$x = \frac{-(-6) \pm \sqrt{(-6)^2 - 4(1)(7)}}{2(1)} = \frac{6 \pm \sqrt{36 - 28}}{2}$$

$$= \frac{6 \pm \sqrt{8}}{2} = \frac{6 \pm 2\sqrt{2}}{2} = 3 \pm \sqrt{2}$$

92. a. $f(x) = -2(x+1)^2 + 12$

Using the graph of $y = x^2$, horizontally shift to the left 1 unit, vertically stretch by a factor of 2, reflect about the x-axis, and then

vertically shift upward 12 units.



b. $f(x) = 0$

$$-2(x+1)^2 + 12 = 0$$

$$-2(x^2 + 2x + 1) + 12 = 0$$

$$-2x^2 - 4x - 2 + 12 = 0$$

$$-2x^2 - 4x + 10 = 0$$

$$-2(x^2 + 2x - 5) = 0$$

$$a = 1, \quad b = 2, \quad c = -5$$

$$x = \frac{-(2) \pm \sqrt{(2)^2 - 4(1)(-5)}}{2(1)} = \frac{-2 \pm \sqrt{4 + 20}}{2}$$

$$= \frac{-2 \pm \sqrt{24}}{2} = \frac{-2 \pm 2\sqrt{6}}{2} = -1 \pm \sqrt{6}$$

93. $f(x) = g(x)$

$$5x(x-1) = -7x^2 + 2$$

$$5x^2 - 5x = -7x^2 + 2$$

$$12x^2 - 5x - 2 = 0$$

$$(3x-2)(4x+1) = 0 \Rightarrow x = \frac{2}{3} \text{ or } x = -\frac{1}{4}$$

$$f\left(\frac{2}{3}\right) = 5\left(\frac{2}{3}\right)\left[\left(\frac{2}{3}\right) - 1\right]$$

$$= \left(\frac{10}{3}\right)\left(-\frac{1}{3}\right) = -\frac{10}{9}$$

$$f\left(-\frac{1}{4}\right) = 5\left(-\frac{1}{4}\right)\left[\left(-\frac{1}{4}\right) - 1\right]$$

$$= \left(-\frac{5}{4}\right)\left(-\frac{5}{4}\right) = \frac{25}{16}$$

The points of intersection are:

$$\left(\frac{2}{3}, -\frac{10}{9}\right) \text{ and } \left(-\frac{1}{4}, \frac{25}{16}\right)$$

Chapter 2: Linear and Quadratic Functions

$$\begin{aligned}
 94. \quad & f(x) = g(x) \\
 & 10x(x+2) = -3x+5 \\
 & 10x^2 + 20x = -3x+5 \\
 & 10x^2 + 23x - 5 = 0 \\
 & (2x+5)(5x-1) = 0 \Rightarrow x = -\frac{5}{2} \text{ or } x = \frac{1}{5} \\
 & f\left(-\frac{5}{2}\right) = 10\left(-\frac{5}{2}\right)\left[\left(-\frac{5}{2}\right)+2\right] \\
 & \quad = (-25)\left(-\frac{1}{2}\right) = \frac{25}{2} \\
 & f\left(\frac{1}{5}\right) = 10\left(\frac{1}{5}\right)\left[\left(\frac{1}{5}\right)+2\right] \\
 & \quad = (2)\left(\frac{11}{5}\right) = \frac{22}{5}
 \end{aligned}$$

The points of intersection are:

$$\left(-\frac{5}{2}, \frac{25}{2}\right) \text{ and } \left(\frac{1}{5}, \frac{22}{5}\right)$$

$$\begin{aligned}
 95. \quad & f(x) = g(x) \\
 & 3(x^2 - 4) = 3x^2 + 2x + 4 \\
 & 3x^2 - 12 = 3x^2 + 2x + 4 \\
 & -12 = 2x + 4 \\
 & -16 = 2x \Rightarrow x = -8 \\
 & f(-8) = 3[(-8)^2 - 4] \\
 & \quad = 3[64 - 4] = 180
 \end{aligned}$$

The point of intersection is: $(-8, 180)$

$$\begin{aligned}
 96. \quad & f(x) = g(x) \\
 & 4(x^2 + 1) = 4x^2 - 3x - 8 \\
 & 4x^2 + 4 = 4x^2 - 3x - 8 \\
 & 4 = -3x - 8 \\
 & 12 = -3x \Rightarrow x = -4 \\
 & f(-4) = 4[(-4)^2 + 1] \\
 & \quad = 4[16 + 1] = 68
 \end{aligned}$$

The point of intersection is: $(-4, 68)$

$$\begin{aligned}
 97. \quad & f(x) = g(x) \\
 & \frac{3x}{x+2} - \frac{5}{x+1} = \frac{-5}{x^2+3x+2} \\
 & \frac{3x}{x+2} - \frac{5}{x+1} = \frac{-5}{(x+2)(x+1)} \\
 & 3x(x+1) - 5(x+2) = -5 \\
 & 3x^2 + 3x - 5x - 10 = -5 \\
 & 3x^2 - 2x - 5 = 0 \\
 & (3x-5)(x+1) = 0 \\
 & x = \frac{5}{3} \text{ or } x = -1 \\
 & f\left(\frac{5}{3}\right) = \frac{3\left(\frac{5}{3}\right)}{\left(\frac{5}{3}\right)+2} - \frac{5}{\left(\frac{5}{3}\right)+1} \\
 & \quad = \frac{\left(\frac{5}{3}\right)}{\left(\frac{11}{3}\right)} - \frac{5}{\left(\frac{8}{3}\right)} \\
 & \quad = \frac{15}{11} - \frac{15}{8} \\
 & \quad = -\frac{45}{88}
 \end{aligned}$$

The point of intersection is: $\left(\frac{5}{3}, -\frac{45}{88}\right)$

$$\begin{aligned}
 98. \quad & f(x) = g(x) \\
 & \frac{2x}{x-3} - \frac{3}{x+1} = \frac{2x+18}{x^2-2x-3} \\
 & \frac{2x}{x-3} - \frac{3}{x+1} = \frac{2x+18}{(x-3)(x+1)} \\
 & 2x(x+1) - 3(x-3) = 2x+18 \\
 & 2x^2 + 2x - 3x + 9 = 2x+18 \\
 & 2x^2 - 3x - 9 = 0 \\
 & (2x+3)(x-3) = 0 \\
 & x = -\frac{3}{2} \text{ or } x = 3
 \end{aligned}$$

Section 2.3: Quadratic Functions and Their Zeros

$$\begin{aligned} f\left(-\frac{3}{2}\right) &= \frac{2\left(-\frac{3}{2}\right)}{\left(-\frac{3}{2}\right)-3} - \frac{3}{\left(-\frac{3}{2}\right)+1} \\ &= \frac{(-3)}{\left(-\frac{9}{2}\right)} - \frac{3}{\left(-\frac{1}{2}\right)} \\ &= \frac{6}{9} + 6 = \frac{2}{3} + 6 = \frac{20}{3} \end{aligned}$$

The point of intersection is: $\left(-\frac{3}{2}, \frac{20}{3}\right)$

99. a. $(f+g)(x) =$

$$\begin{aligned} &= x^2 + 5x - 14 + x^2 + 3x - 4 \\ &= 2x^2 + 8x - 18 = 0 \\ &x^2 + 4x - 9 = 0 \\ x &= \frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(-9)}}{2(1)} = \frac{-4 \pm \sqrt{16+36}}{2} \\ &= \frac{-4 \pm \sqrt{52}}{2} = \frac{-4 \pm 2\sqrt{13}}{2} = -2 \pm \sqrt{13} \end{aligned}$$

b. $(f-g)(x) =$

$$\begin{aligned} &= (x^2 + 5x - 14) - (x^2 + 3x - 4) \\ &= x^2 + 5x - 14 - x^2 - 3x + 4 \\ &= 2x - 10 \\ 2x - 10 &= 0 \Rightarrow x = 5 \end{aligned}$$

c. $(f \cdot g)(x) =$

$$\begin{aligned} &= (x^2 + 5x - 14)(x^2 + 3x - 4) \\ &= (x+7)(x-2)(x+4)(x-1) \end{aligned}$$

$$\begin{aligned} (f \cdot g)(x) &= 0 \\ 0 &= (x+7)(x-2)(x+4)(x-1) \\ \Rightarrow x &= -7 \text{ or } x = 2 \text{ or } x = -4 \text{ or } x = 1 \end{aligned}$$

100. a. $(f+g)(x) =$

$$\begin{aligned} &= x^2 - 3x - 18 + x^2 + 2x - 3 \\ &= 2x^2 - x - 21 \\ 2x^2 - x - 21 &= 0 \\ (2x-7)(x+3) &= 0 \Rightarrow x = \frac{7}{2} \text{ or } x = -3 \end{aligned}$$

b. $(f-g)(x) =$

$$\begin{aligned} &= (x^2 - 3x - 18) - (x^2 + 2x - 3) \\ &= x^2 - 3x - 18 - x^2 - 2x + 3 \\ &= -5x - 15 \\ -5x - 15 &= 0 \Rightarrow x = -3 \end{aligned}$$

c. $(f \cdot g)(x) =$

$$\begin{aligned} &= (x^2 - 3x - 18)(x^2 + 2x - 3) \\ &= (x+3)(x-6)(x+3)(x-1) \\ (f \cdot g)(x) &= 0 \\ 0 &= (x+3)(x-6)(x+3)(x-1) \\ \Rightarrow x &= -3 \text{ or } x = 6 \text{ or } x = 1 \end{aligned}$$

101. $A(x) = 143$

$$\begin{aligned} x(x+2) &= 143 \\ x^2 + 2x - 143 &= 0 \\ (x+13)(x-11) &= 0 \\ \cancel{x = -13} \text{ or } x &= 11 \end{aligned}$$

Discard the negative solution since width cannot be negative. The width of the rectangular window is 11 feet and the length is 13 feet.

102. $A(x) = 306$

$$\begin{aligned} x(x+1) &= 306 \\ x^2 + x - 306 &= 0 \\ (x+18)(x-17) &= 0 \\ \cancel{x = -18} \text{ or } x &= 17 \end{aligned}$$

Discard the negative solution since width cannot be negative. The width of the rectangular window is 17 cm and the length is 18 cm.

103. $V(x) = 4$

$$\begin{aligned} (x-2)^2 &= 4 \\ x-2 &= \pm\sqrt{4} \\ x-2 &= \pm 2 \\ x &= 2 \pm 2 \\ x &= 4 \text{ or } x = 0 \end{aligned}$$

Discard $x = 0$ since that is not a feasible length for the original sheet. Therefore, the original sheet should measure 4 feet on each side.

Chapter 2: Linear and Quadratic Functions

104. $V(x) = 4$

$$(x-2)^2 = 16$$

$$x-2 = \pm\sqrt{16}$$

$$x-2 = \pm 4$$

$$x = 2 \pm 4$$

$$x = 6 \quad \text{or} \quad \cancel{x = -2}$$

Discard $x = -2$ since width cannot be negative.
Therefore, the original sheet should measure 6 feet on each side.

105. a. When the ball strikes the ground, the distance from the ground will be 0.

Therefore, we solve

$$s = 0$$

$$96 + 80t - 16t^2 = 0$$

$$-16t^2 + 80t + 96 = 0$$

$$t^2 - 5t - 6 = 0$$

$$(t-6)(t+1) = 0$$

$$t = 6 \quad \text{or} \quad \cancel{t = -1}$$

Discard the negative solution since the time of flight must be positive. The ball will strike the ground after 6 seconds.

- b. When the ball passes the top of the building, it will be 96 feet from the ground. Therefore, we solve

$$s = 96$$

$$96 + 80t - 16t^2 = 96$$

$$-16t^2 + 80t = 0$$

$$t^2 - 5t = 0$$

$$t(t-5) = 0$$

$$t = 0 \quad \text{or} \quad t = 5$$

The ball is at the top of the building at time $t = 0$ seconds when it is thrown. It will pass the top of the building on the way down after 5 seconds.

106. a. To find when the object will be 15 meters above the ground, we solve

$$s = 15$$

$$-4.9t^2 + 20t = 15$$

$$-4.9t^2 + 20t - 15 = 0$$

$$a = -4.9, b = 20, c = -15$$

$$t = \frac{-20 \pm \sqrt{20^2 - 4(-4.9)(-15)}}{2(-4.9)}$$

$$= \frac{-20 \pm \sqrt{106}}{-9.8}$$

$$= \frac{20 \pm \sqrt{106}}{9.8}$$

$$t \approx 0.99 \quad \text{or} \quad t \approx 3.09$$

The object will be 15 meters above the ground after about 0.99 seconds (on the way up) and about 3.09 seconds (on the way down).

- b. The object will strike the ground when the distance from the ground is 0. Thus, we solve

$$s = 0$$

$$-4.9t^2 + 20t = 0$$

$$t(-4.9t + 20) = 0$$

$$t = 0 \quad \text{or} \quad -4.9t + 20 = 0$$

$$-4.9t = -20$$

$$t \approx 4.08$$

The object will strike the ground after about 4.08 seconds.

- c. $s = 100$

$$-4.9t^2 + 20t = 100$$

$$-4.9t^2 + 20t - 100 = 0$$

$$a = -4.9, b = 20, c = -100$$

$$t = \frac{-20 \pm \sqrt{20^2 - 4(-4.9)(-100)}}{2(-4.9)}$$

$$= \frac{-20 \pm \sqrt{-1560}}{-9.8}$$

There is no real solution. The object never reaches a height of 100 meters.

107. For the sum to be 210, we solve

$$S(n) = 210$$

$$\frac{1}{2}n(n+1) = 210$$

$$n(n+1) = 420$$

$$n^2 + n - 420 = 0$$

$$(n-20)(n+21) = 0$$

$$n-20 = 0 \quad \text{or} \quad n+21 = 0$$

$$n = 20 \quad \text{or} \quad \cancel{n = -21}$$

Discard the negative solution since the number of consecutive integers must be positive. For a sum of 210, we must add the 20 consecutive integers, starting at 1.

- 108.** To determine the number of sides when a polygon has 65 diagonals, we solve

$$D(n) = 65$$

$$\frac{1}{2}n(n-3) = 65$$

$$n(n-3) = 130$$

$$n^2 - 3n - 130 = 0$$

$$(n+10)(n-13) = 0$$

$$n+10 = 0 \quad \text{or} \quad n-13 = 0$$

$$\cancel{n = -10} \quad n = 13$$

Discard the negative solution since the number of sides must be positive. A polygon with 65 diagonals will have 13 sides.

To determine the number of sides if a polygon has 80 diagonals, we solve

$$D(n) = 80$$

$$\frac{1}{2}n(n-3) = 80$$

$$n(n-3) = 160$$

$$n^2 - 3n - 160 = 0$$

$$a = 1, b = -3, c = -160$$

$$t = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(1)(-160)}}{2(1)}$$

$$= \frac{3 \pm \sqrt{649}}{2}$$

Since the solutions are not integers, a polygon with 80 diagonals is not possible.

- 109.** The roots of a quadratic equation are

$$x_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a} \quad \text{and} \quad x_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a},$$

so the sum of the roots is

$$\begin{aligned} x_1 + x_2 &= \frac{-b - \sqrt{b^2 - 4ac}}{2a} + \frac{-b + \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-b - \sqrt{b^2 - 4ac} - b + \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-2b}{2a} = -\frac{b}{a} \end{aligned}$$

- 110.** The roots of a quadratic equation are

$$x_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a} \quad \text{and} \quad x_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a},$$

so the product of the roots is

$$\begin{aligned} x_1 \cdot x_2 &= \left(\frac{-b - \sqrt{b^2 - 4ac}}{2a} \right) \left(\frac{-b + \sqrt{b^2 - 4ac}}{2a} \right) \\ &= \frac{(-b)^2 - (\sqrt{b^2 - 4ac})^2}{(2a)^2} = \frac{b^2 - (b^2 - 4ac)}{4a^2} \\ &= \frac{b^2 - b^2 + 4ac}{4a^2} = \frac{4ac}{4a^2} = \frac{c}{a} \end{aligned}$$

- 111.** In order to have one repeated real zero, we need the discriminant to be 0.

$$b^2 - 4ac = 0$$

$$1^2 - 4(k)(k) = 0$$

$$1 - 4k^2 = 0$$

$$4k^2 = 1$$

$$k^2 = \frac{1}{4}$$

$$k = \pm \sqrt{\frac{1}{4}}$$

$$k = \frac{1}{2} \quad \text{or} \quad k = -\frac{1}{2}$$

- 112.** In order to have one repeated real zero, we need the discriminant to be 0.

$$b^2 - 4ac = 0$$

$$(-k)^2 - 4(1)(4) = 0$$

$$k^2 - 16 = 0$$

$$(k-4)(k+4) = 0$$

$$k = 4 \quad \text{or} \quad k = -4$$

- 113.** For $f(x) = ax^2 + bx + c = 0$:

$$x_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a} \quad \text{and} \quad x_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$

For $f(x) = ax^2 - bx + c = 0$:

$$\begin{aligned} x_1^* &= \frac{-(-b) - \sqrt{(-b)^2 - 4ac}}{2a} \\ &= \frac{b - \sqrt{b^2 - 4ac}}{2a} = -\left(\frac{-b + \sqrt{b^2 - 4ac}}{2a} \right) = -x_2 \end{aligned}$$

Chapter 2: Linear and Quadratic Functions

and

$$\begin{aligned}x_2^* &= \frac{-(-b) + \sqrt{(-b)^2 - 4ac}}{2a} \\&= \frac{b + \sqrt{b^2 - 4ac}}{2a} = -\left(\frac{-b - \sqrt{b^2 - 4ac}}{2a}\right) = -x_1\end{aligned}$$

- 114.** For $f(x) = ax^2 + bx + c = 0$:

$$x_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a} \quad \text{and} \quad x_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$

For $f(x) = cx^2 + bx + a = 0$:

$$\begin{aligned}x_1^* &= \frac{-b - \sqrt{b^2 - 4(c)(a)}}{2c} = \frac{-b - \sqrt{b^2 - 4ac}}{2c} \\&= \frac{-b - \sqrt{b^2 - 4ac}}{2c} \cdot \frac{-b + \sqrt{b^2 - 4ac}}{-b + \sqrt{b^2 - 4ac}} \\&= \frac{b^2 - (b^2 - 4ac)}{2c(-b + \sqrt{b^2 - 4ac})} = \frac{4ac}{2c(-b + \sqrt{b^2 - 4ac})} \\&= \frac{2a}{-b + \sqrt{b^2 - 4ac}} = \frac{1}{x_2}\end{aligned}$$

and

$$\begin{aligned}x_2^* &= \frac{-b + \sqrt{b^2 - 4(c)(a)}}{2c} = \frac{-b + \sqrt{b^2 - 4ac}}{2c} \\&= \frac{-b + \sqrt{b^2 - 4ac}}{2c} \cdot \frac{-b - \sqrt{b^2 - 4ac}}{-b - \sqrt{b^2 - 4ac}} \\&= \frac{b^2 - (b^2 - 4ac)}{2c(-b - \sqrt{b^2 - 4ac})} = \frac{4ac}{2c(-b - \sqrt{b^2 - 4ac})} \\&= \frac{2a}{-b - \sqrt{b^2 - 4ac}} = \frac{1}{x_1}\end{aligned}$$

- 115. a.** $x^2 = 9$ and $x = 3$ are not equivalent because they do not have the same solution set. In the first equation we can also have $x = -3$.
- b.** $x = \sqrt{9}$ and $x = 3$ are equivalent because $\sqrt{9} = 3$.
- c.** $(x-1)(x-2) = (x-1)^2$ and $x-2 = x-1$ are not equivalent because they do not have the same solution set. The first equation has the

solution set $\{1\}$ while the second equation has no solutions.

- 116.** Answers may vary. Methods discussed in this section include factoring, the square root method, completing the square, and the quadratic formula.
- 117.** Answers will vary. Knowing the discriminant allows us to know how many real solutions the equation will have.
- 118.** Answers will vary. One possibility:
Two distinct: $f(x) = x^2 - 3x - 18$
One repeated: $f(x) = x^2 - 14x + 49$
No real: $f(x) = x^2 + x + 4$
- 119.** Answers will vary.
- 120.** Two quadratic functions can intersect 0, 1, or 2 times.

Section 2.4

- 1.** $y = x^2 - 9$

To find the y-intercept, let $x = 0$:

$$y = 0^2 - 9 = -9.$$

To find the x-intercept(s), let $y = 0$:

$$x^2 - 9 = 0$$

$$x^2 = 9$$

$$x = \pm\sqrt{9} = \pm 3$$

The intercepts are $(0, -9)$, $(-3, 0)$, and $(3, 0)$.

- 2.** $2x^2 + 7x - 4 = 0$

$$(2x-1)(x+4) = 0$$

$$2x-1=0 \quad \text{or} \quad x+4=0$$

$$2x=1 \quad \text{or} \quad x=-4$$

$$x = \frac{1}{2} \quad \text{or} \quad x = -4$$

The solution set is $\left\{-4, \frac{1}{2}\right\}$.

- 3.** $\left(\frac{1}{2} \cdot (-5)\right)^2 = \frac{25}{4}$

- 4.** right; 4

Section 2.4: Properties of Quadratic Functions

5. parabola

6. axis (or axis of symmetry)

7. $-\frac{b}{2a}$

8. True; $a = 2 > 0$.

9. True; $-\frac{b}{2a} = -\frac{4}{2(-1)} = 2$

10. True

11. C

12. E

13. F

14. A

15. G


16. B

17. H

18. D

19. $f(x) = \frac{1}{4}x^2 - 2$


Using the graph of $y = x^2$, compress vertically by a factor of $\frac{1}{4}$, then shift down 2 units.



20. $f(x) = 2x^2 + 4$


Using the graph of $y = x^2$, stretch vertically by a

factor of 2, then shift up 4 units.




21. $f(x) = (x+2)^2 - 2$

Using the graph of $y = x^2$, shift left 2 units, then shift down 2 units.



22. $f(x) = (x-3)^2 - 10$

Using the graph of $y = x^2$, shift right 3 units, then shift down 10 units.




23. $f(x) = 2(x-1)^2 - 1$

Using the graph of $y = x^2$, shift right 1 unit, stretch vertically by a factor of 2, then shift


Chapter 2: Linear and Quadratic Functions

down 1 unit.



24. $f(x) = 3(x+1)^2 - 3$


Using the graph of $y = x^2$, shift left 1 unit, stretch vertically by a factor of 3, then shift down 3 units.



25. $f(x) = -x^2 - 2x$


$$\begin{aligned} &= -(x^2 + 2x) \\ &= -(x^2 + 2x + 1) + 1 \\ &= -(x+1)^2 + 1 \end{aligned}$$

Using the graph of $y = x^2$, shift left 1 unit, reflect across the x -axis, then shift up 1 unit.




26.
$$\begin{aligned} f(x) &= -2x^2 + 6x + 2 \\ &= -2(x^2 - 3x) + 2 \\ &= -2\left(x^2 - 3x + \frac{9}{4}\right) + 2 + \frac{9}{2} \\ &= -2\left(x - \frac{3}{2}\right)^2 + \frac{13}{2} \end{aligned}$$

Using the graph of $y = x^2$, shift right $\frac{3}{2}$ units, reflect about the x -axis, stretch vertically by a factor of 2, then shift up $\frac{13}{2}$ units.



27.
$$\begin{aligned} f(x) &= 2x^2 + 4x - 2 \\ &= 2(x^2 + 2x) - 2 \\ &= 2(x^2 + 2x + 1) - 2 - 2 \\ &= 2(x+1)^2 - 4 \end{aligned}$$

Using the graph of $y = x^2$, shift left 1 unit, stretch vertically by a factor of 2, then shift down 4 units.




28.
$$\begin{aligned} f(x) &= 3x^2 + 12x + 5 \\ &= 3(x^2 + 4x) + 5 \\ &= 3(x^2 + 4x + 4) + 5 - 12 \\ &= 3(x+2)^2 - 7 \end{aligned}$$

Using the graph of $y = x^2$, shift left 2 units,

Section 2.4: Properties of Quadratic Functions

stretch vertically by a factor of 3, then shift down 7 units.



- 29. a.** For $f(x) = x^2 + 2x$, $a = 1$, $b = 2$, $c = 0$.

Since $a = 1 > 0$, the graph opens up.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(2)}{2(1)} = \frac{-2}{2} = -1.$$

The y -coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(-1) = (-1)^2 + 2(-1) = 1 - 2 = -1.$$

Thus, the vertex is $(-1, -1)$.

The axis of symmetry is the line $x = -1$.

The discriminant is

$$b^2 - 4ac = (2)^2 - 4(1)(0) = 4 > 0, \text{ so the graph}$$

has two x -intercepts.

The x -intercepts are found by solving:


$$x^2 + 2x = 0$$

$$x(x + 2) = 0$$

$$x = 0 \text{ or } x = -2$$

The x -intercepts are -2 and 0 .

The y -intercept is $f(0) = 0$.



- b.** The domain is $(-\infty, \infty)$.

The range is $[-1, \infty)$.

- c.** Decreasing on $(-\infty, -1)$.

Increasing on $(-1, \infty)$.

- 30. a.** For $f(x) = x^2 - 4x$, $a = 1$, $b = -4$, $c = 0$.

Since $a = 1 > 0$, the graph opens up.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-4)}{2(1)} = \frac{4}{2} = 2.$$

The y -coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(2) = (2)^2 - 4(2) = 4 - 8 = -4.$$

Thus, the vertex is $(2, -4)$.

The axis of symmetry is the line $x = 2$.

The discriminant is:

$$b^2 - 4ac = (-4)^2 - 4(1)(0) = 16 > 0, \text{ so the}$$

graph has two x -intercepts.

The x -intercepts are found by solving:


$$x^2 - 4x = 0$$

$$x(x - 4) = 0$$

$$x = 0 \text{ or } x = 4.$$

The x -intercepts are 0 and 4 .

The y -intercept is $f(0) = 0$.



- b.** The domain is $(-\infty, \infty)$.

The range is $[-4, \infty)$.

- c.** Decreasing on $(-\infty, 2)$.

Increasing on $(2, \infty)$.

- 31. a.** For $f(x) = -x^2 - 6x$, $a = -1$, $b = -6$,

$c = 0$. Since $a = -1 < 0$, the graph opens down. The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-6)}{2(-1)} = \frac{6}{-2} = -3.$$

The y -coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(-3) = -(-3)^2 - 6(-3) = -9 + 18 = 9.$$

Thus, the vertex is $(-3, 9)$.

The axis of symmetry is the line $x = -3$.

Chapter 2: Linear and Quadratic Functions

The discriminant is:

$$b^2 - 4ac = (-6)^2 - 4(-1)(0) = 36 > 0,$$

so the graph has two x -intercepts.

The x -intercepts are found by solving:


$$-x^2 - 6x = 0$$

$$-x(x + 6) = 0$$

$$x = 0 \text{ or } x = -6.$$

The x -intercepts are -6 and 0 .

The y -intercept is $f(0) = 0$.



- b. The domain is $(-\infty, \infty)$.

The range is $(-\infty, 9]$.

- c. Increasing on $(-\infty, -3)$.

Decreasing on $(-3, \infty)$.

32. a. For $f(x) = -x^2 + 4x$, $a = -1$, $b = 4$, $c = 0$.

Since $a = -1 < 0$, the graph opens down.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-4}{2(-1)} = \frac{-4}{-2} = 2.$$

The y -coordinate of the vertex is

$$\begin{aligned} f\left(\frac{-b}{2a}\right) &= f(2) \\ &= -(2)^2 + 4(2) \\ &= 4. \end{aligned}$$

Thus, the vertex is $(2, 4)$.

The axis of symmetry is the line $x = 2$.

The discriminant is:

$$b^2 - 4ac = 4^2 - 4(-1)(0) = 16 > 0,$$

so the graph has two x -intercepts.

The x -intercepts are found by solving:


$$-x^2 + 4x = 0$$

$$-x(x - 4) = 0$$

$$x = 0 \text{ or } x = 4.$$

The x -intercepts are 0 and 4 .

The y -intercept is $f(0) = 0$.



- b. The domain is $(-\infty, \infty)$.

The range is $(-\infty, 4]$.

- c. Increasing on $(-\infty, 2)$.

Decreasing on $(2, \infty)$.

33. a. For $f(x) = x^2 + 2x - 8$, $a = 1$, $b = 2$, $c = -8$.

Since $a = 1 > 0$, the graph opens up.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-2}{2(1)} = \frac{-2}{2} = -1.$$

The y -coordinate of the vertex is

$$\begin{aligned} f\left(\frac{-b}{2a}\right) &= f(-1) = (-1)^2 + 2(-1) - 8 \\ &= 1 - 2 - 8 = -9. \end{aligned}$$

Thus, the vertex is $(-1, -9)$.

The axis of symmetry is the line $x = -1$.

The discriminant is:

$$b^2 - 4ac = 2^2 - 4(1)(-8) = 4 + 32 = 36 > 0,$$

so the graph has two x -intercepts.

The x -intercepts are found by solving:


$$x^2 + 2x - 8 = 0$$

$$(x + 4)(x - 2) = 0$$

$$x = -4 \text{ or } x = 2.$$

The x -intercepts are -4 and 2 .

The y -intercept is $f(0) = -8$.



Section 2.4: Properties of Quadratic Functions

- b. The domain is $(-\infty, \infty)$.

The range is $[-9, \infty)$.

- c. Decreasing on $(-\infty, -1)$.

Increasing on $(-1, \infty)$.

34. a. For $f(x) = x^2 - 2x - 3$, $a = 1$, $b = -2$,
 $c = -3$.

Since $a = 1 > 0$, the graph opens up.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-2)}{2(1)} = \frac{2}{2} = 1.$$

The y -coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(1) = 1^2 - 2(1) - 3 = -4.$$

Thus, the vertex is $(1, -4)$.

The axis of symmetry is the line $x = 1$.

The discriminant is:

$$b^2 - 4ac = (-2)^2 - 4(1)(-3) = 4 + 12 = 16 > 0,$$

so the graph has two x -intercepts.

The x -intercepts are found by solving:


$$x^2 - 2x - 3 = 0$$

$$(x+1)(x-3) = 0$$

$$x = -1 \text{ or } x = 3.$$

The x -intercepts are -1 and 3 .

The y -intercept is $f(0) = -3$.



- b. The domain is $(-\infty, \infty)$.

The range is $[-4, \infty)$.

- c. Decreasing on $(-\infty, 1)$.

Increasing on $(1, \infty)$.

35. a. For $f(x) = x^2 + 2x + 1$, $a = 1$, $b = 2$, $c = 1$.

Since $a = 1 > 0$, the graph opens up.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-2}{2(1)} = \frac{-2}{2} = -1.$$

The y -coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(-1)$$

$$= (-1)^2 + 2(-1) + 1 = 1 - 2 + 1 = 0.$$

Thus, the vertex is $(-1, 0)$.

The axis of symmetry is the line $x = -1$.

The discriminant is:

$$b^2 - 4ac = 2^2 - 4(1)(1) = 4 - 4 = 0,$$

so the graph has one x -intercept.

The x -intercept is found by solving:


$$x^2 + 2x + 1 = 0$$

$$(x+1)^2 = 0$$

$$x = -1.$$

The x -intercept is -1 .

The y -intercept is $f(0) = 1$.



- b. The domain is $(-\infty, \infty)$. The range is $[0, \infty)$.

- c. Decreasing on $(-\infty, -1)$.

Increasing on $(-1, \infty)$.

36. a. For $f(x) = x^2 + 6x + 9$, $a = 1$, $b = 6$, $c = 9$.

Since $a = 1 > 0$, the graph opens up.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-6}{2(1)} = \frac{-6}{2} = -3.$$

The y -coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(-3) = (-3)^2 + 6(-3) + 9$$

$$= 9 - 18 + 9 = 0.$$

Thus, the vertex is $(-3, 0)$.

The axis of symmetry is the line $x = -3$.

The discriminant is:

$$b^2 - 4ac = 6^2 - 4(1)(9) = 36 - 36 = 0,$$

so the graph has one x -intercept.

The x -intercept is found by solving:

$$x^2 + 6x + 9 = 0$$


$$(x+3)^2 = 0$$

$$x = -3.$$

Chapter 2: Linear and Quadratic Functions

The x -intercept is -3 .

The y -intercept is $f(0) = 9$.



- b. The domain is $(-\infty, \infty)$. The range is $[0, \infty)$.
- c. Decreasing on $(-\infty, -3)$.
Increasing on $(-3, \infty)$.

37. a. For $f(x) = 2x^2 - x + 2$, $a = 2$, $b = -1$, $c = 2$.

Since $a = 2 > 0$, the graph opens up.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-1)}{2(2)} = \frac{1}{4}.$$

The y -coordinate of the vertex is

$$\begin{aligned} f\left(\frac{-b}{2a}\right) &= f\left(\frac{1}{4}\right) = 2\left(\frac{1}{4}\right)^2 - \frac{1}{4} + 2 \\ &= \frac{1}{8} - \frac{1}{4} + 2 = \frac{15}{8}. \end{aligned}$$

Thus, the vertex is $\left(\frac{1}{4}, \frac{15}{8}\right)$.


The axis of symmetry is the line $x = \frac{1}{4}$.

The discriminant is:

$$b^2 - 4ac = (-1)^2 - 4(2)(2) = 1 - 16 = -15,$$

so the graph has no x -intercepts.

The y -intercept is $f(0) = 2$.



- b. The domain is $(-\infty, \infty)$.
The range is $\left[\frac{15}{8}, \infty\right)$.

- c. Decreasing on $\left(-\infty, \frac{1}{4}\right)$.
Increasing on $\left(\frac{1}{4}, \infty\right)$.

38. a. For $f(x) = 4x^2 - 2x + 1$, $a = 4$, $b = -2$, $c = 1$.

Since $a = 4 > 0$, the graph opens up.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-2)}{2(4)} = \frac{2}{8} = \frac{1}{4}.$$

The y -coordinate of the vertex is

$$\begin{aligned} f\left(\frac{-b}{2a}\right) &= f\left(\frac{1}{4}\right) = 4\left(\frac{1}{4}\right)^2 - 2\left(\frac{1}{4}\right) + 1 \\ &= \frac{1}{4} - \frac{1}{2} + 1 = \frac{3}{4}. \end{aligned}$$

Thus, the vertex is $\left(\frac{1}{4}, \frac{3}{4}\right)$.


The axis of symmetry is the line $x = \frac{1}{4}$.

The discriminant is:

$$b^2 - 4ac = (-2)^2 - 4(4)(1) = 4 - 16 = -12,$$

so the graph has no x -intercepts.

The y -intercept is $f(0) = 1$.



- b. The domain is $(-\infty, \infty)$.

The range is $\left[\frac{3}{4}, \infty\right)$.

- c. Decreasing on $\left(-\infty, \frac{1}{4}\right)$.
Increasing on $\left(\frac{1}{4}, \infty\right)$.

39. a. For $f(x) = -2x^2 + 2x - 3$, $a = -2$, $b = 2$, $c = -3$. Since $a = -2 < 0$, the graph opens down.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(2)}{2(-2)} = \frac{-2}{-4} = \frac{1}{2}.$$

The y -coordinate of the vertex is

Section 2.4: Properties of Quadratic Functions

$$\begin{aligned} f\left(\frac{-b}{2a}\right) &= f\left(\frac{1}{2}\right) = -2\left(\frac{1}{2}\right)^2 + 2\left(\frac{1}{2}\right) - 3 \\ &= -\frac{1}{2} + 1 - 3 = -\frac{5}{2}. \end{aligned}$$

Thus, the vertex is $\left(\frac{1}{2}, -\frac{5}{2}\right)$.


The axis of symmetry is the line $x = \frac{1}{2}$.

The discriminant is:

$$b^2 - 4ac = 2^2 - 4(-2)(-3) = 4 - 24 = -20,$$

so the graph has no x -intercepts.

The y -intercept is $f(0) = -3$.



- b. The domain is $(-\infty, \infty)$.

The range is $\left(-\infty, -\frac{5}{2}\right]$.

- c. Increasing on $\left(-\infty, \frac{1}{2}\right)$.

Decreasing on $\left(\frac{1}{2}, \infty\right)$.

40. a. For $f(x) = -3x^2 + 3x - 2$, $a = -3$, $b = 3$, $c = -2$. Since $a = -3 < 0$, the graph opens down.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-3}{2(-3)} = \frac{-3}{-6} = \frac{1}{2}.$$

The y -coordinate of the vertex is

$$\begin{aligned} f\left(\frac{-b}{2a}\right) &= f\left(\frac{1}{2}\right) = -3\left(\frac{1}{2}\right)^2 + 3\left(\frac{1}{2}\right) - 2 \\ &= -\frac{3}{4} + \frac{3}{2} - 2 = -\frac{5}{4}. \end{aligned}$$

Thus, the vertex is $\left(\frac{1}{2}, -\frac{5}{4}\right)$.


The axis of symmetry is the line $x = \frac{1}{2}$.

The discriminant is:

$$b^2 - 4ac = 3^2 - 4(-3)(-2) = 9 - 24 = -15,$$

so the graph has no x -intercepts.

The y -intercept is $f(0) = -2$.



- b. The domain is $(-\infty, \infty)$.

The range is $\left(-\infty, -\frac{5}{4}\right]$.

- c. Increasing on $\left(-\infty, \frac{1}{2}\right)$.

Decreasing on $\left(\frac{1}{2}, \infty\right)$.

41. a. For $f(x) = -4x^2 - 6x + 2$, $a = -4$, $b = -6$, $c = 2$. Since $a = -4 < 0$, the graph opens down.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-6)}{2(-4)} = \frac{6}{-8} = -\frac{3}{4}.$$

The y -coordinate of the vertex is

$$\begin{aligned} f\left(\frac{-b}{2a}\right) &= f\left(-\frac{3}{4}\right) = -4\left(-\frac{3}{4}\right)^2 - 6\left(-\frac{3}{4}\right) + 2 \\ &= -\frac{9}{4} + \frac{9}{2} + 2 = \frac{17}{4}. \end{aligned}$$

Thus, the vertex is $\left(-\frac{3}{4}, \frac{17}{4}\right)$.

The axis of symmetry is the line $x = -\frac{3}{4}$.

The discriminant is:

$$b^2 - 4ac = (-6)^2 - 4(-4)(2) = 36 + 32 = 68,$$

so the graph has two x -intercepts.

The x -intercepts are found by solving:

$$-4x^2 - 6x + 2 = 0$$


$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-6) \pm \sqrt{68}}{2(-4)}$$

$$= \frac{6 \pm \sqrt{68}}{-8} = \frac{6 \pm 2\sqrt{17}}{-8} = \frac{3 \pm \sqrt{17}}{-4}$$

Chapter 2: Linear and Quadratic Functions

The x -intercepts are $\frac{-3+\sqrt{17}}{4}$ and $\frac{-3-\sqrt{17}}{4}$.

The y -intercept is $f(0) = 2$.



- b. The domain is $(-\infty, \infty)$.

The range is $(-\infty, \frac{17}{4}]$.

- c. Decreasing on $(-\frac{3}{4}, \infty)$.

Increasing on $(-\infty, -\frac{3}{4})$.

42. a. For $f(x) = 3x^2 - 8x + 2$, $a = 3$, $b = -8$, $c = 2$.

Since $a = 3 > 0$, the graph opens up.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-8)}{2(3)} = \frac{8}{6} = \frac{4}{3}.$$

The y -coordinate of the vertex is

$$\begin{aligned} f\left(\frac{-b}{2a}\right) &= f\left(\frac{4}{3}\right) = 3\left(\frac{4}{3}\right)^2 - 8\left(\frac{4}{3}\right) + 2 \\ &= \frac{16}{3} - \frac{32}{3} + 2 = -\frac{10}{3}. \end{aligned}$$

Thus, the vertex is $(\frac{4}{3}, -\frac{10}{3})$.

The axis of symmetry is the line $x = \frac{4}{3}$.

The discriminant is:

$$b^2 - 4ac = (-8)^2 - 4(3)(2) = 64 - 24 = 40,$$

so the graph has two x -intercepts.

The x -intercepts are found by solving:


$$3x^2 - 8x + 2 = 0$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-8) \pm \sqrt{40}}{2(3)}$$

$$= \frac{8 \pm \sqrt{40}}{6} = \frac{8 \pm 2\sqrt{10}}{6} = \frac{4 \pm \sqrt{10}}{3}$$

The x -intercepts are $\frac{4+\sqrt{10}}{3}$ and $\frac{4-\sqrt{10}}{3}$.

The y -intercept is $f(0) = 2$.



- b. The domain is $(-\infty, \infty)$.

The range is $[-\frac{10}{3}, \infty)$.

- c. Decreasing on $(-\infty, \frac{4}{3})$.

Increasing on $(\frac{4}{3}, \infty)$.

43. Consider the form $y = a(x-h)^2 + k$. From the graph we know that the vertex is $(-1, -2)$ so we have $h = -1$ and $k = -2$. The graph also passes through the point $(x, y) = (0, -1)$. Substituting these values for x , y , h , and k , we can solve for a :

$$-1 = a(0 - (-1))^2 + (-2)$$

$$-1 = a(1)^2 - 2$$

$$-1 = a - 2$$

$$1 = a$$

The quadratic function is

$$f(x) = (x+1)^2 - 2 = x^2 + 2x - 1.$$

44. Consider the form $y = a(x-h)^2 + k$. From the graph we know that the vertex is $(2, 1)$ so we have $h = 2$ and $k = 1$. The graph also passes through the point $(x, y) = (0, 5)$. Substituting these values for x , y , h , and k , we can solve for a :

Section 2.4: Properties of Quadratic Functions

$$5 = a(0-2)^2 + 1$$

$$5 = a(-2)^2 + 1$$

$$5 = 4a + 1$$

$$4 = 4a$$

$$1 = a$$

The quadratic function is

$$f(x) = (x-2)^2 + 1 = x^2 - 4x + 5.$$

- 45.** Consider the form $y = a(x-h)^2 + k$. From the graph we know that the vertex is $(-3, 5)$ so we have $h = -3$ and $k = 5$. The graph also passes through the point $(x, y) = (0, -4)$. Substituting these values for x, y, h , and k , we can solve for a :

$$-4 = a(0 - (-3))^2 + 5$$

$$-4 = a(3)^2 + 5$$

$$-4 = 9a + 5$$

$$-9 = 9a$$

$$-1 = a$$

The quadratic function is

$$f(x) = -(x+3)^2 + 5 = -x^2 - 6x - 4.$$

- 46.** Consider the form $y = a(x-h)^2 + k$. From the graph we know that the vertex is $(2, 3)$ so we have $h = 2$ and $k = 3$. The graph also passes through the point $(x, y) = (0, -1)$. Substituting these values for x, y, h , and k , we can solve for a :

$$-1 = a(0-2)^2 + 3$$

$$-1 = a(-2)^2 + 3$$

$$-1 = 4a + 3$$

$$-4 = 4a$$

$$-1 = a$$

The quadratic function is

$$f(x) = -(x-2)^2 + 3 = -x^2 + 4x - 1.$$

- 47.** Consider the form $y = a(x-h)^2 + k$. From the graph we know that the vertex is $(1, -3)$ so we have $h = 1$ and $k = -3$. The graph also passes through the point $(x, y) = (3, 5)$. Substituting these values for x, y, h , and k , we can solve for a :

$$5 = a(3-1)^2 + (-3)$$

$$5 = a(2)^2 - 3$$

$$5 = 4a - 3$$

$$8 = 4a$$

$$2 = a$$

The quadratic function is

$$f(x) = 2(x-1)^2 - 3 = 2x^2 - 4x - 1.$$

- 48.** Consider the form $y = a(x-h)^2 + k$. From the graph we know that the vertex is $(-2, 6)$ so we have $h = -2$ and $k = 6$. The graph also passes through the point $(x, y) = (-4, -2)$. Substituting these values for x, y, h , and k , we can solve for a :

$$-2 = a(-4 - (-2))^2 + 6$$

$$-2 = a(-2)^2 + 6$$

$$-2 = 4a + 6$$

$$-8 = 4a$$

$$-2 = a$$

The quadratic function is

$$f(x) = -2(x+2)^2 + 6 = -2x^2 - 8x - 2.$$

- 49.** For $f(x) = 2x^2 + 12x$, $a = 2$, $b = 12$, $c = 0$.

Since $a = 2 > 0$, the graph opens up, so the vertex is a minimum point. The minimum

$$\text{occurs at } x = \frac{-b}{2a} = \frac{-12}{2(2)} = \frac{-12}{4} = -3.$$

The minimum value is

$$f(-3) = 2(-3)^2 + 12(-3) = 18 - 36 = -18.$$

- 50.** For $f(x) = -2x^2 + 12x$, $a = -2$, $b = 12$, $c = 0$.

Since $a = -2 < 0$, the graph opens down, so the vertex is a maximum point. The maximum

$$\text{occurs at } x = \frac{-b}{2a} = \frac{-12}{2(-2)} = \frac{-12}{-4} = 3.$$

The maximum value is

$$f(3) = -2(3)^2 + 12(3) = -18 + 36 = 18.$$

- 51.** For $f(x) = 2x^2 + 12x - 3$, $a = 2$, $b = 12$, $c = -3$.

Since $a = 2 > 0$, the graph opens up, so the vertex is a minimum point. The minimum occurs at

$$x = \frac{-b}{2a} = \frac{-12}{2(2)} = \frac{-12}{4} = -3. \text{ The minimum value is}$$

$$f(-3) = 2(-3)^2 + 12(-3) - 3 = 18 - 36 - 3 = -21.$$

Chapter 2: Linear and Quadratic Functions

- 52.** For $f(x) = 4x^2 - 8x + 3$, $a = 4$, $b = -8$, $c = 3$.
Since $a = 4 > 0$, the graph opens up, so the vertex is a minimum point. The minimum occurs at

$$x = \frac{-b}{2a} = \frac{-(-8)}{2(4)} = \frac{8}{8} = 1. \text{ The minimum value is}$$

$$f(1) = 4(1)^2 - 8(1) + 3 = 4 - 8 + 3 = -1.$$

- 53.** For $f(x) = -x^2 + 10x - 4$, $a = -1$, $b = 10$, $c = -4$.
Since $a = -1 < 0$, the graph opens down, so the vertex is a maximum point. The maximum occurs

$$\text{at } x = \frac{-b}{2a} = \frac{-10}{2(-1)} = \frac{-10}{-2} = 5. \text{ The maximum}$$

value is

$$f(5) = -(5)^2 + 10(5) - 4 = -25 + 50 - 4 = 21.$$

- 54.** For $f(x) = -2x^2 + 8x + 3$, $a = -2$, $b = 8$, $c = 3$.
Since $a = -2 < 0$, the graph opens down, so the vertex is a maximum point. The maximum

$$\text{occurs at } x = \frac{-b}{2a} = \frac{-8}{2(-2)} = \frac{-8}{-4} = 2. \text{ The}$$

maximum value is

$$f(2) = -2(2)^2 + 8(2) + 3 = -8 + 16 + 3 = 11.$$

- 55.** For $f(x) = -3x^2 + 12x + 1$, $a = -3$, $b = 12$, $c = 1$.
Since $a = -3 < 0$, the graph opens down, so the vertex is a maximum point. The maximum

$$\text{occurs at } x = \frac{-b}{2a} = \frac{-12}{2(-3)} = \frac{-12}{-6} = 2. \text{ The}$$

maximum value is

$$f(2) = -3(2)^2 + 12(2) + 1 = -12 + 24 + 1 = 13.$$

- 56.** For $f(x) = 4x^2 - 4x$, $a = 4$, $b = -4$, $c = 0$.
Since $a = 4 > 0$, the graph opens up, so the vertex is a minimum point. The minimum occurs at

$$x = \frac{-b}{2a} = \frac{-(-4)}{2(4)} = \frac{4}{8} = \frac{1}{2}. \text{ The minimum value is}$$

$$f\left(\frac{1}{2}\right) = 4\left(\frac{1}{2}\right)^2 - 4\left(\frac{1}{2}\right) = 1 - 2 = -1.$$

- 57. a.** For $f(x) = x^2 - 2x - 15$, $a = 1$, $b = -2$, $c = -15$. Since $a = 1 > 0$, the graph opens up. The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-2)}{2(1)} = \frac{2}{2} = 1.$$

The y -coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(1) = (1)^2 - 2(1) - 15 = 1 - 2 - 15 = -16.$$

Thus, the vertex is $(1, -16)$.

The discriminant is:

$$b^2 - 4ac = (-2)^2 - 4(1)(-15) = 4 + 60 = 64 > 0, \text{ so the graph has two } x\text{-intercepts.}$$

The x -intercepts are found by solving:


$$x^2 - 2x - 15 = 0$$

$$(x + 3)(x - 5) = 0$$

$$x = -3 \text{ or } x = 5$$

The x -intercepts are -3 and 5 .

The y -intercept is $f(0) = -15$.



- b.** The domain is $(-\infty, \infty)$.

The range is $[-16, \infty)$.

- c.** Decreasing on $(-\infty, 1)$.

Increasing on $(1, \infty)$.

- 58. a.** For $f(x) = x^2 - 2x - 8$, $a = 1$, $b = -2$, $c = -8$. Since $a = 1 > 0$, the graph opens up. The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-2)}{2(1)} = \frac{2}{2} = 1.$$

The y -coordinate of the vertex is

$$f\left(\frac{-b}{2a}\right) = f(1) = (1)^2 - 2(1) - 8 = 1 - 2 - 8 = -9.$$

Thus, the vertex is $(1, -9)$.

The discriminant is:

$$b^2 - 4ac = (-2)^2 - 4(1)(-8) = 4 + 32 = 36 > 0, \text{ so the graph has two } x\text{-intercepts.}$$

The x -intercepts are found by solving:


$$x^2 - 2x - 8 = 0$$

$$(x + 2)(x - 4) = 0$$

$$x = -2 \text{ or } x = 4$$

Section 2.4: Properties of Quadratic Functions

The x -intercepts are -2 and 4 .
The y -intercept is $f(0) = -8$.




- b. The domain is $(-\infty, \infty)$. The range is $[-9, \infty)$.
- c. Decreasing on $(-\infty, 1)$. Increasing on $(1, \infty)$.

59. a. $F(x) = 2x - 5$ is a linear function.
The x -intercept is found by solving:
 $2x - 5 = 0$
 $2x = 5$
 $x = \frac{5}{2}$

The x -intercept is $\frac{5}{2}$.

The y -intercept is $F(0) = -5$.




- b. The domain is $(-\infty, \infty)$.
The range is $(-\infty, \infty)$.
- c. Increasing on $(-\infty, \infty)$.

60. a. $f(x) = \frac{3}{2}x - 2$ is a linear function.
The x -intercept is found by solving:
 $\frac{3}{2}x - 2 = 0$
 $\frac{3}{2}x = 2$
 $x = 2 \cdot \frac{2}{3} = \frac{4}{3}$

The x -intercept is $\frac{4}{3}$.


The y -intercept is $f(0) = -2$.



- b. The domain is $(-\infty, \infty)$.
The range is $(-\infty, \infty)$.
- c. Increasing on $(-\infty, \infty)$.

61. a. $g(x) = -2(x - 3)^2 + 2$

Using the graph of $y = x^2$, shift right 3 units, reflect about the x -axis, stretch vertically by a factor of 2, then shift up 2 units.




- b. The domain is $(-\infty, \infty)$.
The range is $(-\infty, 2]$.
- c. Increasing on $(-\infty, 3)$.
Decreasing on $(3, \infty)$.

62. a. $h(x) = -3(x + 1)^2 + 4$

Using the graph of $y = x^2$, shift left 1 unit, reflect about the x -axis, stretch vertically by a factor of 3, then shift up 4 units.

Chapter 2: Linear and Quadratic Functions



- b. The domain is $(-\infty, \infty)$.
The range is $(-\infty, 4]$.
- c. Increasing on $(-\infty, -1)$.
Decreasing on $(-1, \infty)$.

63. a. For $f(x) = 2x^2 + x + 1$, $a = 2$, $b = 1$, $c = 1$.

Since $a = 2 > 0$, the graph opens up.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-1}{2(2)} = \frac{-1}{4} = -\frac{1}{4}.$$

The y -coordinate of the vertex is

$$\begin{aligned} f\left(\frac{-b}{2a}\right) &= f\left(-\frac{1}{4}\right) = 2\left(-\frac{1}{4}\right)^2 + \left(-\frac{1}{4}\right) + 1 \\ &= \frac{1}{8} - \frac{1}{4} + 1 = \frac{7}{8}. \end{aligned}$$


Thus, the vertex is $\left(-\frac{1}{4}, \frac{7}{8}\right)$.

The discriminant is:

$$b^2 - 4ac = 1^2 - 4(2)(1) = 1 - 8 = -7,$$

so the graph has no x -intercepts.

The y -intercept is $f(0) = 1$.



- b. The domain is $(-\infty, \infty)$.
The range is $\left[\frac{7}{8}, \infty\right)$.

- c. Decreasing on $\left(-\infty, -\frac{1}{4}\right)$.

Increasing on $\left(-\frac{1}{4}, \infty\right)$.

64. a. For $G(x) = 3x^2 + 2x + 5$, $a = 3$, $b = 2$, $c = 5$.

Since $a = 3 > 0$, the graph opens up.

The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-2}{2(3)} = \frac{-2}{6} = -\frac{1}{3}.$$

The y -coordinate of the vertex is

$$\begin{aligned} G\left(\frac{-b}{2a}\right) &= G\left(-\frac{1}{3}\right) = 3\left(-\frac{1}{3}\right)^2 + 2\left(-\frac{1}{3}\right) + 5 \\ &= \frac{1}{3} - \frac{2}{3} + 5 = \frac{14}{3}. \end{aligned}$$


Thus, the vertex is $\left(-\frac{1}{3}, \frac{14}{3}\right)$.

The discriminant is:

$$b^2 - 4ac = 2^2 - 4(3)(5) = 4 - 60 = -56,$$

so the graph has no x -intercepts.

The y -intercept is $G(0) = 5$.



- b. The domain is $(-\infty, \infty)$.

The range is $\left[\frac{14}{3}, \infty\right)$.

- c. Decreasing on $\left(-\infty, -\frac{1}{3}\right)$.

Increasing on $\left(-\frac{1}{3}, \infty\right)$.

65. a. $h(x) = -\frac{2}{5}x + 4$ is a linear function.

The x -intercept is found by solving:

Section 2.4: Properties of Quadratic Functions


$$-\frac{2}{5}x + 4 = 0$$

$$-\frac{2}{5}x = -4$$

$$x = -4\left(-\frac{5}{2}\right) = 10$$

The x -intercept is 10.

The y -intercept is $h(0) = 4$.



b. The domain is $(-\infty, \infty)$.

The range is $(-\infty, \infty)$.

c. Decreasing on $(-\infty, \infty)$.

66. a. $f(x) = -3x + 2$ is a linear function.

The x -intercept is found by solving:


$$-3x + 2 = 0$$

$$-3x = -2$$

$$x = \frac{-2}{-3} = \frac{2}{3}$$

The x -intercept is $\frac{2}{3}$.

The y -intercept is $f(0) = 2$.



b. The domain is $(-\infty, \infty)$.

The range is $(-\infty, \infty)$.

c. Decreasing on $(-\infty, \infty)$.

67. a. For $H(x) = -4x^2 - 4x - 1$, $a = -4$, $b = -4$, $c = -1$. Since $a = -4 < 0$, the graph opens

down. The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-(-4)}{2(-4)} = \frac{4}{-8} = -\frac{1}{2}.$$

The y -coordinate of the vertex is

$$\begin{aligned} H\left(\frac{-b}{2a}\right) &= H\left(-\frac{1}{2}\right) = -4\left(-\frac{1}{2}\right)^2 - 4\left(-\frac{1}{2}\right) - 1 \\ &= -1 + 2 - 1 = 0 \end{aligned}$$

Thus, the vertex is $\left(-\frac{1}{2}, 0\right)$.

The discriminant is:

$$b^2 - 4ac = (-4)^2 - 4(-4)(-1) = 16 - 16 = 0,$$

so the graph has one x -intercept.

The x -intercept is found by solving:

$$-4x^2 - 4x - 1 = 0$$

$$4x^2 + 4x + 1 = 0$$


$$(2x + 1)^2 = 0$$

$$2x + 1 = 0$$

$$x = -\frac{1}{2}$$

The x -intercept is $-\frac{1}{2}$.

The y -intercept is $H(0) = -1$.



b. The domain is $(-\infty, \infty)$.

The range is $(-\infty, 0]$.

c. Increasing on $\left(-\infty, -\frac{1}{2}\right)$.

Decreasing on $\left(-\frac{1}{2}, \infty\right)$.

68. a. For $F(x) = -4x^2 + 20x - 25$, $a = -4$, $b = 20$, $c = -25$. Since $a = -4 < 0$, the graph opens down. The x -coordinate of the vertex is

$$x = \frac{-b}{2a} = \frac{-20}{2(-4)} = \frac{-20}{-8} = \frac{5}{2}.$$

The y -coordinate of the vertex is

Chapter 2: Linear and Quadratic Functions

$$\begin{aligned} F\left(\frac{-b}{2a}\right) &= F\left(\frac{5}{2}\right) = -4\left(\frac{5}{2}\right)^2 + 20\left(\frac{5}{2}\right) - 25 \\ &= -25 + 50 - 25 = 0 \end{aligned}$$

Thus, the vertex is $\left(\frac{5}{2}, 0\right)$.

The discriminant is:

$$\begin{aligned} b^2 - 4ac &= (20)^2 - 4(-4)(-25) \\ &= 400 - 400 = 0, \end{aligned}$$

so the graph has one x -intercept.

The x -intercept is found by solving:

$$-4x^2 + 20x - 25 = 0$$

$$4x^2 - 20x + 25 = 0$$


$$(2x - 5)^2 = 0$$

$$2x - 5 = 0$$

$$x = \frac{5}{2}$$

The x -intercept is $\frac{5}{2}$.

The y -intercept is $F(0) = -25$.



b. The domain is $(-\infty, \infty)$.

The range is $(-\infty, 0]$.

c. Increasing on $\left(-\infty, \frac{5}{2}\right)$.

Decreasing on $\left(\frac{5}{2}, \infty\right)$.

69. Use the form $f(x) = a(x-h)^2 + k$.

The vertex is $(0, 2)$, so $h = 0$ and $k = 2$.

$$f(x) = a(x-0)^2 + 2 = ax^2 + 2.$$

Since the graph passes through $(1, 8)$, $f(1) = 8$.

$$f(x) = ax^2 + 2$$

$$8 = a(1)^2 + 2$$

$$8 = a + 2$$

$$6 = a$$

$$f(x) = 6x^2 + 2.$$

$$a = 6, b = 0, c = 2$$

70. Use the form $f(x) = a(x-h)^2 + k$.

The vertex is $(1, 4)$, so $h = 1$ and $k = 4$.

$$f(x) = a(x-1)^2 + 4.$$

Since the graph passes through $(-1, -8)$,

$$f(-1) = -8.$$

$$-8 = a(-1-1)^2 + 4$$

$$-8 = a(-2)^2 + 4$$

$$-8 = 4a + 4$$

$$-12 = 4a$$

$$-3 = a$$

$$f(x) = -3(x-1)^2 + 4$$


$$= -3(x^2 - 2x + 1) + 4$$

$$= -3x^2 + 6x - 3 + 4$$

$$= -3x^2 + 6x + 1$$

$$a = -3, b = 6, c = 1$$

71. a and d.



b. $f(x) = g(x)$

$$2x - 1 = x^2 - 4$$

$$0 = x^2 - 2x - 3$$

$$0 = (x+1)(x-3)$$

$$x+1=0 \quad \text{or} \quad x-3=0$$

$$x = -1 \quad x = 3$$

The solution set is $\{-1, 3\}$.

c. $f(-1) = 2(-1) - 1 = -2 - 1 = -3$


$$g(-1) = (-1)^2 - 4 = 1 - 4 = -3$$

$$f(3) = 2(3) - 1 = 6 - 1 = 5$$

$$g(3) = (3)^2 - 4 = 9 - 4 = 5$$

Thus, the graphs of f and g intersect at the points $(-1, -3)$ and $(3, 5)$.

72. a and d.



b. $f(x) = g(x)$

$$-2x - 1 = x^2 - 9$$

$$0 = x^2 + 2x - 8$$

$$0 = (x+4)(x-2)$$

$$x+4=0 \quad \text{or} \quad x-2=0$$

$$x = -4 \quad x = 2$$

The solution set is $\{-4, 2\}$.

c. $f(-4) = -2(-4) - 1 = 8 - 1 = 7$


$$g(-4) = (-4)^2 - 9 = 16 - 9 = 7$$

$$f(2) = -2(2) - 1 = -4 - 1 = -5$$

$$g(2) = (2)^2 - 9 = 4 - 9 = -5$$

Thus, the graphs of f and g intersect at the points $(-4, 7)$ and $(2, -5)$.

73. a and d.



b. $f(x) = g(x)$

$$-x^2 + 4 = -2x + 1$$

$$0 = x^2 - 2x - 3$$

$$0 = (x+1)(x-3)$$

$$x+1=0 \quad \text{or} \quad x-3=0$$

$$x = -1 \quad x = 3$$

The solution set is $\{-1, 3\}$.

c. $f(1) = -(-1)^2 + 4 = -1 + 4 = 3$


$$g(1) = -2(-1) + 1 = 2 + 1 = 3$$

$$f(3) = -(3)^2 + 4 = -9 + 4 = -5$$

$$g(3) = -2(3) + 1 = -6 + 1 = -5$$

Thus, the graphs of f and g intersect at the points $(-1, 3)$ and $(3, -5)$.

74. a and d.



Chapter 2: Linear and Quadratic Functions

b. $f(x) = g(x)$

$$-x^2 + 9 = 2x + 1$$

$$0 = x^2 + 2x - 8$$

$$0 = (x+4)(x-2)$$

$$x+4=0 \quad \text{or} \quad x-2=0$$

$$x=-4 \quad \quad \quad x=2$$

The solution set is $\{-4, 2\}$.

c. $f(-4) = -(-4)^2 + 9 = -16 + 9 = -7$


$$g(-4) = 2(-4) + 1 = -8 + 1 = -7$$

$$f(2) = -(2)^2 + 9 = -4 + 9 = 5$$

$$g(2) = 2(2) + 1 = 4 + 1 = 5$$

Thus, the graphs of f and g intersect at the points $(-4, -7)$ and $(2, 5)$.

75. a and d.



b. $f(x) = g(x)$

$$-x^2 + 5x = x^2 + 3x - 4$$

$$0 = 2x^2 - 2x - 4$$

$$0 = x^2 - x - 2$$

$$0 = (x+1)(x-2)$$

$$x+1=0 \quad \text{or} \quad x-2=0$$

$$x=-1 \quad \quad \quad x=2$$

The solution set is $\{-1, 2\}$.

c. $f(-1) = -(-1)^2 + 5(-1) = -1 - 5 = -6$


$$g(-1) = (-1)^2 + 3(-1) - 4 = 1 - 3 - 4 = -6$$

$$f(2) = -(2)^2 + 5(2) = -4 + 10 = 6$$

$$g(2) = 2^2 + 3(2) - 4 = 4 + 6 - 4 = 6$$

Thus, the graphs of f and g intersect at the points $(-1, -6)$ and $(2, 6)$.

76. a and d.



b. $f(x) = g(x)$

$$-x^2 + 7x - 6 = x^2 + x - 6$$

$$0 = 2x^2 - 6x$$

$$0 = 2x(x-3)$$

$$2x=0 \quad \text{or} \quad x-3=0$$

$$x=0 \quad \quad \quad x=3$$

The solution set is $\{0, 3\}$.

c. $f(0) = -(0)^2 + 7(0) - 6 = -6$

$$g(0) = 0^2 + 0 - 6 = -6$$

$$f(3) = -(3)^2 + 7(3) - 6 = -9 + 21 - 6 = 6$$

$$g(3) = 3^2 + 3 - 6 = 9 + 3 - 6 = 6$$

Thus, the graphs of f and g intersect at the points $(0, -6)$ and $(3, 6)$.

77. a. For $a=1$:

$$f(x) = a(x-r_1)(x-r_2)$$

$$= 1(x-(-3))(x-1)$$

$$= (x+3)(x-1) = x^2 + 2x - 3$$

For $a=2$:

$$f(x) = 2(x-(-3))(x-1)$$

$$= 2(x+3)(x-1)$$

$$= 2(x^2 + 2x - 3) = 2x^2 + 4x - 6$$

For $a=-2$:

$$f(x) = -2(x-(-3))(x-1)$$

$$= -2(x+3)(x-1)$$

$$= -2(x^2 + 2x - 3) = -2x^2 - 4x + 6$$

For $a=5$:

$$f(x) = 5(x-(-3))(x-1)$$

$$= 5(x+3)(x-1)$$

$$= 5(x^2 + 2x - 3) = 5x^2 + 10x - 15$$

Section 2.4: Properties of Quadratic Functions

- b. The x -intercepts are not affected by the value of a . The y -intercept is multiplied by the value of a .
- c. The axis of symmetry is unaffected by the value of a . For this problem, the axis of symmetry is $x = -1$ for all values of a .
- d. The x -coordinate of the vertex is not affected by the value of a . The y -coordinate of the vertex is multiplied by the value of a .
- e. The x -coordinate of the vertex is the mean of the x -intercepts.

78. a. For $a = 1$:

$$f(x) = 1(x - (-5))(x - 3) \\ = (x + 5)(x - 3) = x^2 + 2x - 15$$

For $a = 2$:

$$f(x) = 2(x - (-5))(x - 3) \\ = 2(x + 5)(x - 3) \\ = 2(x^2 + 2x - 15) = 2x^2 + 4x - 30$$

For $a = -2$:

$$f(x) = -2(x - (-5))(x - 3) \\ = -2(x + 5)(x - 3) \\ = -2(x^2 + 2x - 15) = -2x^2 - 4x + 30$$

For $a = 5$:

$$f(x) = 5(x - (-5))(x - 3) \\ = 5(x + 5)(x - 3) \\ = 5(x^2 + 2x - 15) = 5x^2 + 10x - 75$$

- b. The x -intercepts are not affected by the value of a . The y -intercept is multiplied by the value of a .
- c. The axis of symmetry is unaffected by the value of a . For this problem, the axis of symmetry is $x = -1$ for all values of a .
- d. The x -coordinate of the vertex is not affected by the value of a . The y -coordinate of the vertex is multiplied by the value of a .
- e. The x -coordinate of the vertex is the mean of the x -intercepts.
79. a. $x = -\frac{b}{2a} = -\frac{4}{2(1)} = -2$
- $$y = f(-2) = (-2)^2 + 4(-2) - 21 = -25$$
- The vertex is $(-2, -25)$.

b. $f(x) = 0$

$$x^2 + 4x - 21 = 0$$

$$(x + 7)(x - 3) = 0$$

$$x + 7 = 0 \quad \text{or} \quad x - 3 = 0$$

$$x = -7 \quad \quad \quad x = 3$$

The x -intercepts of f are -7 and 3 .

c. $f(x) = -21$

$$x^2 + 4x - 21 = -21$$


$$x^2 + 4x = 0$$

$$x(x + 4) = 0$$

$$x = 0 \quad \text{or} \quad x = -4$$

The solutions $f(x) = -21$ are -4 and 0 .

Thus, the points $(-4, -21)$ and $(0, -21)$ are on the graph of f .



80. a. $x = -\frac{b}{2a} = -\frac{2}{2(1)} = -1$

$$y = f(-1) = (-1)^2 + 2(-1) - 8 = -9$$

The vertex is $(-1, -9)$.

b. $f(x) = 0$

$$x^2 + 2x - 8 = 0$$

$$(x + 4)(x - 2) = 0$$

$$x + 4 = 0 \quad \text{or} \quad x - 2 = 0$$

$$x = -4 \quad \quad \quad x = 2$$

The x -intercepts of f are -4 and 2 .

Chapter 2: Linear and Quadratic Functions

c. $f(x) = -8$

$$x^2 + 2x - 8 = -8$$

$$x^2 + 2x = 0$$

$$x(x+2) = 0$$


$$x = 0 \quad \text{or} \quad x + 2 = 0$$

$$x = -2$$

The solutions $f(x) = -8$ are -2 and 0 .

Thus, the points $(-2, -8)$ and $(0, -8)$ are on the graph of f .

d.



81. $R(p) = -4p^2 + 4000p$, $a = -4$, $b = 4000$, $c = 0$.

Since $a = -4 < 0$ the graph is a parabola that opens down, so the vertex is a maximum point. The

maximum occurs at $p = \frac{-b}{2a} = \frac{-4000}{2(-4)} = 500$.

Thus, the unit price should be \$500 for maximum revenue. The maximum revenue is

$$\begin{aligned} R(500) &= -4(500)^2 + 4000(500) \\ &= -1000000 + 2000000 \\ &= \$1,000,000 \end{aligned}$$

82. $R(p) = -\frac{1}{2}p^2 + 1900p$, $a = -\frac{1}{2}$, $b = 1900$, $c = 0$.

Since $a = -\frac{1}{2} < 0$, the graph is a parabola that opens down, so the vertex is a maximum point.

The maximum occurs at

$$p = \frac{-b}{2a} = \frac{-1900}{2(-1/2)} = \frac{-1900}{-1} = 1900.$$

Thus, the unit price should be \$1900 for maximum revenue. The maximum revenue is

$$\begin{aligned} R(1900) &= -\frac{1}{2}(1900)^2 + 1900(1900) \\ &= -1805000 + 3610000 \\ &= \$1,805,000 \end{aligned}$$

83. a. $C(x) = x^2 - 140x + 7400$,

$$a = 1, b = -140, c = 7400. \text{ Since } a = 1 > 0,$$

the graph opens up, so the vertex is a minimum point. The minimum marginal cost

$$\text{occurs at } x = \frac{-b}{2a} = \frac{-(-140)}{2(1)} = \frac{140}{2} = 70 \text{ mp3}$$

players produced.

b. The minimum marginal cost is

$$\begin{aligned} f\left(\frac{-b}{2a}\right) &= f(70) = (70)^2 - 140(70) + 7400 \\ &= 4900 - 9800 + 7400 \\ &= \$2500 \end{aligned}$$

84. a. $C(x) = 5x^2 - 200x + 4000$,

$$a = 5, b = -200, c = 4000. \text{ Since } a = 5 > 0,$$

the graph opens up, so the vertex is a minimum point. The minimum marginal cost occurs at

$$x = \frac{-b}{2a} = \frac{-(-200)}{2(5)} = \frac{200}{10} = 20 \text{ thousand}$$

cell phones manufactured.

b. The minimum marginal cost is

$$\begin{aligned} f\left(\frac{-b}{2a}\right) &= f(20) = 5(20)^2 - 200(20) + 4000 \\ &= 2000 - 4000 + 4000 \\ &= \$2000 \end{aligned}$$

85. a. $d(v) = 1.1v + 0.06v^2$

$$d(45) = 1.1(45) + 0.06(45)^2$$

$$= 49.5 + 121.5 = 171 \text{ ft.}$$

b. $200 = 1.1v + 0.06v^2$

$$0 = -200 + 1.1v + 0.06v^2$$

$$x = \frac{-(1.1) \pm \sqrt{(1.1)^2 - 4(0.06)(-200)}}{2(0.06)}$$

$$= \frac{-1.1 \pm \sqrt{49.21}}{0.12}$$

$$\approx \frac{-1.1 \pm 7.015}{0.12}$$

$$v \approx 49 \text{ or } v \approx -68$$

Disregard the negative value since we are talking about speed. So the maximum speed you can be traveling would be approximately 49 mph.

Section 2.4: Properties of Quadratic Functions

- c. The 1.1v term might represent the reaction time.
86. a. $a = \frac{-b}{2a} = \frac{-14.23}{2(-0.27)} = \frac{-14.23}{-0.54} \approx 26.4$ years old
- b. $B(26.4) = -0.27(26.4)^2 + 14.23(26.4) - 120.16 \approx 67.3$ births per 1000 unmarried women
- c. $B(40) = -0.27(40)^2 + 14.23(40) - 120.16 \approx 17$ births per 1000 unmarried women
87. a. $R(x) = 75x - 0.2x^2$
 $a = -0.2, b = 75, c = 0$
 The maximum revenue occurs when
 $x = \frac{-b}{2a} = \frac{-75}{2(-0.2)} = \frac{-75}{-0.4} = 187.5$
 The maximum revenue occurs when $x = 187$ or $x = 188$ watches.
 The maximum revenue is:
 $R(187) = 75(187) - 0.2(187)^2 = \7031.20
 $R(188) = 75(188) - 0.2(188)^2 = \7031.20
- b. $P(x) = R(x) - C(x)$
 $= 75x - 0.2x^2 - (32x + 1750)$
 $= -0.2x^2 + 43x - 1750$
- c. $P(x) = -0.2x^2 + 43x - 1750$
 $a = -0.2, b = 43, c = -1750$
 $x = \frac{-b}{2a} = \frac{-43}{2(-0.2)} = \frac{-43}{-0.4} = 107.5$
 The maximum profit occurs when $x = 107$ or $x = 108$ watches.
 The maximum profit is:
 $P(107) = -0.2(107)^2 + 43(107) - 1750 = \561.20
 $P(108) = -0.2(108)^2 + 43(108) - 1750 = \561.20
- d. Answers will vary.
88. a. $R(x) = 9.5x - 0.04x^2$
 $a = -0.04, b = 9.5, c = 0$
 The maximum revenue occurs when

$$x = \frac{-b}{2a} = \frac{-9.5}{2(-0.04)} = \frac{-9.5}{-0.08}$$

$$= 118.75 \approx 119 \text{ boxes of candy}$$

The maximum revenue is:

$$R(119) = 9.5(119) - 0.04(119)^2 = \$564.06$$

- b. $P(x) = R(x) - C(x)$
 $= 9.5x - 0.04x^2 - (1.25x + 250)$
 $= -0.04x^2 + 8.25x - 250$
- c. $P(x) = -0.04x^2 + 8.25x - 250$
 $a = -0.04, b = 8.25, c = -250$
 The maximum profit occurs when
 $x = \frac{-b}{2a} = \frac{-8.25}{2(-0.04)} = \frac{-8.25}{-0.08}$
 $= 103.125 \approx 103$ boxes of candy
 The maximum profit is:
 $P(103) = -0.04(103)^2 + 8.25(103) - 250 = \175.39
- d. Answers will vary.

89. $f(x) = a(x - r_1)(x - r_2)$
 $= a(x + 4)(x - 2)$
 $= ax^2 + 2ax - 8a$
 The x value of the vertex is $x = \frac{-b}{2a} = \frac{-2a}{2a} = -1$.
 The y value of the vertex is 18.
 $-18 = a(-1)^2 + 2a(-1) - 8a$
 $-18 = -9a$
 $a = 2$

So the function is $f(x) = 2(x + 4)(x - 2)$

90. $f(x) = a(x - r_1)(x - r_2)$
 $= a(x + 1)(x - 5)$
 $= ax^2 - 4ax - 5a$
 The x value of the vertex is
 $x = \frac{-b}{2a} = \frac{-(-4a)}{2a} = 2$.
 The y value of the vertex is 9.
 $9 = a(2)^2 - 4a(2) - 5a$
 $9 = -9a$
 $a = -1$


So the function is $f(x) = -(x + 1)(x - 5)$

Chapter 2: Linear and Quadratic Functions

91. If x is even, then ax^2 and bx are even. When two even numbers are added to an odd number the result is odd. Thus, $f(x)$ is odd. If x is odd, then ax^2 and bx are odd. The sum of three odd numbers is an odd number. Thus, $f(x)$ is odd.

92. Answers will vary.


93. $y = x^2 + 2x - 3$; $y = x^2 + 2x + 1$; $y = x^2 + 2x$



Each member of this family will be a parabola with the following characteristics:

- (i) opens upwards since $a > 0$;
- (ii) vertex occurs at $x = -\frac{b}{2a} = -\frac{2}{2(1)} = -1$;
- (iii) There is at least one x -intercept since $b^2 - 4ac \geq 0$.

94. $y = x^2 - 4x + 1$; $y = x^2 + 1$; $y = x^2 + 4x + 1$



Each member of this family will be a parabola with the following characteristics:

- (i) opens upwards since $a > 0$
- (ii) y -intercept occurs at $(0, 1)$.

95. The graph of the quadratic function $f(x) = ax^2 + bx + c$ will not have any x -intercepts whenever $b^2 - 4ac < 0$.

96. By completing the square on the quadratic function $f(x) = ax^2 + bx + c$ we obtain the equation $y = a\left(x + \frac{b}{2a}\right)^2 + c - \frac{b^2}{4a}$. We can then draw the graph by applying transformations to the graph of the basic parabola $y = x^2$, which opens up. When $a > 0$, the basic parabola will either be stretched or compressed vertically. When $a < 0$, the basic parabola will either be stretched or compressed vertically as well as reflected across the x -axis. Therefore, when $a > 0$, the graph of $f(x) = ax^2 + bx + c$ will open up, and when $a < 0$, the graph of $f(x) = ax^2 + bx + c$ will open down.

97. No. We know that the graph of a quadratic function $f(x) = ax^2 + bx + c$ is a parabola with vertex $\left(-\frac{b}{2a}, f\left(-\frac{b}{2a}\right)\right)$. If $a > 0$, then the vertex is a minimum point, so the range is $\left[f\left(-\frac{b}{2a}\right), \infty\right)$. If $a < 0$, then the vertex is a maximum point, so the range is $\left(-\infty, f\left(-\frac{b}{2a}\right)\right]$. Therefore, it is impossible for the range to be $(-\infty, \infty)$.

98. Two quadratic functions can intersect 0, 1, or 2 times.

Section 2.5


1. $-3x - 2 < 7$

$$-3x < 9$$

$$x > -3$$

The solution set is $\{x \mid x > -3\}$ or $(-3, \infty)$.

2. $(-2, 7]$ represents the numbers between -2 and 7 , including 7 but not including -2 . Using inequality notation, this is written as $-2 < x \leq 7$.



3. a. $f(x) > 0$ when the graph of f is above the x -axis. Thus, $\{x \mid x < -2 \text{ or } x > 2\}$ or, using interval notation, $(-\infty, -2) \cup (2, \infty)$.

- b. $f(x) \leq 0$ when the graph of f is below or intersects the x -axis. Thus, $\{x \mid -2 \leq x \leq 2\}$ or, using interval notation, $[-2, 2]$.

4. a. $g(x) < 0$ when the graph of g is below the x -axis. Thus, $\{x \mid x < -1 \text{ or } x > 4\}$ or, using interval notation, $(-\infty, -1) \cup (4, \infty)$.

- b. $g(x) \geq 0$ when the graph of f is above or intersects the x -axis. Thus, $\{x \mid -1 \leq x \leq 4\}$ or, using interval notation, $[-1, 4]$.

5. a. $g(x) \geq f(x)$ when the graph of g is above or intersects the graph of f . Thus $\{x \mid -2 \leq x \leq 1\}$ or, using interval notation, $[-2, 1]$.

- b. $f(x) > g(x)$ when the graph of f is above the graph of g . Thus, $\{x \mid x < -2 \text{ or } x > 1\}$ or, using interval notation, $(-\infty, -2) \cup (1, \infty)$.

6. a. $f(x) < g(x)$ when the graph of f is below the graph of g . Thus, $\{x \mid x < -3 \text{ or } x > 1\}$ or, using interval notation, $(-\infty, -3) \cup (1, \infty)$.

- b. $f(x) \geq g(x)$ when the graph of f is above or intersects the graph of g . Thus, $\{x \mid -3 \leq x \leq 1\}$ or, using interval notation, $[-3, 1]$.

7. $x^2 - 3x - 10 \leq 0$

We graph the function $f(x) = x^2 - 3x - 10$. The intercepts are

y-intercept: $f(0) = -10$


x-intercepts: $x^2 - 3x - 10 = 0$

$$(x - 5)(x + 2) = 0$$

$$x = 5, x = -2$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-3)}{2(1)} = \frac{3}{2}$. Since

$$f\left(\frac{3}{2}\right) = -\frac{49}{4}, \text{ the vertex is } \left(\frac{3}{2}, -\frac{49}{4}\right).$$



The graph is below the x -axis for $-2 < x < 5$. Since the inequality is not strict, the solution set is $\{x \mid -2 \leq x \leq 5\}$ or, using interval notation, $[-2, 5]$.

8. $x^2 + 3x - 10 > 0$

We graph the function $f(x) = x^2 + 3x - 10$. The intercepts are

y-intercept: $f(0) = -10$


x-intercepts: $x^2 + 3x - 10 = 0$

$$(x + 5)(x - 2) = 0$$

$$x = -5, x = 2$$

The vertex is at $x = \frac{-b}{2a} = \frac{-3}{2(1)} = -\frac{3}{2}$. Since

$$f\left(-\frac{3}{2}\right) = -\frac{49}{4}, \text{ the vertex is } \left(-\frac{3}{2}, -\frac{49}{4}\right).$$



The graph is above the x -axis when $x < -5$ or $x > 2$. Since the inequality is strict, the solution set is $\{x \mid x < -5 \text{ or } x > 2\}$ or, using interval notation, $(-\infty, -5) \cup (2, \infty)$.

Chapter 2: Linear and Quadratic Functions

9. $x^2 - 4x > 0$

We graph the function $f(x) = x^2 - 4x$. The intercepts are

y-intercept: $f(0) = 0$


x-intercepts: $x^2 - 4x = 0$

$$x(x - 4) = 0$$

$$x = 0, x = 4$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-4)}{2(1)} = \frac{4}{2} = 2$. Since

$f(2) = -4$, the vertex is $(2, -4)$.



The graph is above the x -axis when $x < 0$ or $x > 4$. Since the inequality is strict, the solution set is $\{x \mid x < 0 \text{ or } x > 4\}$ or, using interval notation, $(-\infty, 0) \cup (4, \infty)$.

10. $x^2 + 8x \leq 0$

We graph the function $f(x) = x^2 + 8x$. The intercepts are

y-intercept: $f(0) = 0$


x-intercepts: $x^2 + 8x = 0$

$$x(x + 8) = 0$$

$$x = 0, x = -8$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(8)}{2(1)} = \frac{-8}{2} = -4$.

Since $f(-4) = -16$, the vertex is $(-4, -16)$.



The graph is below the x -axis when $-8 < x < 0$. Since the inequality is not strict, the solution set is $\{x \mid -8 \leq x \leq 0\}$ or, using interval notation, $[-8, 0]$.

11. $x^2 + x > 12$

$$x^2 + x - 12 > 0$$

We graph the function $f(x) = x^2 + x - 12$.

y-intercept: $f(0) = -12$


x-intercepts: $x^2 + x - 12 = 0$

$$(x + 4)(x - 3) = 0$$

$$x = -4, x = 3$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(1)}{2(1)} = -\frac{1}{2}$. Since

$f\left(-\frac{1}{2}\right) = -\frac{49}{4}$, the vertex is $\left(-\frac{1}{2}, -\frac{49}{4}\right)$.



The graph is above the x -axis when $x < -4$ or $x > 3$. Since the inequality is strict, the solution set is $\{x \mid x < -4 \text{ or } x > 3\}$ or, using interval notation, $(-\infty, -4) \cup (3, \infty)$.

12. $x^2 + 7x < -12$

$$x^2 + 7x + 12 < 0$$

We graph the function $f(x) = x^2 + 7x + 12$.

y-intercept: $f(0) = 12$


x-intercepts: $x^2 + 7x + 12 = 0$

$$(x + 4)(x + 3) = 0$$

$$x = -4, x = -3$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(7)}{2(1)} = -\frac{7}{2}$. Since

$f\left(-\frac{7}{2}\right) = -\frac{1}{4}$, the vertex is $\left(-\frac{7}{2}, -\frac{1}{4}\right)$.



The graph is below the x -axis when $-4 < x < -3$. Since the inequality is strict, the solution set is $\{x \mid -4 < x < -3\}$ or, using interval notation, $(-4, -3)$.

Section 2.5: Inequalities Involving Quadratic Functions

13. $x^2 + 6x + 9 \leq 0$

We graph the function $f(x) = x^2 + 6x + 9$.

y-intercept: $f(0) = 9$


x-intercepts: $x^2 + 6x + 9 = 0$

$$(x+3)(x+3) = 0$$

$$x = -3$$

The vertex is at $x = \frac{-b}{2a} = \frac{-6}{2(1)} = -3$. Since

$f(-3) = 0$, the vertex is $(-3, 0)$.



Since the graph is never below the x -axis and only touches at $x = -3$ then the solution is $\{-3\}$.

14. $x^2 - 4x + 4 \leq 0$

We graph the function $f(x) = x^2 - 4x + 4$.

y-intercept: $f(0) = 4$


x-intercepts: $x^2 - 4x + 4 = 0$

$$(x-2)(x-2) = 0$$

$$x = 2$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-4)}{2(1)} = 2$. Since

$f(2) = 0$, the vertex is $(2, 0)$.



Since the graph is never below the x -axis and only touches at $x = 2$ then the solution is $\{2\}$.

15. $2x^2 \leq 5x + 3$

$$2x^2 - 5x - 3 \leq 0$$

We graph the function $f(x) = 2x^2 - 5x - 3$. The intercepts are

y-intercept: $f(0) = -3$


x-intercepts: $2x^2 - 5x - 3 = 0$

$$(2x+1)(x-3) = 0$$

$$x = -\frac{1}{2}, x = 3$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-5)}{2(2)} = \frac{5}{4}$. Since

$f\left(\frac{5}{4}\right) = -\frac{49}{8}$, the vertex is $\left(\frac{5}{4}, -\frac{49}{8}\right)$.



The graph is below the x -axis when $-\frac{1}{2} < x < 3$.

Since the inequality is not strict, the solution set is $\left\{x \mid -\frac{1}{2} \leq x \leq 3\right\}$ or, using interval notation,

$$\left[-\frac{1}{2}, 3\right].$$

16. $6x^2 \leq 6 + 5x$

$$6x^2 - 5x - 6 \leq 0$$

We graph the function $f(x) = 6x^2 - 5x - 6$. The intercepts are

y-intercept: $f(0) = -6$


x-intercepts: $6x^2 - 5x - 6 = 0$

$$(3x+2)(2x-3) = 0$$

$$x = -\frac{2}{3}, x = \frac{3}{2}$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-5)}{2(6)} = \frac{5}{12}$. Since

$f\left(\frac{5}{12}\right) = -\frac{169}{24}$, the vertex is $\left(\frac{5}{12}, -\frac{169}{24}\right)$.



The graph is below the x -axis when $-\frac{2}{3} < x < \frac{3}{2}$.

Since the inequality is not strict, the solution set

Chapter 2: Linear and Quadratic Functions

is $\left\{x \mid -\frac{2}{3} \leq x \leq \frac{3}{2}\right\}$ or, using interval notation,
 $\left[-\frac{2}{3}, \frac{3}{2}\right]$.

17. $x(x-7) > 8$

$$x^2 - 7x > 8$$

$$x^2 - 7x - 8 > 0$$

We graph the function $f(x) = x^2 - 7x - 8$. The intercepts are

y-intercept: $f(0) = -8$


x-intercepts: $x^2 - 7x - 8 = 0$

$$(x+1)(x-8) = 0$$

$$x = -1, x = 8$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-7)}{2(1)} = \frac{7}{2}$. Since

$$f\left(\frac{7}{2}\right) = -\frac{81}{4}, \text{ the vertex is } \left(\frac{7}{2}, -\frac{81}{4}\right).$$



The graph is above the x -axis when $x < -1$ or $x > 8$. Since the inequality is strict, the solution set is $\{x \mid x < -1 \text{ or } x > 8\}$ or, using interval notation, $(-\infty, -1) \cup (8, \infty)$.

18. $x(x+1) > 20$

$$x^2 + x > 20$$

$$x^2 + x - 20 > 0$$

We graph the function $f(x) = x^2 + x - 20$.

y-intercept: $f(0) = -20$


x-intercepts: $x^2 + x - 20 = 0$

$$(x+5)(x-4) = 0$$

$$x = -5, x = 4$$

The vertex is at $x = \frac{-b}{2a} = \frac{-1}{2(1)} = -\frac{1}{2}$. Since

$$f\left(-\frac{1}{2}\right) = -\frac{81}{4}, \text{ the vertex is } \left(-\frac{1}{2}, -\frac{81}{4}\right).$$



The graph is above the x -axis when $x < -5$ or $x > 4$. Since the inequality is strict, the solution set is $\{x \mid x < -5 \text{ or } x > 4\}$ or, using interval notation, $(-\infty, -5) \cup (4, \infty)$.

19. $4x^2 + 9 < 6x$

$$4x^2 - 6x + 9 < 0$$

We graph the function $f(x) = 4x^2 - 6x + 9$.


y-intercept: $f(0) = 9$

$$\begin{aligned} \text{x-intercepts: } x &= \frac{-(-6) \pm \sqrt{(-6)^2 - 4(4)(9)}}{2(4)} \\ &= \frac{6 \pm \sqrt{-108}}{8} \quad (\text{not real}) \end{aligned}$$

Therefore, f has no x -intercepts.

The vertex is at $x = \frac{-b}{2a} = \frac{-(-6)}{2(4)} = \frac{6}{8} = \frac{3}{4}$. Since

$$f\left(\frac{3}{4}\right) = \frac{27}{4}, \text{ the vertex is } \left(\frac{3}{4}, \frac{27}{4}\right).$$



The graph is never below the x -axis. Thus, there is no real solution.

20. $25x^2 + 16 < 40x$

$$25x^2 - 40x + 16 < 0$$

We graph the function $f(x) = 25x^2 - 40x + 16$.

y-intercept: $f(0) = 16$

x-intercepts: $25x^2 - 40x + 16 = 0$

$$(5x-4)^2 = 0$$


$$5x - 4 = 0$$

$$x = \frac{4}{5}$$

Section 2.5: Inequalities Involving Quadratic Functions

The vertex is at $x = \frac{-b}{2a} = \frac{-(-40)}{2(25)} = \frac{40}{50} = \frac{4}{5}$.

Since $f\left(\frac{4}{5}\right) = 0$, the vertex is $\left(\frac{4}{5}, 0\right)$.



The graph is never below the x -axis. Thus, there is no real solution.

21. $6(x^2 + 1) > 5x$

$$6x^2 + 6 > 5x$$

$$6x^2 - 5x + 6 > 0$$

We graph the function $f(x) = 6x^2 - 5x + 6$.

y -intercept: $f(0) = 6$


$$x\text{-intercepts: } x = \frac{-(-5) \pm \sqrt{(-5)^2 - 4(6)(6)}}{2(6)}$$

$$= \frac{6 \pm \sqrt{-119}}{12} \quad (\text{not real})$$

Therefore, f has no x -intercepts.

The vertex is at $x = \frac{-b}{2a} = \frac{-(-5)}{2(6)} = \frac{5}{12}$. Since

$f\left(\frac{5}{12}\right) = \frac{119}{24}$, the vertex is $\left(\frac{5}{12}, \frac{119}{24}\right)$.



The graph is always above the x -axis. Thus, the solution set is all real numbers or, using interval notation, $(-\infty, \infty)$.

22. $2(2x^2 - 3x) > -9$

$$4x^2 - 6x > -9$$

$$4x^2 - 6x + 9 > 0$$

We graph the function $f(x) = 4x^2 - 6x + 9$.


y -intercept: $f(0) = 9$

$$\begin{aligned} x\text{-intercepts: } x &= \frac{-(-6) \pm \sqrt{(-6)^2 - 4(4)(9)}}{2(4)} \\ &= \frac{6 \pm \sqrt{-108}}{8} \quad (\text{not real}) \end{aligned}$$

Therefore, f has no x -intercepts.

The vertex is at $x = \frac{-b}{2a} = \frac{-(-6)}{2(4)} = \frac{6}{8} = \frac{3}{4}$. Since

$f\left(\frac{3}{4}\right) = \frac{27}{4}$, the vertex is $\left(\frac{3}{4}, \frac{27}{4}\right)$.



The graph is always above the x -axis. Thus, the solution set is all real numbers or, using interval notation, $(-\infty, \infty)$.

23. The domain of the expression $f(x) = \sqrt{x^2 - 16}$ includes all values for which $x^2 - 16 \geq 0$.

We graph the function $p(x) = x^2 - 16$. The intercepts of p are

y -intercept: $p(0) = -16$


x -intercepts: $x^2 - 16 = 0$

$$(x+4)(x-4) = 0$$

$$x = -4, x = 4$$

The vertex of p is at $x = \frac{-b}{2a} = \frac{-(0)}{2(1)} = 0$. Since

$p(0) = -16$, the vertex is $(0, -16)$.



The graph of p is above the x -axis when $x < -4$ or $x > 4$. Since the inequality is not strict, the solution set of $x^2 - 16 \geq 0$ is $\{x \mid x \leq -4 \text{ or } x \geq 4\}$. Thus, the domain of f is also $\{x \mid x \leq -4 \text{ or } x \geq 4\}$ or, using interval notation, $(-\infty, -4] \cup [4, \infty)$.

24. The domain of the expression $f(x) = \sqrt{x - 3x^2}$ includes all values for which $x - 3x^2 \geq 0$.

Chapter 2: Linear and Quadratic Functions

We graph the function $p(x) = x - 3x^2$. The intercepts of p are

y-intercept: $p(0) = -6$


x-intercepts: $x - 3x^2 = 0$

$$x(1 - 3x) = 0$$

$$x = 0, x = \frac{1}{3}.$$

The vertex of p is at $x = \frac{-b}{2a} = \frac{-(-1)}{2(-3)} = \frac{-1}{-6} = \frac{1}{6}$.

Since $p\left(\frac{1}{6}\right) = \frac{1}{12}$, the vertex is $\left(\frac{1}{6}, \frac{1}{12}\right)$.



The graph of p is above the x -axis when

$0 < x < \frac{1}{3}$. Since the inequality is not strict, the

solution set of $x - 3x^2 \geq 0$ is $\left\{x \mid 0 \leq x \leq \frac{1}{3}\right\}$.

Thus, the domain of f is also $\left\{x \mid 0 \leq x \leq \frac{1}{3}\right\}$ or,

using interval notation, $\left[0, \frac{1}{3}\right]$.

25. $f(x) = x^2 - 1$; $g(x) = 3x + 3$

a. $f(x) = 0$

$$x^2 - 1 = 0$$

$$(x - 1)(x + 1) = 0$$

$$x = 1; x = -1$$

Solution set: $\{-1, 1\}$.

b. $g(x) = 0$

$$3x + 3 = 0$$

$$3x = -3$$

$$x = -1$$

Solution set: $\{-1\}$.

c. $f(x) = g(x)$

$$x^2 - 1 = 3x + 3$$

$$x^2 - 3x - 4 = 0$$

$$(x - 4)(x + 1) = 0$$

$$x = 4; x = -1$$

Solution set: $\{-1, 4\}$.

d. $f(x) > 0$

We graph the function $f(x) = x^2 - 1$.

y-intercept: $f(0) = -1$


x-intercepts: $x^2 - 1 = 0$

$$(x + 1)(x - 1) = 0$$

$$x = -1, x = 1$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-0)}{2(1)} = 0$. Since

$f(0) = -1$, the vertex is $(0, -1)$.



The graph is above the x -axis when $x < -1$ or $x > 1$. Since the inequality is strict, the solution set is $\{x \mid x < -1 \text{ or } x > 1\}$ or, using interval notation, $(-\infty, -1) \cup (1, \infty)$.

e. $g(x) \leq 0$

$$3x + 3 \leq 0$$

$$3x \leq -3$$

$$x \leq -1$$

The solution set is $\{x \mid x \leq -1\}$ or, using interval notation, $(-\infty, -1]$.

f. $f(x) > g(x)$

$$x^2 - 1 > 3x + 3$$

$$x^2 - 3x - 4 > 0$$

We graph the function $p(x) = x^2 - 3x - 4$.

The intercepts of p are

y-intercept: $p(0) = -4$

x-intercepts: $x^2 - 3x - 4 = 0$


$$(x - 4)(x + 1) = 0$$

$$x = 4, x = -1$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-3)}{2(1)} = \frac{3}{2}$. Since

$p\left(\frac{3}{2}\right) = -\frac{25}{4}$, the vertex is $\left(\frac{3}{2}, -\frac{25}{4}\right)$.

Section 2.5: Inequalities Involving Quadratic Functions



The graph of p is above the x -axis when $x < -1$ or $x > 4$. Since the inequality is strict, the solution set is

$\{x \mid x < -1 \text{ or } x > 4\}$ or, using interval notation, $(-\infty, -1) \cup (4, \infty)$.

g. $f(x) \geq 1$
 $x^2 - 1 \geq 1$
 $x^2 - 2 \geq 0$

We graph the function $p(x) = x^2 - 2$. The intercepts of p are

y-intercept: $p(0) = -2$


x-intercepts: $x^2 - 2 = 0$

$$x^2 = 2$$

$$x = \pm\sqrt{2}$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(1)} = 0$. Since

$p(0) = -2$, the vertex is $(0, -2)$.



The graph of p is above the x -axis when $x < -\sqrt{2}$ or $x > \sqrt{2}$. Since the inequality is not strict, the solution set is

$\{x \mid x \leq -\sqrt{2} \text{ or } x \geq \sqrt{2}\}$ or, using interval notation, $(-\infty, -\sqrt{2}] \cup [\sqrt{2}, \infty)$.

26. $f(x) = -x^2 + 3$; $g(x) = -3x + 3$

a. $f(x) = 0$
 $-x^2 + 3 = 0$
 $x^2 = 3$
 $x = \pm\sqrt{3}$

Solution set: $\{-\sqrt{3}, \sqrt{3}\}$.

b. $g(x) = 0$
 $-3x + 3 = 0$
 $-3x = -3$
 $x = 1$

Solution set: $\{1\}$.

c. $f(x) = g(x)$
 $-x^2 + 3 = -3x + 3$
 $0 = x^2 - 3x$
 $0 = x(x - 3)$
 $x = 0; x = 3$

Solution set: $\{0, 3\}$.

d. $f(x) > 0$

We graph the function $f(x) = -x^2 + 3$.

y-intercept: $f(0) = 3$


x-intercepts: $-x^2 + 3 = 0$

$$x^2 = 3$$

$$x = \pm\sqrt{3}$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(-1)} = 0$. Since

$f(0) = 3$, the vertex is $(0, 3)$.



The graph is above the x -axis when

$-\sqrt{3} < x < \sqrt{3}$. Since the inequality is strict, the solution set is $\{x \mid -\sqrt{3} < x < \sqrt{3}\}$ or, using interval notation, $(-\sqrt{3}, \sqrt{3})$.

e. $g(x) \leq 0$
 $-3x + 3 \leq 0$
 $-3x \leq -3$
 $x \geq 1$

The solution set is $\{x \mid x \geq 1\}$ or, using interval notation, $[1, \infty)$.

Chapter 2: Linear and Quadratic Functions

f. $f(x) > g(x)$
 $-x^2 + 3 > -3x + 3$
 $-x^2 + 3x > 0$

We graph the function $p(x) = -x^2 + 3x$.

The intercepts of p are

y-intercept: $p(0) = 0$


x-intercepts: $-x^2 + 3x = 0$

$$-x(x-3) = 0$$

$$x = 0; x = 3$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-3)}{2(-1)} = \frac{-3}{-2} = \frac{3}{2}$.

Since $p\left(\frac{3}{2}\right) = \frac{9}{4}$, the vertex is $\left(\frac{3}{2}, \frac{9}{4}\right)$.



The graph of p is above the x -axis when $0 < x < 3$. Since the inequality is strict, the solution set is $\{x \mid 0 < x < 3\}$ or, using interval notation, $(0, 3)$.

g. $f(x) \geq 1$
 $-x^2 + 3 \geq 1$
 $-x^2 + 2 \geq 0$

We graph the function $p(x) = -x^2 + 2$. The

intercepts of p are

y-intercept: $p(0) = 2$


x-intercepts: $-x^2 + 2 = 0$

$$x^2 = 2$$

$$x = \pm\sqrt{2}$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(-1)} = 0$. Since

$p(0) = 2$, the vertex is $(0, 2)$.



The graph of p is above the x -axis when $-\sqrt{2} < x < \sqrt{2}$. Since the inequality is not

strict, the solution set is $\{x \mid -\sqrt{2} \leq x \leq \sqrt{2}\}$

or, using interval notation, $[-\sqrt{2}, \sqrt{2}]$.

27. $f(x) = -x^2 + 1$; $g(x) = 4x + 1$

a. $f(x) = 0$

$$-x^2 + 1 = 0$$

$$1 - x^2 = 0$$

$$(1-x)(1+x) = 0$$

$$x = 1; x = -1$$

Solution set: $\{-1, 1\}$.

b. $g(x) = 0$

$$4x + 1 = 0$$

$$4x = -1$$

$$x = -\frac{1}{4}$$

Solution set: $\left\{-\frac{1}{4}\right\}$.

c. $f(x) = g(x)$

$$-x^2 + 1 = 4x + 1$$

$$0 = x^2 + 4x$$

$$0 = x(x+4)$$

$$x = 0; x = -4$$

Solution set: $\{-4, 0\}$.

d. $f(x) > 0$

We graph the function $f(x) = -x^2 + 1$.

y-intercept: $f(0) = 1$

x-intercepts: $-x^2 + 1 = 0$


$$x^2 - 1 = 0$$

$$(x+1)(x-1) = 0$$

$$x = -1; x = 1$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(-1)} = 0$. Since

$f(0) = 1$, the vertex is $(0, 1)$.



The graph is above the x -axis when $-1 < x < 1$. Since the inequality is strict, the

Section 2.5: Inequalities Involving Quadratic Functions

solution set is $\{x \mid -1 < x < 1\}$ or, using interval notation, $(-1, 1)$.

e. $g(x) \leq 0$

$$4x + 1 \leq 0$$

$$4x \leq -1$$

$$x \leq -\frac{1}{4}$$

The solution set is $\left\{x \mid x \leq -\frac{1}{4}\right\}$ or, using

interval notation, $\left(-\infty, -\frac{1}{4}\right]$.

f. $f(x) > g(x)$

$$-x^2 + 1 > 4x + 1$$

$$-x^2 - 4x > 0$$

We graph the function $p(x) = -x^2 - 4x$.

The intercepts of p are

y-intercept: $p(0) = 0$


x-intercepts: $-x^2 - 4x = 0$

$$-x(x + 4) = 0$$

$$x = 0; x = -4$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-4)}{2(-1)} = \frac{4}{-2} = -2$.

Since $p(-2) = 4$, the vertex is $(-2, 4)$.



The graph of p is above the x -axis when $-4 < x < 0$. Since the inequality is strict, the solution set is $\{x \mid -4 < x < 0\}$ or, using interval notation, $(-4, 0)$.

g. $f(x) \geq 1$

$$-x^2 + 1 \geq 1$$


$$-x^2 \geq 0$$

We graph the function $p(x) = -x^2$. The

vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(-1)} = 0$. Since

$p(0) = 0$, the vertex is $(0, 0)$. Since

$a = -1 < 0$, the parabola opens downward.



The graph of p is never above the x -axis, but it does touch the x -axis at $x = 0$. Since the inequality is not strict, the solution set is $\{0\}$.

28. $f(x) = -x^2 + 4$; $g(x) = -x - 2$

a. $f(x) = 0$

$$-x^2 + 4 = 0$$

$$x^2 - 4 = 0$$

$$(x + 2)(x - 2) = 0$$

$$x = -2; x = 2$$

Solution set: $\{-2, 2\}$.

b. $g(x) = 0$

$$-x - 2 = 0$$

$$-2 = x$$

Solution set: $\{-2\}$.

c. $f(x) = g(x)$

$$-x^2 + 4 = -x - 2$$

$$0 = x^2 - x - 6$$

$$0 = (x - 3)(x + 2)$$

$$x = 3; x = -2$$

Solution set: $\{-2, 3\}$.

d. $f(x) > 0$

$$-x^2 + 4 > 0$$

We graph the function $f(x) = -x^2 + 4$.

y-intercept: $f(0) = 4$

x-intercepts: $-x^2 + 4 = 0$


$$x^2 - 4 = 0$$

$$(x + 2)(x - 2) = 0$$

$$x = -2; x = 2$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(-1)} = 0$. Since

$f(0) = 4$, the vertex is $(0, 4)$.



Chapter 2: Linear and Quadratic Functions

The graph is above the x -axis when $-2 < x < 2$. Since the inequality is strict, the solution set is $\{x \mid -2 < x < 2\}$ or, using interval notation, $(-2, 2)$.

e. $g(x) \leq 0$

$$-x - 2 \leq 0$$

$$-x \leq 2$$

$$x \geq -2$$

The solution set is $\{x \mid x \geq -2\}$ or, using interval notation, $[-2, \infty)$.

f. $f(x) > g(x)$

$$-x^2 + 4 > -x - 2$$

$$-x^2 + x + 6 > 0$$

We graph the function $p(x) = -x^2 + x + 6$.

The intercepts of p are

y-intercept: $p(0) = 6$

x-intercepts: $-x^2 + x + 6 = 0$


$$x^2 - x - 6 = 0$$

$$(x+2)(x-3) = 0$$

$$x = -2; x = 3$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(1)}{2(-1)} = \frac{-1}{-2} = \frac{1}{2}$.

Since $p\left(\frac{1}{2}\right) = \frac{25}{4}$, the vertex is $\left(\frac{1}{2}, \frac{25}{4}\right)$.



The graph of p is above the x -axis when $-2 < x < 3$. Since the inequality is strict, the solution set is $\{x \mid -2 < x < 3\}$ or, using interval notation, $(-2, 3)$.

g. $f(x) \geq 1$

$$-x^2 + 4 > 1$$

$$-x^2 + 3 > 0$$

We graph the function $p(x) = -x^2 + 3$. The

intercepts of p are

y-intercept: $p(0) = 3$


x-intercepts: $-x^2 + 3 = 0$

$$x^2 = 3$$

$$x = \pm\sqrt{3}$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(-1)} = 0$. Since

$p(0) = 3$, the vertex is $(0, 3)$.



The graph of p is above the x -axis when $-\sqrt{3} < x < \sqrt{3}$. Since the inequality is not strict, the solution set is $\{x \mid -\sqrt{3} \leq x \leq \sqrt{3}\}$ or, using interval notation, $[-\sqrt{3}, \sqrt{3}]$.

29. $f(x) = x^2 - 4$; $g(x) = -x^2 + 4$

a. $f(x) = 0$

$$x^2 - 4 = 0$$

$$(x-2)(x+2) = 0$$

$$x = 2; x = -2$$

Solution set: $\{-2, 2\}$.

b. $g(x) = 0$

$$-x^2 + 4 = 0$$

$$x^2 - 4 = 0$$

$$(x+2)(x-2) = 0$$

$$x = -2; x = 2$$

Solution set: $\{-2, 2\}$.

c. $f(x) = g(x)$

$$x^2 - 4 = -x^2 + 4$$

$$2x^2 - 8 = 0$$

$$2(x-2)(x+2) = 0$$

$$x = 2; x = -2$$

Solution set: $\{-2, 2\}$.

d. $f(x) > 0$

$$x^2 - 4 > 0$$


We graph the function $f(x) = x^2 - 4$.

y-intercept: $f(0) = -4$

Section 2.5: Inequalities Involving Quadratic Functions

$$\begin{aligned}x\text{-intercepts: } x^2 - 4 &= 0 \\(x+2)(x-2) &= 0 \\x &= -2; x = 2\end{aligned}$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(-1)} = 0$. Since $f(0) = -4$, the vertex is $(0, -4)$.



The graph is above the x -axis when $x < -2$ or $x > 2$. Since the inequality is strict, the solution set is $\{x \mid x < -2 \text{ or } x > 2\}$ or, using interval notation, $(-\infty, -2) \cup (2, \infty)$.

e. $g(x) \leq 0$


$$-x^2 + 4 \leq 0$$

We graph the function $g(x) = -x^2 + 4$.

y-intercept: $g(0) = 4$

$$\begin{aligned}x\text{-intercepts: } -x^2 + 4 &= 0 \\x^2 - 4 &= 0 \\(x+2)(x-2) &= 0 \\x &= -2; x = 2\end{aligned}$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(-1)} = 0$. Since $g(0) = 4$, the vertex is $(0, 4)$.



The graph is below the x -axis when $x < -2$ or $x > 2$. Since the inequality is not strict, the solution set is $\{x \mid x \leq -2 \text{ or } x \geq 2\}$ or, using interval notation, $(-\infty, -2] \cup [2, \infty)$.

f. $f(x) > g(x)$

$$x^2 - 4 > -x^2 + 4$$


$$2x^2 - 8 > 0$$

We graph the function $p(x) = 2x^2 - 8$.

y-intercept: $p(0) = -8$

$$\begin{aligned}x\text{-intercepts: } 2x^2 - 8 &= 0 \\2(x+2)(x-2) &= 0 \\x &= -2; x = 2\end{aligned}$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(2)} = 0$. Since $p(0) = -8$, the vertex is $(0, -8)$.



The graph is above the x -axis when $x < -2$ or $x > 2$. Since the inequality is strict, the solution set is $\{x \mid x < -2 \text{ or } x > 2\}$ or, using interval notation, $(-\infty, -2) \cup (2, \infty)$.

g. $f(x) \geq 1$

$$x^2 - 4 \geq 1$$


$$x^2 - 5 \geq 0$$

We graph the function $p(x) = x^2 - 5$.

y-intercept: $p(0) = -5$

$$\begin{aligned}x\text{-intercepts: } x^2 - 5 &= 0 \\x^2 &= 5 \\x &= \pm\sqrt{5}\end{aligned}$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(1)} = 0$. Since $p(0) = -5$, the vertex is $(0, -5)$.



The graph of p is above the x -axis when $x < -\sqrt{5}$ or $x > \sqrt{5}$. Since the inequality is not strict, the solution set is $\{x \mid x \leq -\sqrt{5} \text{ or } x \geq \sqrt{5}\}$ or, using interval notation, $(-\infty, -\sqrt{5}] \cup [\sqrt{5}, \infty)$.

Chapter 2: Linear and Quadratic Functions

30. $f(x) = x^2 - 2x + 1$; $g(x) = -x^2 + 1$

a. $f(x) = 0$

$$x^2 - 2x + 1 = 0$$

$$(x-1)^2 = 0$$

$$x - 1 = 0$$

$$x = 1$$

Solution set: $\{1\}$.

b. $g(x) = 0$

$$-x^2 + 1 = 0$$

$$x^2 - 1 = 0$$

$$(x+1)(x-1) = 0$$

$$x = -1; x = 1$$

Solution set: $\{-1, 1\}$.

c. $f(x) = g(x)$

$$x^2 - 2x + 1 = -x^2 + 1$$

$$2x^2 - 2x = 0$$

$$2x(x-1) = 0$$

$$x = 0, x = 1$$

Solution set: $\{0, 1\}$.

d. $f(x) > 0$

$$x^2 - 2x + 1 > 0$$

We graph the function $f(x) = x^2 - 2x + 1$.

y-intercept: $f(0) = 1$

x-intercepts: $x^2 - 2x + 1 = 0$


$$(x-1)^2 = 0$$

$$x - 1 = 0$$

$$x = 1$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-2)}{2(1)} = \frac{2}{2} = 1$.

Since $f(1) = 0$, the vertex is $(1, 0)$.



The graph is above the x -axis when $x < 1$ or $x > 1$. Since the inequality is strict, the solution set is $\{x \mid x < 1 \text{ or } x > 1\}$ or, using interval notation, $(-\infty, 1) \cup (1, \infty)$.

e. $g(x) \leq 0$

$$-x^2 + 1 \leq 0$$

We graph the function $g(x) = -x^2 + 1$.

y-intercept: $g(0) = 1$

x-intercepts: $-x^2 + 1 = 0$


$$x^2 - 1 = 0$$

$$(x+1)(x-1) = 0$$

$$x = -1; x = 1$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(0)}{2(-1)} = 0$. Since

$g(0) = 1$, the vertex is $(0, 1)$.



The graph is below the x -axis when $x < -1$ or $x > 1$. Since the inequality is not strict, the solution set is $\{x \mid x \leq -1 \text{ or } x \geq 1\}$ or, using interval notation, $(-\infty, -1] \cup [1, \infty)$.

f. $f(x) > g(x)$

$$x^2 - 2x + 1 > -x^2 + 1$$

$$2x^2 - 2x > 0$$

We graph the function $p(x) = 2x^2 - 2x$.

y-intercept: $p(0) = 0$


x-intercepts: $2x^2 - 2x = 0$

$$2x(x-1) = 0$$

$$x = 0; x = 1$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-2)}{2(2)} = \frac{2}{4} = \frac{1}{2}$.

Since $p\left(\frac{1}{2}\right) = \frac{1}{2}$, the vertex is $\left(\frac{1}{2}, \frac{1}{2}\right)$.



The graph is above the x -axis when $x < 0$ or $x > 1$. Since the inequality is strict, the

solution set is $\{x \mid x < 0 \text{ or } x > 1\}$ or, using interval notation, $(-\infty, 0) \cup (1, \infty)$.

g. $f(x) \geq 1$

$$x^2 - 2x + 1 \geq 1$$

$$x^2 - 2x \geq 0$$

We graph the function $p(x) = x^2 - 2x$.

y-intercept: $p(0) = 0$


x-intercepts: $x^2 - 2x = 0$

$$x(x - 2) = 0$$

$$x = 0; x = 2$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-2)}{2(1)} = \frac{2}{2} = 1$.

Since $p(1) = -1$, the vertex is $(1, -1)$.



The graph of p is above the x -axis when $x < 0$ or $x > 2$. Since the inequality is not strict, the solution set is $\{x \mid x \leq 0 \text{ or } x \geq 2\}$ or, using interval notation, $(-\infty, 0] \cup [2, \infty)$.

31. $f(x) = x^2 - x - 2$; $g(x) = x^2 + x - 2$

a. $f(x) = 0$

$$x^2 - x - 2 = 0$$

$$(x - 2)(x + 1) = 0$$

$$x = 2, x = -1$$

Solution set: $\{-1, 2\}$.

b. $g(x) = 0$

$$x^2 + x - 2 = 0$$

$$(x + 2)(x - 1) = 0$$

$$x = -2; x = 1$$

Solution set: $\{-2, 1\}$.

c. $f(x) = g(x)$

$$x^2 - x - 2 = x^2 + x - 2$$

$$-2x = 0$$

$$x = 0$$

Solution set: $\{0\}$.

d. $f(x) > 0$

$$x^2 - x - 2 > 0$$

We graph the function $f(x) = x^2 - x - 2$.

y-intercept: $f(0) = -2$


x-intercepts: $x^2 - x - 2 = 0$

$$(x - 2)(x + 1) = 0$$

$$x = 2; x = -1$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-1)}{2(1)} = \frac{1}{2}$. Since

$$f\left(\frac{1}{2}\right) = -\frac{9}{4}, \text{ the vertex is } \left(\frac{1}{2}, -\frac{9}{4}\right).$$



The graph is above the x -axis when $x < -1$ or $x > 2$. Since the inequality is strict, the solution set is $\{x \mid x < -1 \text{ or } x > 2\}$ or, using interval notation, $(-\infty, -1) \cup (2, \infty)$.

e. $g(x) \leq 0$

$$x^2 + x - 2 \leq 0$$

We graph the function $g(x) = x^2 + x - 2$.

y-intercept: $g(0) = -2$


x-intercepts: $x^2 + x - 2 = 0$

$$(x + 2)(x - 1) = 0$$

$$x = -2; x = 1$$

The vertex is at $x = \frac{-b}{2a} = \frac{-1}{2(1)} = -\frac{1}{2}$. Since

$$g\left(-\frac{1}{2}\right) = -\frac{7}{4}, \text{ the vertex is } \left(-\frac{1}{2}, -\frac{7}{4}\right).$$



The graph is below the x -axis when $-2 < x < 1$. Since the inequality is not strict, the solution set is $\{x \mid -2 \leq x \leq 1\}$ or, using interval notation, $[-2, 1]$.

Chapter 2: Linear and Quadratic Functions

f. $f(x) > g(x)$

$$x^2 - x - 2 > x^2 + x - 2$$

$$-2x > 0$$

$$x < 0$$

The solution set is $\{x \mid x < 0\}$ or, using interval notation, $(-\infty, 0)$.

g. $f(x) \geq 1$

$$x^2 - x - 2 \geq 1$$

$$x^2 - x - 3 \geq 0$$

We graph the function $p(x) = x^2 - x - 3$.

y-intercept: $p(0) = -3$

x-intercepts: $x^2 - x - 3 = 0$


$$x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(1)(-3)}}{2(1)}$$

$$= \frac{1 \pm \sqrt{1+12}}{2} = \frac{1 \pm \sqrt{13}}{2}$$

$$x \approx -1.30 \text{ or } x \approx 2.30$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-1)}{2(1)} = \frac{1}{2}$. Since

$$p\left(\frac{1}{2}\right) = -\frac{13}{4}, \text{ the vertex is } \left(\frac{1}{2}, -\frac{13}{4}\right).$$



The graph of p is above the x -axis when

$$x < \frac{1-\sqrt{13}}{2} \text{ or } x > \frac{1+\sqrt{13}}{2}.$$

Since the inequality is not strict, the solution set is

$$\left\{x \mid x \leq \frac{1-\sqrt{13}}{2} \text{ or } x \geq \frac{1+\sqrt{13}}{2}\right\} \text{ or, using}$$

interval notation,

$$\left(-\infty, \frac{1-\sqrt{13}}{2}\right] \cup \left[\frac{1+\sqrt{13}}{2}, \infty\right).$$

32. $f(x) = -x^2 - x + 1$; $g(x) = -x^2 + x + 6$

a. $f(x) = 0$

$$-x^2 - x + 1 = 0$$

$$x^2 + x - 1 = 0$$

$$x = \frac{-(1) \pm \sqrt{(1)^2 - 4(1)(-1)}}{2(1)}$$

$$= \frac{-1 \pm \sqrt{1+4}}{2} = \frac{-1 \pm \sqrt{5}}{2}$$

$$\text{Solution set: } \left\{\frac{-1-\sqrt{5}}{2}, \frac{-1+\sqrt{5}}{2}\right\}.$$

b. $g(x) = 0$

$$-x^2 + x + 6 = 0$$

$$x^2 - x - 6 = 0$$

$$(x-3)(x+2) = 0$$

$$x = 3; x = -2$$

$$\text{Solution set: } \{-2, 3\}.$$

c. $f(x) = g(x)$

$$-x^2 - x + 1 = -x^2 + x + 6$$

$$-2x - 5 = 0$$

$$-2x = 5$$

$$x = -\frac{5}{2}$$

$$\text{Solution set: } \left\{-\frac{5}{2}\right\}.$$

d. $f(x) > 0$

$$-x^2 - x + 1 > 0$$

We graph the function $f(x) = -x^2 - x + 1$.

y-intercept: $f(0) = -1$

x-intercepts: $-x^2 - x + 2 = 0$

$$x^2 + x - 2 = 0$$

$$x = \frac{-(1) \pm \sqrt{(1)^2 - 4(1)(-2)}}{2(1)}$$


$$= \frac{-1 \pm \sqrt{1+8}}{2} = \frac{-1 \pm \sqrt{9}}{2}$$

$$x \approx -1.62 \text{ or } x \approx 0.62$$

$$\text{The vertex is at } x = \frac{-b}{2a} = \frac{-(-1)}{2(-1)} = \frac{1}{-2} = -\frac{1}{2}.$$

$$\text{Since } f\left(-\frac{1}{2}\right) = \frac{5}{4}, \text{ the vertex is } \left(-\frac{1}{2}, \frac{5}{4}\right).$$

Section 2.5: Inequalities Involving Quadratic Functions



The graph is above the x -axis when

$$\frac{-1-\sqrt{5}}{2} < x < \frac{-1+\sqrt{5}}{2}.$$

Since the inequality

$$\left\{ x \mid \frac{-1-\sqrt{5}}{2} < x < \frac{-1+\sqrt{5}}{2} \right\}$$

$$\text{or, using interval notation, } \left(\frac{-1-\sqrt{5}}{2}, \frac{-1+\sqrt{5}}{2} \right).$$

e. $g(x) \leq 0$

$$-x^2 + x + 6 \leq 0$$

We graph the function $g(x) = -x^2 + x + 6$.

y-intercept: $g(0) = 6$

x-intercepts: $-x^2 + x + 6 = 0$


$$x^2 - x - 6 = 0$$

$$(x-3)(x+2) = 0$$

$$x = 3; x = -2$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-1)}{2(-1)} = \frac{-1}{-2} = \frac{1}{2}$.

Since $f\left(\frac{1}{2}\right) = \frac{25}{4}$, the vertex is $\left(\frac{1}{2}, \frac{25}{4}\right)$.



The graph is below the x -axis when $x < -2$

or $x > 3$. Since the inequality is not strict,

the solution set is $\{x \mid x \leq -2 \text{ or } x \geq 3\}$ or,

using interval notation, $(-\infty, -2] \cup [3, \infty)$.

f. $f(x) > g(x)$

$$-x^2 - x + 1 > -x^2 + x + 6$$

$$-2x > 5$$

$$x < -\frac{5}{2}$$

The solution set is $\{x \mid x < -\frac{5}{2}\}$ or, using

interval notation, $(-\infty, -\frac{5}{2})$.

g. $f(x) \geq 1$

$$-x^2 - x + 1 \geq 1$$

$$-x^2 - x \geq 0$$

We graph the function $p(x) = -x^2 - x$.

y-intercept: $p(0) = 0$


x-intercepts: $-x^2 - x = 0$

$$-x(x+1) = 0$$

$$x = 0; x = -1$$

The vertex is at $x = \frac{-b}{2a} = \frac{-(-1)}{2(-1)} = \frac{1}{-2} = -\frac{1}{2}$.

Since $p\left(-\frac{1}{2}\right) = \frac{1}{4}$, the vertex is $\left(-\frac{1}{2}, \frac{1}{4}\right)$.



The graph of p is above the x -axis when

$-1 < x < 0$. Since the inequality is not

strict, the solution set is $\{x \mid -1 \leq x < 0\}$ or,

using interval notation, $[-1, 0)$.

33. a. The ball strikes the ground when

$$s(t) = 80t - 16t^2 = 0.$$

$$80t - 16t^2 = 0$$

$$16t(5-t) = 0$$

$$t = 0, t = 5$$

The ball strikes the ground after 5 seconds.

b. Find the values of t for which

$$80t - 16t^2 > 96$$

$$-16t^2 + 80t - 96 > 0$$

We graph the function

$$f(t) = -16t^2 + 80t - 96.$$

The intercepts are

y-intercept: $f(0) = -96$

t-intercepts: $-16t^2 + 80t - 96 = 0$

$$-16(t^2 - 5t + 6) = 0$$


$$16(t-2)(t-3) = 0$$

$$t = 2, t = 3$$

The vertex is at $t = \frac{-b}{2a} = \frac{-(80)}{2(-16)} = 2.5$.

Since $f(2.5) = 4$, the vertex is $(2.5, 4)$.

Chapter 2: Linear and Quadratic Functions



The graph of f is above the t -axis when $2 < t < 3$. Since the inequality is strict, the solution set is $\{t \mid 2 < t < 3\}$ or, using interval notation, $(2, 3)$. The ball is more than 96 feet above the ground for times between 2 and 3 seconds.

34. a. The ball strikes the ground when $s(t) = 96t - 16t^2 = 0$.

$$96t - 16t^2 = 0$$

$$16t(6 - t) = 0$$

$$t = 0, t = 6$$

The ball strikes the ground after 6 seconds.

- b. Find the values of t for which

$$96t - 16t^2 > 128$$

$$-16t^2 + 96t - 128 > 0$$

We graph $f(t) = -16t^2 + 96t - 128$. The intercepts are

$$\text{y-intercept: } f(0) = -128$$

$$\text{t-intercepts: } -16t^2 + 96t - 128 = 0$$


$$16(t^2 - 6t + 8) = 0$$

$$-16(t - 4)(t - 2) = 0$$

$$t = 4, t = 2$$

The vertex is at $t = \frac{-b}{2a} = \frac{-(96)}{2(-16)} = 3$. Since

$$f(3) = 16, \text{ the vertex is } (3, 16).$$



The graph of f is above the t -axis when $2 < t < 4$. Since the inequality is strict, the solution set is $\{t \mid 2 < t < 4\}$ or, using interval notation, $(2, 4)$. The ball is more than 128 feet above the ground for times between 2 and 4 seconds.

35. a. $R(p) = -4p^2 + 4000p = 0$

$$-4p(p - 1000) = 0$$

$$p = 0, p = 1000$$

Thus, the revenue equals zero when the price is \$0 or \$1000.

- b. Find the values of p for which

$$-4p^2 + 4000p > 800,000$$

$$-4p^2 + 4000p - 800,000 > 0$$

$$\text{We graph } f(p) = -4p^2 + 4000p - 800,000.$$

The intercepts are

$$\text{y-intercept: } f(0) = -800,000$$

p -intercepts:

$$-4p^2 + 4000p - 800,000 = 0$$

$$p^2 - 1000p + 200,000 = 0$$

$$p = \frac{-(-1000) \pm \sqrt{(-1000)^2 - 4(1)(200,000)}}{2(1)}$$

$$= \frac{1000 \pm \sqrt{200,000}}{2}$$


$$= \frac{1000 \pm 200\sqrt{5}}{2}$$

$$= 500 \pm 100\sqrt{5}$$

$$p \approx 276.39; p \approx 723.61.$$

$$\text{The vertex is at } p = \frac{-b}{2a} = \frac{-(4000)}{2(-4)} = 500.$$

Since $f(500) = 200,000$, the vertex is $(500, 200,000)$.



The graph of f is above the p -axis when $276.39 < p < 723.61$. Since the inequality is strict, the solution set is $\{p \mid 276.39 < p < 723.61\}$ or, using interval notation, $(276.39, 723.61)$. The revenue is more than \$800,000 for prices between \$276.39 and \$723.61.

36. a. $R(p) = -\frac{1}{2}p^2 + 1900p = 0$

$$-\frac{1}{2}p(p - 3800) = 0$$

$$p = 0, p = 3800$$

Thus, the revenue equals zero when the price is \$0 or \$3800.

b. Find the values of p for which

$$-\frac{1}{2}p^2 + 1900p > 1200000$$

$$-\frac{1}{2}p^2 + 1900p - 1200000 > 0$$

We graph $f(p) = -\frac{1}{2}p^2 + 1900p - 1200000$.

The intercepts are

y-intercept: $f(0) = -1,200,000$

$$p\text{-intercepts: } -\frac{1}{2}p^2 + 1900p - 1200000 = 0$$


$$p^2 - 3800p + 2400000 = 0$$

$$(p - 800)(p - 3000) = 0$$

$$p = 800; p = 3000$$

The vertex is at $p = \frac{-b}{2a} = \frac{-(-1900)}{2(1/2)} = 1900$.

Since $f(1900) = 605,000$, the vertex is $(1900, 605000)$.



The graph of f is above the p -axis when $800 < p < 3000$. Since the inequality is strict, the solution set is $\{p \mid 800 < p < 3000\}$ or, using interval notation, $(800, 3000)$. The revenue is more than \$1,200,000 for prices between \$800 and \$3000.

37. $(x - 4)^2 \leq 0$

We graph the function $f(x) = (x - 4)^2$.


y-intercept: $f(0) = 16$

$$x\text{-intercepts: } (x - 4)^2 = 0$$

$$x - 4 = 0$$

$$x = 4$$

The vertex is the vertex is $(4, 0)$.



The graph is never below the x -axis. Since the inequality is not strict, the only solution comes from the x -intercept. Therefore, the given inequality has exactly one real solution, namely $x = 4$.

38. $(x - 2)^2 > 0$

We graph the function $f(x) = (x - 2)^2$.


y-intercept: $f(0) = 4$

$$x\text{-intercepts: } (x - 2)^2 = 0$$

$$x - 2 = 0$$

$$x = 2$$

The vertex is the vertex is $(2, 0)$.



The graph is above the x -axis when $x < 2$ or $x > 2$. Since the inequality is strict, the solution set is $\{x \mid x < 2 \text{ or } x > 2\}$. Therefore, the given inequality has exactly one real number that is not a solution, namely $x \neq 2$.

39. Solving $x^2 + x + 1 > 0$

We graph the function $f(x) = x^2 + x + 1$.


y-intercept: $f(0) = 1$

x -intercepts: $b^2 - 4ac = 1^2 - 4(1)(1) = -3$, so f has no x -intercepts.

The vertex is at $x = \frac{-b}{2a} = \frac{-(1)}{2(1)} = -\frac{1}{2}$. Since

$$f\left(-\frac{1}{2}\right) = \frac{3}{4}, \text{ the vertex is } \left(-\frac{1}{2}, \frac{3}{4}\right).$$

Chapter 2: Linear and Quadratic Functions



The graph is always above the x -axis. Thus, the solution is the set of all real numbers or, using interval notation, $(-\infty, \infty)$.

40. Solving $x^2 - x + 1 < 0$


We graph the function $f(x) = x^2 - x + 1$.

y-intercept: $f(0) = 1$

x-intercepts: $b^2 - 4ac = (-1)^2 - 4(1)(1) = -3$, so f has no x -intercepts.

The vertex is at $x = \frac{-b}{2a} = \frac{-(-1)}{2(1)} = \frac{1}{2}$. Since

$f\left(-\frac{1}{2}\right) = \frac{3}{4}$, the vertex is $\left(-\frac{1}{2}, \frac{3}{4}\right)$.



The graph is never below the x -axis. Thus, the inequality has no solution. That is, the solution set is $\{ \}$ or $\emptyset$.

41. If the inequality is not strict, we include the x -intercepts in the solution.

Section 2.6

1. $R = 3x$

2. Use LIN REGression to get
 $y = 1.7826x + 4.0652$

3. a. $R(x) = x\left(-\frac{1}{6}x + 100\right) = -\frac{1}{6}x^2 + 100x$

b. The quantity sold price cannot be negative, so $x \geq 0$. Similarly, the price should be positive, so $p > 0$.

$$-\frac{1}{6}x + 100 > 0$$

$$-\frac{1}{6}x > -100$$

$$x < 600$$

Thus, the implied domain for R is $\{x \mid 0 \leq x < 600\}$ or $[0, 600)$.

$$\begin{aligned} \text{c. } R(200) &= -\frac{1}{6}(200)^2 + 100(200) \\ &= \frac{-20000}{3} + 20000 \\ &= \frac{40000}{3} \approx \$13,333.33 \end{aligned}$$

$$\text{d. } x = \frac{-b}{2a} = \frac{-100}{2\left(-\frac{1}{6}\right)} = \frac{-100}{\left(-\frac{1}{3}\right)} = \frac{300}{1} = 300$$

The maximum revenue is

$$\begin{aligned} R(300) &= -\frac{1}{6}(300)^2 + 100(300) \\ &= -15000 + 30000 \\ &= \$15,000 \end{aligned}$$

$$\text{e. } p = -\frac{1}{6}(300) + 100 = -50 + 100 = \$50$$

$$\text{4. a. } R(x) = x\left(-\frac{1}{3}x + 100\right) = -\frac{1}{3}x^2 + 100x$$

b. The quantity sold price cannot be negative, so $x \geq 0$. Similarly, the price should be positive, so $p > 0$.

$$-\frac{1}{3}x + 100 > 0$$

$$-\frac{1}{3}x > -100$$

$$x < 300$$

Thus, the implied domain for R is $\{x \mid 0 \leq x < 300\}$ or $[0, 300)$.

$$\begin{aligned} \text{c. } R(100) &= -\frac{1}{3}(100)^2 + 100(100) \\ &= \frac{-10000}{3} + 10000 \\ &= \frac{20000}{3} \approx \$6,666.67 \end{aligned}$$

Section 2.6: Building Quadratic Models from Verbal Descriptions and from Data

d. $x = \frac{-b}{2a} = \frac{-100}{2(-\frac{1}{3})} = \frac{-100}{(-\frac{2}{3})} = \frac{300}{2} = 150$

The maximum revenue is

$$R(150) = -\frac{1}{3}(150)^2 + 100(150) \\ = -7500 + 15000 = \$7,500$$

e. $p = -\frac{1}{3}(150) + 100 = -50 + 100 = \50

5. a. If $x = -5p + 100$, then $p = \frac{100-x}{5}$.

$$R(x) = x\left(\frac{100-x}{5}\right) = -\frac{1}{5}x^2 + 20x$$

b. $R(15) = -\frac{1}{5}(15)^2 + 20(15) \\ = -45 + 300 = \255

c. $x = \frac{-b}{2a} = \frac{-20}{2(-\frac{1}{5})} = \frac{-20}{(-\frac{2}{5})} = \frac{100}{2} = 50$

The maximum revenue is


$$R(50) = -\frac{1}{5}(50)^2 + 20(50) \\ = -500 + 1000 = \$500$$

d. $p = \frac{100-50}{5} = \frac{50}{5} = \10

e. Graph $R = -\frac{1}{5}x^2 + 20x$ and $R = 480$. Find

where the graphs intersect by solving

$$480 = -\frac{1}{5}x^2 + 20x.$$



$$\frac{1}{5}x^2 - 20x + 480 = 0$$

$$x^2 - 100x + 2400 = 0$$

$$(x-40)(x-60) = 0$$

$$x = 40, x = 60$$

Solve for price.

$$x = -5p + 100$$

$$40 = -5p + 100 \Rightarrow p = \$12$$

$$60 = -5p + 100 \Rightarrow p = \$8$$

The company should charge between \$8 and \$12.

6. a. If $x = -20p + 500$, then $p = \frac{500-x}{20}$.

$$R(x) = x\left(\frac{500-x}{20}\right) = -\frac{1}{20}x^2 + 25x$$

b. $R(20) = -\frac{1}{20}(20)^2 + 25(20) \\ = -20 + 500 = \480

c. $x = \frac{-b}{2a} = \frac{-25}{2(-\frac{1}{20})} = \frac{-25}{(-\frac{1}{10})} = \frac{250}{1} = 250$.

The maximum revenue is


$$R(250) = -\frac{1}{20}(250)^2 + 25(250) \\ = -3125 + 6250 = \$3125$$

d. $p = \frac{500-250}{20} = \frac{250}{20} = \12.50

e. Graph $R = -\frac{1}{20}x^2 + 25x$ and $R = 3000$.

Find where the graphs intersect by solving

$$3000 = -\frac{1}{20}x^2 + 25x.$$



$$\frac{1}{20}x^2 - 25x + 3000 = 0$$

$$x^2 - 500x + 60000 = 0$$

$$(x-200)(x-300) = 0$$

$$x = 200, x = 300$$

Solve for price.

$$x = -20p + 500$$

$$300 = -20p + 500 \Rightarrow p = \$10$$

$$200 = -20p + 500 \Rightarrow p = \$15$$

The company should charge between \$10 and \$15.

Chapter 2: Linear and Quadratic Functions

7. a. Let w = width and l = length of the rectangular area.
Solving $P = 2w + 2l = 400$ for l :
$$l = \frac{400 - 2w}{2} = 200 - w.$$

Then $A(w) = (200 - w)w = 200w - w^2$
$$= -w^2 + 200w$$
- b. $w = \frac{-b}{2a} = \frac{-200}{2(-1)} = \frac{-200}{-2} = 100$ yards
- c. $A(100) = -100^2 + 200(100)$
$$= -10000 + 20000$$

$$= 10,000 \text{ yd}^2$$
8. a. Let x = width and y = width of the rectangle.
Solving $P = 2x + 2y = 3000$ for y :
$$y = \frac{3000 - 2x}{2} = 1500 - x.$$

Then $A(x) = (1500 - x)x$
$$= 1500x - x^2$$

$$= -x^2 + 1500x.$$
- b. $x = \frac{-b}{2a} = \frac{-1500}{2(-1)} = \frac{-1500}{-2} = 750$ feet
- c. $A(750) = -750^2 + 1500(750)$
$$= -562500 + 1125000$$

$$= 562,500 \text{ ft}^2$$
9. Let x = width and y = length of the rectangle.
Solving $P = 2x + y = 4000$ for y :
$$y = 4000 - 2x.$$

Then $A(x) = (4000 - 2x)x$
$$= 4000x - 2x^2$$

$$= -2x^2 + 4000x$$

$$x = \frac{-b}{2a} = \frac{-4000}{2(-2)} = \frac{-4000}{-4} = 1000 \text{ meters}$$

maximizes area.
$$A(1000) = -2(1000)^2 + 4000(1000).$$

$$= -2000000 + 4000000$$

$$= 2,000,000$$

The largest area that can be enclosed is 2,000,000 square meters.

10. Let x = width and y = length of the rectangle.
$$2x + y = 2000$$

$$y = 2000 - 2x$$

Then $A(x) = (2000 - 2x)x$
$$= 2000x - 2x^2$$

$$= -2x^2 + 2000x$$

$$x = \frac{-b}{2a} = \frac{-2000}{2(-2)} = \frac{-2000}{-4} = 500 \text{ meters}$$

maximizes area.
$$A(500) = -2(500)^2 + 2000(500)$$

$$= -500,000 + 1,000,000$$

$$= 500,000$$

The largest area that can be enclosed is 500,000 square meters.
11. $h(x) = \frac{-32x^2}{(50)^2} + x + 200 = -\frac{8}{625}x^2 + x + 200$
- a. $a = -\frac{8}{625}, b = 1, c = 200.$
The maximum height occurs when
$$x = \frac{-b}{2a} = \frac{-1}{2(-8/625)} = \frac{625}{16} \approx 39 \text{ feet from}$$

base of the cliff.
- b. The maximum height is
$$h\left(\frac{625}{16}\right) = \frac{-8}{625}\left(\frac{625}{16}\right)^2 + \frac{625}{16} + 200$$

$$= \frac{7025}{32} \approx 219.5 \text{ feet.}$$
- c. Solving when $h(x) = 0$:
$$-\frac{8}{625}x^2 + x + 200 = 0$$


$$x = \frac{-1 \pm \sqrt{1^2 - 4(-8/625)(200)}}{2(-8/625)}$$

$$x \approx \frac{-1 \pm \sqrt{11.24}}{-0.0256}$$


$$x \approx -91.90 \text{ or } x \approx 170$$

Since the distance cannot be negative, the projectile strikes the water approximately 170 feet from the base of the cliff.


Section 2.6: Building Quadratic Models from Verbal Descriptions and from Data



e. Using the MAXIMUM function



Using the ZERO function



f.

$$-\frac{8}{625}x^2 + x + 200 = 100$$

$$-\frac{8}{625}x^2 + x + 100 = 0$$

$$x = \frac{\sqrt{1^2 - 4(-8/625)(100)}}{2(-8/625)} = \frac{-1 \pm \sqrt{6.12}}{-0.0256}$$

$$x \approx -57.57 \text{ or } x \approx 135.70$$

Since the distance cannot be negative, the projectile is 100 feet above the water when it is approximately 135.7 feet from the base of the cliff.

12. a. $h(x) = \frac{-32x^2}{(100)^2} + x = -\frac{2}{625}x^2 + x$

$$a = -\frac{2}{625}, b = 1, c = 0.$$

The maximum height occurs when

$$x = \frac{-b}{2a} = \frac{-1}{2(-2/625)} = \frac{625}{4} = 156.25 \text{ feet}$$

b. The maximum height is

$$h\left(\frac{625}{4}\right) = \frac{-2}{625}\left(\frac{625}{4}\right)^2 + \frac{625}{4}$$

$$= \frac{625}{8} = 78.125 \text{ feet}$$

c. Solving when $h(x) = 0$:

$$-\frac{2}{625}x^2 + x = 0$$


$$x\left(-\frac{2}{625}x + 1\right) = 0$$

$$x = 0 \text{ or } -\frac{2}{625}x + 1 = 0$$


$$x = 0 \text{ or } 1 = \frac{2}{625}x$$

$$x = 0 \text{ or } x = \frac{625}{2} = 312.5$$


Since the distance cannot be zero, the projectile lands 312.5 feet from where it was fired.



e. Using the MAXIMUM function



Using the ZERO function



f. Solving when $h(x) = 50$:

$$-\frac{2}{625}x^2 + x = 50$$

$$-\frac{2}{625}x^2 + x - 50 = 0$$

$$x = \frac{-1 \pm \sqrt{1^2 - 4(-2/625)(-50)}}{2(-2/625)}$$

$$= \frac{-1 \pm \sqrt{0.36}}{-0.0064} \approx \frac{-1 \pm 0.6}{-0.0064}$$

$$x = 62.5 \text{ or } x = 250$$

The projectile is 50 feet above the ground 62.5 feet and 250 feet from where it was fired.


Chapter 2: Linear and Quadratic Functions

13. Locate the origin at the point where the cable touches the road. Then the equation of the parabola is of the form: $y = ax^2$, where $a > 0$. Since the point $(200, 75)$ is on the parabola, we can find the constant a :

$$\text{Since } 75 = a(200)^2, \text{ then } a = \frac{75}{200^2} = 0.001875.$$

When $x = 100$, we have:


$$y = 0.001875(100)^2 = 18.75 \text{ meters.}$$



14. Locate the origin at the point directly under the highest point of the arch. Then the equation of the parabola is of the form: $y = -ax^2 + k$, where $a > 0$. Since the maximum height is 25 feet, when $x = 0$, $y = k = 25$. Since the point $(60, 0)$ is on the parabola, we can find the constant a : Since $0 = -a(60)^2 + 25$ then

$$a = \frac{25}{60^2}. \text{ The equation of the parabola is:}$$

$$h(x) = -\frac{25}{60^2}x^2 + 25.$$



At $x = 10$:

$$h(10) = -\frac{25}{60^2}(10)^2 + 25 = -\frac{25}{36} + 25 \approx 24.3 \text{ ft.}$$

At $x = 20$:

$$h(20) = -\frac{25}{60^2}(20)^2 + 25 = -\frac{25}{9} + 25 \approx 22.2 \text{ ft.}$$

At $x = 40$:

$$h(40) = -\frac{25}{60^2}(40)^2 + 25 = -\frac{100}{9} + 25 \approx 13.9 \text{ ft.}$$

15. a. Let x = the depth of the gutter and y the width of the gutter. Then $A = xy$ is the cross-sectional area of the gutter. Since the aluminum sheets for the gutter are 12 inches wide, we have $2x + y = 12$. Solving for y : $y = 12 - 2x$. The area is to be maximized, so:


$A = xy = x(12 - 2x) = -2x^2 + 12x$. This equation is a parabola opening down; thus, it has a maximum

$$\text{when } x = \frac{-b}{2a} = \frac{-12}{2(-2)} = \frac{-12}{-4} = 3.$$

Thus, a depth of 3 inches produces a maximum cross-sectional area.

- b. Graph $A = -2x^2 + 12x$ and $A = 16$. Find where the graphs intersect by solving

$$16 = -2x^2 + 12x.$$



$$2x^2 - 12x + 16 = 0$$

$$x^2 - 6x + 8 = 0$$

$$(x-4)(x-2) = 0$$

$$x = 4, x = 2$$

The graph of $A = -2x^2 + 12x$ is above the graph of $A = 16$ where the depth is between 2 and 4 inches.

16. Let x = width of the window and y = height of the rectangular part of the window. The perimeter of the window is: $x + 2y + \frac{\pi x}{2} = 20$.

$$\text{Solving for } y: y = \frac{40 - 2x - \pi x}{4}.$$

The area of the window is:

$$\begin{aligned} A(x) &= x \left(\frac{40 - 2x - \pi x}{4} \right) + \frac{1}{2} \pi \left(\frac{x}{2} \right)^2 \\ &= 10x - \frac{x^2}{2} - \frac{\pi x^2}{4} + \frac{\pi x^2}{8} \\ &= \left(-\frac{1}{2} - \frac{\pi}{8} \right) x^2 + 10x. \end{aligned}$$

This equation is a parabola opening down; thus, it has a maximum when

Section 2.6: Building Quadratic Models from Verbal Descriptions and from Data

$$x = \frac{-b}{2a} = \frac{-10}{2\left(-\frac{1}{2} - \frac{\pi}{8}\right)} = \frac{10}{\left(1 + \frac{\pi}{4}\right)} \approx 5.6 \text{ feet}$$

$$y = \frac{40 - 2(5.60) - \pi(5.60)}{4} \approx 2.8 \text{ feet}$$

The width of the window is about 5.6 feet and the height of the rectangular part is approximately 2.8 feet. The radius of the semicircle is roughly 2.8 feet, so the total height is about 5.6 feet.

17. Let x = the width of the rectangle or the diameter of the semicircle and let y = the length of the rectangle. The perimeter of each semicircle is $\frac{\pi x}{2}$.

The perimeter of the track is given

$$\text{by: } \frac{\pi x}{2} + \frac{\pi x}{2} + y + y = 1500.$$

Solving for x :

$$\pi x + 2y = 1500$$

$$\pi x = 1500 - 2y$$

$$x = \frac{1500 - 2y}{\pi}$$

The area of the rectangle is:

$$A = xy = \left(\frac{1500 - 2y}{\pi}\right)y = \frac{-2}{\pi}y^2 + \frac{1500}{\pi}y.$$

This equation is a parabola opening down; thus, it has a maximum when

$$y = \frac{-b}{2a} = \frac{-1500}{2\left(\frac{-2}{\pi}\right)} = \frac{-1500}{-4} = 375.$$

$$\text{Thus, } x = \frac{1500 - 2(375)}{\pi} = \frac{750}{\pi} \approx 238.73$$

The dimensions for the rectangle with maximum

$$\text{area are } \frac{750}{\pi} \approx 238.73 \text{ meters by } 375 \text{ meters.}$$

18. Let x = width of the window and y = height of the rectangular part of the window. The perimeter of the window is:

$$3x + 2y = 16$$

$$y = \frac{16 - 3x}{2}$$

The area of the window is

$$\begin{aligned} A(x) &= x\left(\frac{16 - 3x}{2}\right) + \frac{\sqrt{3}}{4}x^2 \\ &= 8x - \frac{3}{2}x^2 + \frac{\sqrt{3}}{4}x^2 \\ &= \left(-\frac{3}{2} + \frac{\sqrt{3}}{4}\right)x^2 + 8x \end{aligned}$$

This equation is a parabola opening down; thus, it has a maximum when

$$\begin{aligned} x &= \frac{-b}{2a} = \frac{-8}{2\left(-\frac{3}{2} + \frac{\sqrt{3}}{4}\right)} \\ &= \frac{-8}{-3 + \frac{\sqrt{3}}{2}} = \frac{-16}{-6 + \sqrt{3}} \approx 3.75 \text{ ft.} \end{aligned}$$

The window is approximately 3.75 feet wide.

$$y = \frac{16 - 3\left(\frac{-16}{-6 + \sqrt{3}}\right)}{2} = \frac{16 + \frac{48}{-6 + \sqrt{3}}}{2} = 8 + \frac{24}{-6 + \sqrt{3}}$$

The height of the equilateral triangle is

$$\begin{aligned} \frac{\sqrt{3}}{2}\left(\frac{-16}{-6 + \sqrt{3}}\right) &= \frac{-8\sqrt{3}}{-6 + \sqrt{3}} \text{ feet, so the total height is} \\ 8 + \frac{24}{-6 + \sqrt{3}} + \frac{-8\sqrt{3}}{-6 + \sqrt{3}} &\approx 5.62 \text{ feet.} \end{aligned}$$

19. We are given: $V(x) = kx(a - x) = -kx^2 + akx$.

The reaction rate is a maximum when:

$$x = \frac{-b}{2a} = \frac{-ak}{2(-k)} = \frac{ak}{2k} = \frac{a}{2}.$$

20. We have:

$$a(-h)^2 + b(-h) + c = ah^2 - bh + c = y_0$$

$$a(0)^2 + b(0) + c = c = y_1$$

$$a(h)^2 + b(h) + c = ah^2 + bh + c = y_2$$

Equating the two equations for the area, we have:

$$\begin{aligned} y_0 + 4y_1 + y_2 &= ah^2 - bh + c + 4c + ah^2 + bh + c \\ &= 2ah^2 + 6c. \end{aligned}$$

Therefore,

$$\text{Area} = \frac{h}{3}(2ah^2 + 6c) = \frac{h}{3}(y_0 + 4y_1 + y_2) \text{ sq. units.}$$

21. $f(x) = -5x^2 + 8$, $h = 1$

$$\begin{aligned} \text{Area} &= \frac{h}{3}(2ah^2 + 6c) = \frac{1}{3}(2(-5)(1)^2 + 6(8)) \\ &= \frac{1}{3}(-10 + 48) = \frac{38}{3} \text{ sq. units} \end{aligned}$$

Chapter 2: Linear and Quadratic Functions

22. $f(x) = 2x^2 + 8$, $h = 2$

$$\begin{aligned} \text{Area} &= \frac{h}{3}(2ah^2 + 6c) = \frac{2}{3}(2(2)(2)^2 + 6(8)) \\ &= \frac{2}{3}(16 + 48) = \frac{2}{3}(64) = \frac{128}{3} \text{ sq. units} \end{aligned}$$


23. $f(x) = x^2 + 3x + 5$, $h = 4$

$$\begin{aligned} \text{Area} &= \frac{h}{3}(2ah^2 + 6c) = \frac{4}{3}(2(1)(4)^2 + 6(5)) \\ &= \frac{4}{3}(32 + 30) = \frac{248}{3} \text{ sq. units} \end{aligned}$$

24. $f(x) = -x^2 + x + 4$, $h = 1$

$$\begin{aligned} \text{Area} &= \frac{h}{3}(2ah^2 + 6c) = \frac{1}{3}(2(-1)(1)^2 + 6(4)) \\ &= \frac{1}{3}(-2 + 24) = \frac{1}{3}(22) = \frac{22}{3} \text{ sq. units} \end{aligned}$$

25. a.



From the graph, the data appear to follow a quadratic relation with $a < 0$.

b. Using the QUADratic REGression program

```
QuadReg
y=ax^2+bx+c
a=-45.46571429
b=4314.374286
c=-55961.675
```


$$I(x) = -45.466x^2 + 4314.374x - 55,961.675$$

c. $x = \frac{-b}{2a} = \frac{-4314.374}{2(-45.466)} \approx 47.4$


An individual will earn the most income at about 47.4 years of age.

d. The maximum income will be: $I(47.4) = -45.466(47.4)^2 + 4314.374(47.4) - 55,961.675 \approx \$46,388$

e.



26. a.



From the graph, the data appear to follow a quadratic relation with $a < 0$.

b. Using the QUADratic REGression program

```
QuadReg
y=ax^2+bx+c
a=-.0037121212
b=1.031818182
c=5.666666667
```


$$h(x) = -0.0037x^2 + 1.0318x + 5.6667$$

c. $x = \frac{-b}{2a} = \frac{-1.0318}{2(-0.0037)} \approx 139.4$


The ball will travel about 139.4 feet before it reaches its maximum height.

d. The maximum height will be: $h(139.4) = -0.0037(139.4)^2 + 1.0318(139.4) + 5.6667 \approx 77.6$ feet

e.



27. a.



From the graph, the data appear to be linearly related with $m > 0$.

Section 2.7: Complex Zeros of a Quadratic Function


- b. Using the LINear REGression program

```
LinReg
y=ax+b
a=.8358556937
b=1032.272871
r2=.8772192055
r=.9365998107
```

$$R(x) = 0.836x + 1032.273$$

- c. $R(850) = 0.836(850) + 1032.273 \approx 1743$
The rent for an 850 square-foot apartment in San Diego will be about \$1743 per month.

28. a.



From the graph, the data appear to follow a quadratic relation with $a < 0$.

- b. Using the QUADratic REGression program


```
QuadReg
y=ax2+bx+c
a=-.0174674623
b=1.934623878
c=-25.34083541
```

$$M(s) = -0.017s^2 + 1.935s - 25.341$$

- c. $M(63) = -0.017(63)^2 + 1.935(63) - 25.341 \approx 29.1$

A Camry traveling 63 miles per hour will get about 29.1 miles per gallon.

29. a.



From the graph, the data appear to follow a quadratic relation with $a > 0$.

- b. Using the QUADratic REGression program


```
QuadReg
y=ax2+bx+c
a=.0152142857
b=-1.332428571
c=39.82285714
```

$$P(a) = 0.015a^2 - 1.332a + 39.823$$

- c. $P(35) = 0.015(35)^2 - 1.332(35) + 39.823 \approx 11.6$

About 11.6% of 35-year-olds do not have a high school diploma.

30. a.



From the graph, the data appear to be linearly related with $m > 0$.

- b. Using the LINear REGression program

```
LinReg
y=ax+b
a=.2330507161
b=-2.037230647
r2=.7610474345
r=.8723803267
```

$$C(x) = 0.233x - 2.037$$

- c. $C(80) = 0.233(80) - 2.037 \approx 16.6$

When the temperature is 80°F, there will be about 16.6 chirps per second.

31. Answers will vary. One possibility follows: If the price is \$140, no one will buy the calculators, thus making the revenue \$0.

Section 2.7

1. Integers: $\{-3, 0\}$

Rationals: $\left\{-3, 0, \frac{6}{5}\right\}$

2. True; the set of real numbers consists of all rational and irrational numbers.

3. $10 - 5i$

4. $2 - 5i$

Chapter 2: Linear and Quadratic Functions

5. True

6. $9i$

7. $2 + 3i$

8. True


9. $f(x) = 0$

$$x^2 + 4 = 0$$

$$x^2 = -4$$

$$x = \pm\sqrt{-4} = \pm 2i$$

The zeros are $-2i$ and $2i$.




10. $f(x) = 0$

$$x^2 - 9 = 0$$

$$x^2 = 9$$

$$x = \pm\sqrt{9} = \pm 3$$

The zeros are -3 and 3 .




11. $f(x) = 0$

$$x^2 - 16 = 0$$

$$x^2 = 16$$

$$x = \pm\sqrt{16} = \pm 4$$

The zeros are -4 and 4 .




12. $f(x) = 0$

$$x^2 + 25 = 0$$

$$x^2 = -25$$

$$x = \pm\sqrt{-25} = \pm 5i$$

The zeros are $-5i$ and $5i$.



13. $f(x) = 0$


$$x^2 - 6x + 13 = 0$$

$$a = 1, b = -6, c = 13,$$

$$b^2 - 4ac = (-6)^2 - 4(1)(13) = 36 - 52 = -16$$

$$x = \frac{-(-6) \pm \sqrt{-16}}{2(1)} = \frac{6 \pm 4i}{2} = 3 \pm 2i$$

The zeros are $3 - 2i$ and $3 + 2i$.



Section 2.7: Complex Zeros of a Quadratic Function

14. $f(x) = 0$


$$x^2 + 4x + 8 = 0$$

$$a = 1, b = 4, c = 8$$

$$b^2 - 4ac = 4^2 - 4(1)(8) = 16 - 32 = -16$$

$$x = \frac{-4 \pm \sqrt{-16}}{2(1)} = \frac{-4 \pm 4i}{2} = -2 \pm 2i$$

The zeros are $-2 - 2i$ and $-2 + 2i$.



15. $f(x) = 0$


$$x^2 - 6x + 10 = 0$$

$$a = 1, b = -6, c = 10$$

$$b^2 - 4ac = (-6)^2 - 4(1)(10) = 36 - 40 = -4$$

$$x = \frac{-(-6) \pm \sqrt{-4}}{2(1)} = \frac{6 \pm 2i}{2} = 3 \pm i$$

The zeros are $3 - i$ and $3 + i$.



16. $f(x) = 0$


$$x^2 - 2x + 5 = 0$$

$$a = 1, b = -2, c = 5$$

$$b^2 - 4ac = (-2)^2 - 4(1)(5) = 4 - 20 = -16$$

$$x = \frac{-(-2) \pm \sqrt{-16}}{2(1)} = \frac{2 \pm 4i}{2} = 1 \pm 2i$$

The zeros are $1 - 2i$ and $1 + 2i$.



17. $f(x) = 0$


$$x^2 - 4x + 1 = 0$$

$$a = 1, b = -4, c = 1$$

$$b^2 - 4ac = (-4)^2 - 4(1)(1) = 16 - 4 = 12$$

$$x = \frac{-(-4) \pm \sqrt{12}}{2(1)} = \frac{4 \pm 2\sqrt{3}}{2} = 2 \pm \sqrt{3}$$

The zeros are $2 - \sqrt{3}$ and $2 + \sqrt{3}$, or approximately 0.27 and 3.73.



18. $f(x) = 0$


$$x^2 + 6x + 1 = 0$$

$$a = 1, b = 6, c = 1$$


$$b^2 - 4ac = 6^2 - 4(1)(1) = 36 - 4 = 32$$

$$x = \frac{-6 \pm \sqrt{32}}{2(1)} = \frac{-6 \pm 4\sqrt{2}}{2} = -3 \pm 2\sqrt{2}$$


The zeros are $-3 - 2\sqrt{2}$ and $-3 + 2\sqrt{2}$, or approximately -5.83 and -0.17.




19. $f(x) = 0$
 $2x^2 + 2x + 1 = 0$
 $a = 2, b = 2, c = 1$
 $b^2 - 4ac = (2)^2 - 4(2)(1) = 4 - 8 = -4$
 $x = \frac{-2 \pm \sqrt{-4}}{2(2)} = \frac{-2 \pm 2i}{4} = -\frac{1}{2} \pm \frac{1}{2}i$
 The zeros are $-\frac{1}{2} - \frac{1}{2}i$ and $-\frac{1}{2} + \frac{1}{2}i$.




20. $f(x) = 0$
 $3x^2 + 6x + 4 = 0$
 $a = 3, b = 6, c = 4$
 $b^2 - 4ac = (6)^2 - 4(3)(4) = 36 - 48 = -12$
 $x = \frac{-6 \pm \sqrt{-12}}{2(3)} = \frac{-6 \pm 2\sqrt{3}i}{6} = -1 \pm \frac{\sqrt{3}}{3}i$
 The zeros are $-1 - \frac{\sqrt{3}}{3}i$ and $-1 + \frac{\sqrt{3}}{3}i$.



21. $f(x) = 0$
 $x^2 + x + 1 = 0$
 $a = 1, b = 1, c = 1$
 $b^2 - 4ac = 1^2 - 4(1)(1) = 1 - 4 = -3$
 $x = \frac{-1 \pm \sqrt{-3}}{2(1)} = \frac{-1 \pm \sqrt{3}i}{2} = -\frac{1}{2} \pm \frac{\sqrt{3}}{2}i$
 The zeros are $-\frac{1}{2} - \frac{\sqrt{3}}{2}i$ and $-\frac{1}{2} + \frac{\sqrt{3}}{2}i$.



22. $f(x) = 0$
 $x^2 - x + 1 = 0$
 $a = 1, b = -1, c = 1$
 $b^2 - 4ac = (-1)^2 - 4(1)(1) = 1 - 4 = -3$
 $x = \frac{-(-1) \pm \sqrt{-3}}{2(1)} = \frac{1 \pm \sqrt{3}i}{2} = \frac{1}{2} \pm \frac{\sqrt{3}}{2}i$
 The zeros are $\frac{1}{2} - \frac{\sqrt{3}}{2}i$ and $\frac{1}{2} + \frac{\sqrt{3}}{2}i$.



Section 2.7: Complex Zeros of a Quadratic Function

23. $f(x) = 0$


$$-2x^2 + 8x + 1 = 0$$

$$a = -2, b = 8, c = 1$$

$$b^2 - 4ac = 8^2 - 4(-2)(1) = 64 + 8 = 72$$

$$x = \frac{-8 \pm \sqrt{72}}{2(-2)} = \frac{-8 \pm 6\sqrt{2}}{-4} = \frac{4 \pm 3\sqrt{2}}{2} = 2 \pm \frac{3\sqrt{2}}{2}$$

The zeros are $\frac{4-3\sqrt{2}}{2}$ and $\frac{4+3\sqrt{2}}{2}$, or approximately -0.12 and 4.12 .



24. $f(x) = 0$


$$-3x^2 + 6x + 1 = 0$$

$$a = -3, b = 6, c = 1$$

$$b^2 - 4ac = 6^2 - 4(-3)(1) = 36 + 12 = 48$$

$$x = \frac{-6 \pm \sqrt{48}}{2(-3)} = \frac{-6 \pm 4\sqrt{3}}{-6} = \frac{3 \pm 2\sqrt{3}}{3} = 1 \pm \frac{2\sqrt{3}}{3}$$

The zeros are $\frac{3-2\sqrt{3}}{3}$ and $\frac{3+2\sqrt{3}}{3}$, or approximately -0.15 and 2.15 .



25. $3x^2 - 3x + 4 = 0$

$$a = 3, b = -3, c = 4$$

$$b^2 - 4ac = (-3)^2 - 4(3)(4) = 9 - 48 = -39$$

The equation has two complex solutions that are conjugates of each other.

26. $2x^2 - 4x + 1 = 0$

$$a = 2, b = -4, c = 1$$

$$b^2 - 4ac = (-4)^2 - 4(2)(1) = 16 - 8 = 8$$

The equation has two unequal real number solutions.

27. $2x^2 + 3x - 4 = 0$

$$a = 2, b = 3, c = -4$$

$$b^2 - 4ac = 3^2 - 4(2)(-4) = 9 + 32 = 41$$

The equation has two unequal real solutions.

28. $x^2 + 2x + 6 = 0$

$$a = 1, b = 2, c = 6$$

$$b^2 - 4ac = (2)^2 - 4(1)(6) = 4 - 24 = -20$$

The equation has two complex solutions that are conjugates of each other.

29. $9x^2 - 12x + 4 = 0$

$$a = 9, b = -12, c = 4$$

$$b^2 - 4ac = (-12)^2 - 4(9)(4) = 144 - 144 = 0$$

The equation has a repeated real solution.

30. $4x^2 + 12x + 9 = 0$

$$a = 4, b = 12, c = 9$$

$$b^2 - 4ac = 12^2 - 4(4)(9) = 144 - 144 = 0$$

The equation has a repeated real solution.

31. $t^4 - 16 = 0$

$$(t^2 - 4)(t^2 + 4) = 0$$

$$t^2 = 4 \quad t^2 = -4$$

$$t = \pm 2 \quad t = \pm 2i$$

32. $y^4 - 81 = 0$

$$(y^2 - 9)(y^2 + 9) = 0$$

$$y^2 = 9 \quad y^2 = -9$$

$$y = \pm 3 \quad y = \pm 3i$$

33. $F(x) = x^6 - 9x^3 + 8 = 0$

$$(x^3 - 8)(x^3 - 1) = 0$$

$$(x - 2)(x^2 + 2x + 4)(x - 1)(x^2 + x + 1) = 0$$

Chapter 2: Linear and Quadratic Functions

$$x^2 + 2x + 4 = 0$$

$$a = 1, b = 2, c = 4$$

$$x = \frac{-2 \pm \sqrt{2^2 - 4(1)(4)}}{2(1)} = \frac{-2 \pm \sqrt{-12}}{2} = \frac{-2 \pm 2i\sqrt{3}}{2}$$

$$= -1 \pm \sqrt{3}i$$

$$x^2 + x + 1 = 0$$

$$a = 1, b = 1, c = 1$$

$$x = \frac{-1 \pm \sqrt{1^2 - 4(1)(1)}}{2(1)} = \frac{-1 \pm \sqrt{-3}}{2} = \frac{-1 \pm i\sqrt{3}}{2}$$

$$= -\frac{1}{2} \pm \frac{\sqrt{3}}{2}i$$

$$\text{The solution set is } \left\{ -1 \pm i\sqrt{3}, -\frac{1}{2} \pm i\frac{\sqrt{3}}{2}, 2, 1 \right\}$$

34. $P(z) = z^6 + 28z^3 + 27 = 0$

$$(z^3 + 27)(z^3 + 1) = 0$$

$$(z + 3)(z^2 - 3z + 9)(z + 1)(z^2 - z + 1) = 0$$

$$z^2 - 3z + 9 = 0$$

$$a = 1, b = -3, c = 9$$

$$z = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(1)(9)}}{2(1)} = \frac{3 \pm \sqrt{-27}}{2}$$

$$= \frac{3 \pm 3i\sqrt{3}}{2} = \frac{3}{2} \pm \frac{3\sqrt{3}}{2}i$$

$$z^2 - z + 1 = 0$$

$$a = 1, b = -1, c = 1$$


$$z = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(1)(1)}}{2(1)} = \frac{1 \pm \sqrt{-3}}{2} = \frac{1 \pm i\sqrt{3}}{2}$$

$$= \frac{1}{2} \pm \frac{\sqrt{3}}{2}i$$

$$\text{The solution set is } \left\{ \frac{3}{2} \pm \frac{3\sqrt{3}}{2}i, \frac{1}{2} \pm \frac{\sqrt{3}}{2}i, -3, -1 \right\}$$

Section 2.8

1. $x \geq -2$



2. The distance on a number line from the origin to a is $|a|$ for any real number a .

3. $4x - 3 = 9$

$$4x = 12$$

$$x = 3$$

The solution set is $\{3\}$.

4. $3x - 2 > 7$

$$3x > 9$$

$$x > 3$$

The solution set is $\{x \mid x > 3\}$ or, using interval notation, $(3, \infty)$.


5. $-1 < 2x + 5 < 13$

$$-6 < 2x < 8$$

$$-3 < x < 4$$

The solution set is $\{x \mid -3 < x < 4\}$ or, using interval notation, $(-3, 4)$.

6. To graph $f(x) = |x - 3|$, shift the graph of $y = |x|$ to the right 3 units.



7. $-a; a$

8. $-a < u < a$

9. $\leq$

10. True

11. False. Any real number will be a solution of $|x| > -2$ since the absolute value of any real number is positive.

12. False. $|u| > a$ is equivalent to $u < -a$ or $u > a$.

13. a. Since the graphs of f and g intersect at the points $(-9, 6)$ and $(3, 6)$, the solution set of $f(x) = g(x)$ is $\{-9, 3\}$.

- b. Since the graph of f is below the graph of g when x is between -9 and 3 , the solution

Section 2.8: Equations and Inequalities Involving the Absolute Value Function

- set of $f(x) \leq g(x)$ is $\{x \mid -9 \leq x \leq 3\}$ or, using interval notation, $[-9, 3]$.
- c. Since the graph of f is above the graph of g to the left of $x = -9$ and to the right of $x = 3$, the solution set of $f(x) > g(x)$ is $\{x \mid x < -9 \text{ or } x > 3\}$ or, using interval notation, $(-\infty, -9) \cup (3, \infty)$.
14. a. Since the graphs of f and g intersect at the points $(0, 2)$ and $(4, 2)$, the solution set of $f(x) = g(x)$ is $\{0, 4\}$.
- b. Since the graph of f is below the graph of g when x is between 0 and 4, the solution set of $f(x) \leq g(x)$ is $\{x \mid 0 \leq x \leq 4\}$ or, using interval notation, $[0, 4]$.
- c. Since the graph of f is above the graph of g to the left of $x = 0$ and to the right of $x = 4$, the solution set of $f(x) > g(x)$ is $\{x \mid x < 0 \text{ or } x > 4\}$ or, using interval notation, $(-\infty, 0) \cup (4, \infty)$.
15. a. Since the graphs of f and g intersect at the points $(-2, 5)$ and $(3, 5)$, the solution set of $f(x) = g(x)$ is $\{-2, 3\}$.
- b. Since the graph of f is above the graph of g to the left of $x = -2$ and to the right of $x = 3$, the solution set of $f(x) \geq g(x)$ is $\{x \mid x \leq -2 \text{ or } x \geq 3\}$ or, using interval notation, $(-\infty, -2] \cup [3, \infty)$.
- c. Since the graph of f is below the graph of g when x is between -2 and 3 , the solution set of $f(x) < g(x)$ is $\{x \mid -2 < x < 3\}$ or, using interval notation, $(-2, 3)$.
16. a. Since the graphs of f and g intersect at the points $(-4, 7)$ and $(3, 7)$, the solution set of $f(x) = g(x)$ is $\{-4, 3\}$.
- b. Since the graph of f is above the graph of g to the left of $x = -4$ and to the right of $x = 3$, the solution set of $f(x) \geq g(x)$ is $\{x \mid x \leq -4 \text{ or } x \geq 3\}$ or, using interval notation, $(-\infty, -4] \cup [3, \infty)$.
- c. Since the graph of f is below the graph of g when x is between -4 and 3 , the solution set of $f(x) < g(x)$ is $\{x \mid -4 < x < 3\}$ or, using interval notation, $(-4, 3)$.
17. $|x| = 6$
 $x = 6$ or $x = -6$
 The solution set is $\{-6, 6\}$.
18. $|x| = 12$
 $x = 12$ or $x = -12$
 The solution set is $\{-12, 12\}$.
19. $|2x + 3| = 5$
 $2x + 3 = 5$ or $2x + 3 = -5$
 $2x = 2$ or $2x = -8$
 $x = 1$ or $x = -4$
 The solution set is $\{-4, 1\}$.
20. $|3x - 1| = 2$
 $3x - 1 = 2$ or $3x - 1 = -2$
 $3x = 3$ or $3x = -1$
 $x = 1$ or $x = -\frac{1}{3}$
 The solution set is $\{-\frac{1}{3}, 1\}$.
21. $|1 - 4t| + 8 = 13$
 $|1 - 4t| = 5$
 $1 - 4t = 5$ or $1 - 4t = -5$
 $-4t = 4$ or $-4t = -6$
 $t = -1$ or $t = \frac{3}{2}$
 The solution set is $\{-1, \frac{3}{2}\}$.
22. $|1 - 2z| + 6 = 9$
 $|1 - 2z| = 3$
 $1 - 2z = 3$ or $1 - 2z = -3$
 $-2z = 2$ or $-2z = -4$
 $z = -1$ or $z = 2$
 The solution set is $\{-1, 2\}$.
23. $|-2x| = 8$
 $-2x = 8$ or $-2x = -8$
 $x = -4$ or $x = 4$
 The solution set is $\{-4, 4\}$.

Chapter 2: Linear and Quadratic Functions

24. $|-x| = 1$

$-x = 1 \quad \text{or} \quad -x = -1$

The solution set is $\{-1, 1\}$.

25. $4 - |2x| = 3$

$-|2x| = -1$

$|2x| = 1$

$2x = 1 \quad \text{or} \quad 2x = -1$

$x = \frac{1}{2} \quad \text{or} \quad x = -\frac{1}{2}$

The solution set is $\left\{-\frac{1}{2}, \frac{1}{2}\right\}$.

26. $5 - \left|\frac{1}{2}x\right| = 3$

$-\left|\frac{1}{2}x\right| = -2$

$\left|\frac{1}{2}x\right| = 2$

$\frac{1}{2}x = 2 \quad \text{or} \quad \frac{1}{2}x = -2$

$x = 4 \quad \text{or} \quad x = -4$

The solution set is $\{-4, 4\}$.

27. $\frac{2}{3}|x| = 9$

$|x| = \frac{27}{2}$

$x = \frac{27}{2} \quad \text{or} \quad x = -\frac{27}{2}$

The solution set is $\left\{-\frac{27}{2}, \frac{27}{2}\right\}$.

28. $\frac{3}{4}|x| = 9$

$|x| = 12$

$x = 12 \quad \text{or} \quad x = -12$

The solution set is $\{-12, 12\}$.

29. $\left|\frac{x}{3} + \frac{2}{5}\right| = 2$

$\frac{x}{3} + \frac{2}{5} = 2 \quad \text{or} \quad \frac{x}{3} + \frac{2}{5} = -2$

$5x + 6 = 30 \quad \text{or} \quad 5x + 6 = -30$

$5x = 24 \quad \text{or} \quad 5x = -36$

$x = \frac{24}{5} \quad \text{or} \quad x = -\frac{36}{5}$

The solution set is $\left\{-\frac{36}{5}, \frac{24}{5}\right\}$.

30. $\left|\frac{x}{2} - \frac{1}{3}\right| = 1$

$\frac{x}{2} - \frac{1}{3} = 1 \quad \text{or} \quad \frac{x}{2} - \frac{1}{3} = -1$

$3x - 2 = 6 \quad \text{or} \quad 3x - 2 = -6$

$3x = 8 \quad \text{or} \quad 3x = -4$

$x = \frac{8}{3} \quad \text{or} \quad x = -\frac{4}{3}$

The solution set is $\left\{-\frac{4}{3}, \frac{8}{3}\right\}$.

31. $|u - 2| = -\frac{1}{2}$

No solution, since absolute value always yields a non-negative number.

32. $|2 - v| = -1$

No solution, since absolute value always yields a non-negative number.

33. $|x^2 - 9| = 0$

$x^2 - 9 = 0$

$x^2 = 9$

$x = \pm 3$

The solution set is $\{-3, 3\}$.

34. $|x^2 - 16| = 0$

$x^2 - 16 = 0$

$x^2 = 16$

$x = \pm 4$

The solution set is $\{-4, 4\}$.

Section 2.8: Equations and Inequalities Involving the Absolute Value Function

35. $|x^2 - 2x| = 3$

$$x^2 - 2x = 3 \quad \text{or} \quad x^2 - 2x = -3$$

$$x^2 - 2x - 3 = 0 \quad \text{or} \quad x^2 - 2x + 3 = 0$$

$$(x-3)(x+1) = 0 \quad \text{or} \quad x = \frac{2 \pm \sqrt{4-12}}{2}$$

$$= \frac{2 \pm \sqrt{-8}}{2} \quad \text{no real sol.}$$

$$x = 3 \quad \text{or} \quad x = -1$$

The solution set is $\{-1, 3\}$.

36. $|x^2 + x| = 12$

$$x^2 + x = 12 \quad \text{or} \quad x^2 + x = -12$$

$$x^2 + x - 12 = 0 \quad \text{or} \quad x^2 + x + 12 = 0$$

$$(x-3)(x+4) = 0 \quad \text{or} \quad x = \frac{-1 \pm \sqrt{1-48}}{2}$$

$$= \frac{1 \pm \sqrt{-47}}{2} \quad \text{no real sol.}$$

$$x = 3 \quad \text{or} \quad x = -4$$

The solution set is $\{-4, 3\}$.

37. $|x^2 + x - 1| = 1$

$$x^2 + x - 1 = 1 \quad \text{or} \quad x^2 + x - 1 = -1$$

$$x^2 + x - 2 = 0 \quad \text{or} \quad x^2 + x = 0$$

$$(x-1)(x+2) = 0 \quad \text{or} \quad x(x+1) = 0$$

$$x = 1, x = -2 \quad \text{or} \quad x = 0, x = -1$$

The solution set is $\{-2, -1, 0, 1\}$.

38. $|x^2 + 3x - 2| = 2$

$$x^2 + 3x - 2 = 2 \quad \text{or} \quad x^2 + 3x - 2 = -2$$

$$x^2 + 3x = 4 \quad \text{or} \quad x^2 + 3x = 0$$

$$x^2 + 3x - 4 = 0 \quad \text{or} \quad x(x+3) = 0$$

$$(x+4)(x-1) = 0 \quad \text{or} \quad x = 0, x = -3$$


$$x = -4, x = 1$$

The solution set is $\{-4, -3, 0, 1\}$.

39. $|x| < 6$

$$-6 < x < 6$$


$$\{x | -6 < x < 6\} \quad \text{or} \quad (-6, 6)$$



40. $|x| < 9$

$$-9 < x < 9$$


$$\{x | -9 < x < 9\} \quad \text{or} \quad (-9, 9)$$



41. $|x| > 4$

$$x < -4 \quad \text{or} \quad x > 4$$


$$\{x | x < -4 \text{ or } x > 4\} \quad \text{or} \quad (-\infty, -4) \cup (4, \infty)$$



42. $|x| > 1$

$$x < -1 \quad \text{or} \quad x > 1$$

$$\{x | x < -1 \text{ or } x > 1\} \quad \text{or} \quad (-\infty, -1) \cup (1, \infty)$$




43. $|2x| < 8$

$$-8 < 2x < 8$$

$$-4 < x < 4$$

$$\{x | -4 < x < 4\} \quad \text{or} \quad (-4, 4)$$



44. $|3x| < 15$

$$-15 < 3x < 15$$

$$-5 < x < 5$$

$$\{x | -5 < x < 5\} \quad \text{or} \quad (-5, 5)$$




45. $|3x| > 12$

$$3x < -12 \quad \text{or} \quad 3x > 12$$

$$x < -4 \quad \text{or} \quad x > 4$$

$$\{x | x < -4 \text{ or } x > 4\} \quad \text{or} \quad (-\infty, -4) \cup (4, \infty)$$




46. $|2x| > 6$

$$2x < -6 \quad \text{or} \quad 2x > 6$$

$$x < -3 \quad \text{or} \quad x > 3$$

$$\{x | x < -3 \text{ or } x > 3\} \quad \text{or} \quad (-\infty, -3) \cup (3, \infty)$$



Chapter 2: Linear and Quadratic Functions


47. $|x-2|+2<3$

$$|x-2|<1$$

$$-1<x-2<1$$

$$1<x<3$$

$$\{x|1<x<3\} \text{ or } (1,3)$$




48. $|x+4|+3<5$

$$|x+4|<2$$

$$-2<x+4<2$$

$$-6<x<-2$$

$$\{x|-6<x<-2\} \text{ or } (-6,-2)$$




49. $|3t-2|\leq 4$

$$-4\leq 3t-2\leq 4$$

$$-2\leq 3t\leq 6$$

$$-\frac{2}{3}\leq t\leq 2$$

$$\left\{t\left|-\frac{2}{3}\leq t\leq 2\right.\right\} \text{ or } \left[-\frac{2}{3}, 2\right]$$




50. $|2u+5|\leq 7$

$$-7\leq 2u+5\leq 7$$

$$-12\leq 2u\leq 2$$

$$-6\leq u\leq 1$$

$$\{u|-6\leq u\leq 1\} \text{ or } [-6, 1]$$




51. $|x-3|\geq 2$

$$x-3\leq -2 \text{ or } x-3\geq 2$$

$$x\leq 1 \text{ or } x\geq 5$$

$$\{x|x\leq 1 \text{ or } x\geq 5\} \text{ or } (-\infty, 1]\cup[5, \infty)$$




52. $|x+4|\geq 2$

$$x+4\leq -2 \text{ or } x+4\geq 2$$

$$x\leq -6 \text{ or } x\geq -2$$

$$\{x|x\leq -6 \text{ or } x\geq -2\} \text{ or } (-\infty, -6]\cup[-2, \infty)$$



53. $|1-4x|-7<-2$

$$|1-4x|<5$$


$$-5<1-4x<5$$

$$-6<-4x<4$$

$$-\frac{6}{4}>x>\frac{4}{-4}$$

$$\frac{3}{2}>x>-1 \text{ or } -1<x<\frac{3}{2}$$

$$\{x|-1<x<\frac{3}{2}\} \text{ or } (-1, \frac{3}{2})$$



54. $|1-2x|-4<-1$

$$|1-2x|<3$$


$$-3<1-2x<3$$

$$-4<-2x<2$$

$$-\frac{4}{-2}>x>\frac{2}{-2}$$

$$2>x>-1 \text{ or } -1<x<2$$

$$\{x|-1<x<2\} \text{ or } (-1, 2)$$



55. $|1-2x|>|-3|$


$$|1-2x|>3$$

$$1-2x<-3 \text{ or } 1-2x>3$$

$$-2x<-4 \text{ or } -2x>2$$

$$x>2 \text{ or } x<-1$$

$$\{x|x<-1 \text{ or } x>2\} \text{ or } (-\infty, -1)\cup(2, \infty)$$



Section 2.8: Equations and Inequalities Involving the Absolute Value Function

56. $|2-3x| > |-1|$


$$|2-3x| > 1$$

$$2-3x < -1 \quad \text{or} \quad 2-3x > 1$$

$$-3x < -3 \quad \text{or} \quad -3x > -1$$


$$x > 1 \quad \text{or} \quad x < \frac{1}{3}$$

$$\left\{x \mid x < \frac{1}{3} \text{ or } x > 1\right\} \text{ or } \left(-\infty, \frac{1}{3}\right) \cup (1, \infty)$$



57. $|2x+1| < -1$


No solution since absolute value is always non-negative.



58. $|3x-4| \geq 0$

All real numbers since absolute value is always non-negative.

$$\{x \mid x \text{ is any real number}\} \text{ or } (-\infty, \infty)$$



59. $|(3x-2)-7| < \frac{1}{2}$


$$|3x-9| < \frac{1}{2}$$

$$-\frac{1}{2} < 3x-9 < \frac{1}{2}$$

$$\frac{17}{2} < 3x < \frac{19}{2}$$

$$\frac{17}{6} < x < \frac{19}{6}$$

$$\left\{x \mid \frac{17}{6} < x < \frac{19}{6}\right\} \text{ or } \left(\frac{17}{6}, \frac{19}{6}\right)$$



60. $|(4x-1)-11| < \frac{1}{4}$


$$|4x-12| < \frac{1}{4}$$

$$-\frac{1}{4} < 4x-12 < \frac{1}{4}$$

$$\frac{47}{4} < 4x < \frac{49}{4}$$

$$\frac{47}{16} < x < \frac{49}{16}$$

$$\left\{x \mid \frac{47}{16} < x < \frac{49}{16}\right\} \text{ or } \left(\frac{47}{16}, \frac{49}{16}\right)$$



61. $5-|x-1| > 2$


$$-|x-1| > -3$$

$$|x-1| < 3$$

$$-3 < x-1 < 3$$

$$-2 < x < 4$$

$$\{x \mid -2 < x < 4\} \text{ or } (-2, 4)$$



62. $6-|x+3| \geq 2$


$$-|x+3| \geq -4$$

$$|x+3| \leq 4$$

$$-4 \leq x+3 \leq 4$$

$$-7 \leq x \leq 1$$

$$\{x \mid -7 \leq x \leq 1\} \text{ or } [-7, 1]$$



63. a. $f(x) = g(x)$

$$-3|5x-2| = -9$$

$$|5x-2| = 3$$

$$5x-2 = 3 \quad \text{or} \quad 5x-2 = -3$$

$$5x = 5 \quad \text{or} \quad 5x = -1$$

$$x = 1 \quad \text{or} \quad x = -\frac{1}{5}$$

Chapter 2: Linear and Quadratic Functions

b. $f(x) > g(x)$

$$-3|5x-2| > -9$$

$$|5x-2| < 3$$

$$-3 < 5x-2 < 3$$

$$-1 < 5x < 5$$

$$-\frac{1}{5} < x < 1$$

$$\left\{x \mid -\frac{1}{5} < x < 1\right\} \text{ or } \left(-\frac{1}{5}, 1\right)$$

c. $f(x) \leq g(x)$

$$-3|5x-2| \leq -9$$

$$|5x-2| \geq 3$$

$$5x-2 \geq 3 \text{ or } 5x-2 \leq -3$$

$$5x \geq 5 \text{ or } 5x \leq -1$$

$$x \geq 1 \text{ or } x \leq -\frac{1}{5}$$

$$\left\{x \mid x \leq -\frac{1}{5} \text{ or } x \geq 1\right\} \text{ or } \left(-\infty, -\frac{1}{5}\right] \cup [1, \infty)$$

64. a. $f(x) = g(x)$

$$-2|2x-3| = -12$$

$$|2x-3| = 6$$

$$2x-3 = 6 \text{ or } 2x-3 = -6$$

$$2x = 9 \text{ or } 2x = -3$$

$$x = \frac{9}{2} \text{ or } x = -\frac{3}{2}$$

b. $f(x) = g(x)$

$$-2|2x-3| \geq -12$$

$$|2x-3| \leq 6$$

$$-6 \leq 2x-3 \leq 6$$

$$-3 \leq 2x \leq 9$$

$$-\frac{3}{2} \leq x \leq \frac{9}{2}$$

$$\left\{x \mid -\frac{3}{2} \leq x \leq \frac{9}{2}\right\} \text{ or } \left[-\frac{3}{2}, \frac{9}{2}\right]$$

c. $f(x) = g(x)$

$$-2|2x-3| < -12$$

$$|2x-3| > 6$$

$$2x-3 > 6 \text{ or } 2x-3 < -6$$

$$2x > 9 \text{ or } 2x < -3$$

$$x > \frac{9}{2} \text{ or } x < -\frac{3}{2}$$

$$\left\{x \mid x < -\frac{3}{2} \text{ or } x > \frac{9}{2}\right\} \text{ or } \left(-\infty, -\frac{3}{2}\right) \cup \left(\frac{9}{2}, \infty\right)$$

65. a. $f(x) = g(x)$

$$|-3x+2| = x+10$$


$$-3x+2 = x+10 \text{ or } -3x+2 = -(x+10)$$

$$-4x = 8 \text{ or } -3x+2 = -x-10$$

$$x = -2 \text{ or } -2x = -12$$

$$\text{or } x = 6$$

b.



Look at the graph of $f(x)$ and $g(x)$ and see where the graph of $f(x) \geq g(x)$. We see that this occurs where $x \leq -2$ or $x \geq 6$.

So the solution set is: $\{x \mid x \leq -2 \text{ or } x \geq 6\}$

or $(-\infty, -2] \cup [6, \infty)$.

c. Look at the graph of $f(x)$ and $g(x)$ and see where the graph of $f(x) < g(x)$. We see that this occurs where x is between -2 and 6. So the solution set is:

$\{x \mid -2 < x < 6\}$ or $(-2, 6)$.

66. a. $f(x) = g(x)$

$$|4x-3| = x+2$$


$$4x-3 = x+2 \text{ or } 4x-3 = -(x+2)$$

$$3x = 5 \text{ or } 4x-3 = -x-2$$

$$x = \frac{5}{3} \text{ or } 5x = 1$$

$$\text{or } x = \frac{1}{5}$$

b.



Section 2.8: Equations and Inequalities Involving the Absolute Value Function

Look at the graph of $f(x)$ and $g(x)$ and see where the graph of $f(x) > g(x)$. We see that this occurs where $x < \frac{1}{5}$ or $x > \frac{5}{3}$.

So the solution set is: $\{x \mid x < \frac{1}{5} \text{ or } x > \frac{5}{3}\}$ or

$$\left(-\infty, \frac{1}{5}\right) \cup \left(\frac{5}{3}, \infty\right)$$

- c. Look at the graph of $f(x)$ and $g(x)$ and see where the graph of $f(x) \leq g(x)$. We see that this occurs where x is between $\frac{1}{5}$ and $\frac{5}{3}$. So the solution set is:

$$\left\{x \mid \frac{1}{5} \leq x \leq \frac{5}{3}\right\} \text{ or } \left[\frac{1}{5}, \frac{5}{3}\right].$$

67. $|x-10| < 2$

$$-2 < x-10 < 2$$

$$8 < x < 12$$

Solution set: $\{x \mid 8 < x < 12\}$ or $(8, 12)$

68. $|x-(-6)| < 3$

$$|x+6| < 3$$

$$-3 < x+6 < 3$$

$$-9 < x < -3$$

Solution set: $\{x \mid -9 < x < -3\}$ or $(-9, -3)$

69. $|2x-(-1)| > 5$

$$|2x+1| > 5$$

$$2x+1 < -5 \text{ or } 2x+1 > 5$$

$$2x < -6 \text{ or } 2x > 4$$

$$x < -3 \text{ or } x > 2$$

Solution set: $\{x \mid x < -3 \text{ or } x > 2\}$ or

$$(-\infty, -3) \cup (2, \infty)$$

70. $|2x-3| > 1$

$$2x-3 < -1 \text{ or } 2x-3 > 1$$

$$2x < 2 \text{ or } 2x > 4$$

$$x < 1 \text{ or } x > 2$$

Solution set: $\{x \mid x < 1 \text{ or } x > 2\}$ or

$$(-\infty, 1) \cup (2, \infty)$$

71. $|x-5.7| \leq 0.0005$

$$-0.0005 < x-5.7 < 0.0005$$

$$5.6995 < x < 5.7005$$

The acceptable lengths of the rod is from 5.6995 inches to 5.7005 inches.

72. $|x-6.125| \leq 0.0005$

$$-0.0005 < x-6.125 < 0.0005$$

$$6.1245 < x < 6.1255$$

The acceptable lengths of the rod is from 6.1245 inches to 6.1255 inches.

73. $\left|\frac{x-100}{15}\right| > 1.96$

$$\frac{x-100}{15} < -1.96 \text{ or } \frac{x-100}{15} > 1.96$$

$$x-100 < -29.4 \text{ or } x-100 > 29.4$$

$$x < 70.6 \text{ or } x > 129.4$$

Since IQ scores are whole numbers, any IQ less than 71 or greater than 129 would be considered unusual.

74. $\left|\frac{x-266}{16}\right| > 1.96$

$$\frac{x-266}{16} < -1.96 \text{ or } \frac{x-266}{16} > 1.96$$

$$x-266 < -31.36 \text{ or } x-266 > 31.36$$

$$x < 234.64 \text{ or } x > 297.36$$

Pregnancies less than 235 days long or greater than 297 days long would be considered unusual.

75. $|5x+1|+7=5$

$$|5x+1| = -2$$

No matter what real number is substituted for x , the absolute value expression on the left side of the equation must always be zero or larger. Thus, it can never equal -2 .

76. $|2x+5|+3 > 1 \Rightarrow |2x+5| > -2$

No matter what real number is substituted for x , the absolute value expression on the left side of the equation must always be zero or larger.

Thus, it will always be larger than -2 . Thus, the solution is the set of all real numbers.

77. $|2x-1| \leq 0$

No matter what real number is substituted for x , the absolute value expression on the left side of

Chapter 2: Linear and Quadratic Functions

the equation must always be zero or larger.
will be when the absolute value expression
equals 0:

$$|2x-1| = 0$$

$$2x-1 = 0$$

$$2x = 1$$


$$x = \frac{1}{2}$$

Thus, the solution set is $\left\{\frac{1}{2}\right\}$.

Chapter 2 Review Exercises

1. $f(x) = 2x - 5$


- Slope = 2; y-intercept = -5
- Plot the point $(0, -5)$. Use the slope to find an additional point by moving 1 unit to the right and 2 units up.



- Domain and Range: $(-\infty, \infty)$
- Average rate of change = slope = 2
- Increasing

2. $g(x) = -4x + 7$

- Slope = -4; y-intercept = 7
- Plot the point $(0, 7)$. Use the slope to find an additional point by moving 1 unit to the right and 4 units down.




Thus, the only solution to the inequality above

- Domain and Range: $(-\infty, \infty)$
- Average rate of change = slope = -4
- Decreasing

3. $h(x) = \frac{4}{5}x - 6$


- Slope = $\frac{4}{5}$; y-intercept = -6
- Plot the point $(0, -6)$. Use the slope to find an additional point by moving 5 units to the right and 4 units up.



- Domain and Range: $(-\infty, \infty)$
- Average rate of change = slope = $\frac{4}{5}$
- Increasing

4. $F(x) = -\frac{1}{3}x + 1$

- Slope = $-\frac{1}{3}$; y-intercept = 1
- Plot the point $(0, 1)$. Use the slope to find an additional point by moving 3 units to the right and 1 unit down.




- Domain and Range: $(-\infty, \infty)$
- Average rate of change = slope = $-\frac{1}{3}$

e. Decreasing

5. $G(x) = 4$

a. Slope = 0; y-intercept = 4

b. Plot the point (0, 4) and draw a horizontal line through it.



c. Domain: $(-\infty, \infty)$

Range: $\{y \mid y = 4\}$


d. Average rate of change = slope = 0

e. Constant

6. $H(x) = -3$

a. Slope = 0; y-intercept = -3

b. Plot the point (0, -3) and draw a horizontal line through it.



c. Domain: $(-\infty, \infty)$

Range: $\{y \mid y = -3\}$

d. Average rate of change = slope = 0

e. Constant


7. $f(x) = 2x + 14$

zero: $f(x) = 2x + 14 = 0$

$$2x = -14$$

$$x = -7$$

y-intercept = 14




8. $f(x) = -\frac{1}{4}x + 2$

zero: $f(x) = -\frac{1}{4}x + 2 = 0$

$$-\frac{1}{4}x = -2$$

$$x = 8$$

y-intercept = 2



9.

x	$y = f(x)$	Avg. rate of change = $\frac{\Delta y}{\Delta x}$
-2	-7	
0	3	$\frac{3 - (-7)}{0 - (-2)} = \frac{10}{2} = 5$
1	8	$\frac{8 - 3}{1 - 0} = \frac{5}{1} = 5$
3	18	$\frac{18 - 8}{3 - 1} = \frac{10}{2} = 5$
6	33	$\frac{33 - 18}{6 - 3} = \frac{15}{3} = 5$

This is a linear function with slope = 5, since the average rate of change is constant at 5. To find the equation of the line, we use the point-slope formula and one of the points.

$$y - y_1 = m(x - x_1)$$

$$y - 3 = 5(x - 0)$$

$$y = 5x + 3$$

Chapter 2: Linear and Quadratic Functions

10.

x	$y = f(x)$	Avg. rate of change = $\frac{\Delta y}{\Delta x}$
-1	-3	
0	4	$\frac{4 - (-3)}{0 - (-1)} = \frac{7}{1} = 7$
1	7	$\frac{7 - 4}{1 - 0} = \frac{3}{1} = 3$
2	6	
3	1	

This is not a linear function, since the average rate of change is not constant.

11. $f(x) = 0$

$$x^2 + x - 72 = 0$$

$$(x+9)(x-8) = 0$$

$$x+9=0 \quad \text{or} \quad x-8=0$$

$$x = -9 \quad \quad \quad x = 8$$

The zeros of $f(x) = x^2 + x - 72$ are -9 and 8.

The x -intercepts of the graph of f are -9 and 8.

12. $G(x) = 0$

$$x^2 - 3x - 40 = 0$$

$$(x+5)(x-8) = 0$$

$$x+5=0 \quad \text{or} \quad x-8=0$$

$$x = -5 \quad \quad \quad x = 8$$

The zeros of $G(x) = x^2 - 3x - 40$ are -5 and 8.

The x -intercepts of the graph of G are -5 and 8.

13. $P(t) = 0$

$$6t^2 - 13t - 5 = 0$$

$$(3t+1)(2t-5) = 0$$

$$3t+1=0 \quad \text{or} \quad 2t-5=0$$

$$t = -\frac{1}{3} \quad \quad \quad t = \frac{5}{2}$$

The zeros of $P(t) = 6t^2 - 13t - 5$ are $-\frac{1}{3}$ and $\frac{5}{2}$.

The t -intercepts of the graph of P are $-\frac{1}{3}$ and $\frac{5}{2}$.

14. $H(z) = 0$

$$3z^2 - 17z + 20 = 0$$

$$(3z-5)(z-4) = 0$$

$$3z-5=0 \quad \text{or} \quad z-4=0$$

$$z = \frac{5}{3} \quad \quad \quad z = 4$$

The zeros of $H(z) = 3z^2 - 17z + 20$ are $\frac{5}{3}$ and 4.

The x -intercepts of the graph of H are $\frac{5}{3}$ and 4.

15. $g(x) = 0$

$$(x-3)^2 - 4 = 0$$

$$(x-3)^2 = 4$$

$$x-3 = \pm\sqrt{4}$$

$$x-3 = \pm 2$$

$$x = 3 \pm 2$$

$$x = 3-2=1 \quad \text{or} \quad x = 3+2=5$$

The zeros of $g(x) = (x-3)^2 - 4$ are 1 and 5. The

x -intercepts of the graph of g are 1 and 5.

16. $F(x) = 0$

$$(x+5)^2 - 9 = 0$$

$$(x+5)^2 = 9$$

$$x+5 = \pm\sqrt{9}$$

$$x+5 = \pm 3$$

$$x = -5 \pm 3$$

$$x = -5-3=-8 \quad \text{or} \quad x = -5+3=-2$$

The zeros of $F(x) = (x+5)^2 - 9$ are -8 and -2.

The x -intercepts of the graph of F are -8 and -2.

17. $h(x) = 0$

$$9x^2 + 6x + 1 = 0$$

$$(3x+1)(3x+1) = 0$$

$$3x+1=0 \quad \text{or} \quad 3x+1=0$$

$$x = -\frac{1}{3} \quad \quad \quad x = -\frac{1}{3}$$

The only zero of $h(x) = 9x^2 + 6x + 1$ is $-\frac{1}{3}$.

The only x -intercept of the graph of h is $-\frac{1}{3}$.

18. $f(x) = 0$

$$4x^2 - 4x + 1 = 0$$

$$(2x-1)(2x-1) = 0$$

$$2x-1=0 \quad \text{or} \quad 2x-1=0$$

$$x = \frac{1}{2} \qquad x = \frac{1}{2}$$

The only zero of $f(x) = 4x^2 - 4x + 1$ is $\frac{1}{2}$. The

only x -intercept of the graph of f is $\frac{1}{2}$.

19. $G(x) = 0$

$$2x^2 - 4x - 1 = 0$$

$$x^2 - 2x - \frac{1}{2} = 0$$

$$x^2 - 2x = \frac{1}{2}$$

$$x^2 - 2x + 1 = \frac{1}{2} + 1$$

$$(x-1)^2 = \frac{3}{2}$$

$$x-1 = \pm\sqrt{\frac{3}{2}} = \pm\frac{\sqrt{3}}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \pm\frac{\sqrt{6}}{2}$$

$$x = 1 \pm \frac{\sqrt{6}}{2} = \frac{2 \pm \sqrt{6}}{2}$$

The zeros of $G(x) = 2x^2 - 4x - 1$ are $\frac{2-\sqrt{6}}{2}$

and $\frac{2+\sqrt{6}}{2}$. The x -intercepts of the graph of G

are $\frac{2-\sqrt{6}}{2}$ and $\frac{2+\sqrt{6}}{2}$.

20. $P(x) = 0$

$$3x^2 + 5x + 1 = 0$$

$$x^2 + \frac{5}{3}x + \frac{1}{3} = 0$$

$$x^2 + \frac{5}{3}x = -\frac{1}{3}$$

$$x^2 + \frac{5}{3}x + \frac{25}{36} = -\frac{1}{3} + \frac{25}{36}$$

$$\left(x + \frac{5}{6}\right)^2 = \frac{13}{36}$$

$$x + \frac{5}{6} = \pm\sqrt{\frac{13}{36}} = \pm\frac{\sqrt{13}}{6}$$

$$x = -\frac{5}{6} \pm \frac{\sqrt{13}}{6} = \frac{-5 \pm \sqrt{13}}{6}$$

The zeros of $P(x) = 3x^2 + 5x + 1$ are $\frac{-5-\sqrt{13}}{6}$

and $\frac{-5+\sqrt{13}}{6}$. The x -intercepts of the graph of

P are $\frac{-5-\sqrt{13}}{6}$ and $\frac{-5+\sqrt{13}}{6}$.

21. $f(x) = 0$

$$-2x^2 + x + 1 = 0$$

$$2x^2 - x - 1 = 0$$

$$(2x+1)(x-1) = 0$$

$$2x+1=0 \quad \text{or} \quad x-1=0$$

$$x = -\frac{1}{2} \qquad x = 1$$

The zeros of $f(x) = -2x^2 + x + 1$ are $-\frac{1}{2}$ and 1.

The x -intercepts of the graph of f are $-\frac{1}{2}$ and 1.

22. $H(x) = 0$

$$-x^2 - 2x + 5 = 0$$

$$x^2 + 2x - 5 = 0$$

$$x^2 + 2x = 5$$

$$x^2 + 2x + 1 = 5 + 1$$

$$(x+1)^2 = 6$$

$$x+1 = \pm\sqrt{6}$$

$$x = -1 \pm \sqrt{6}$$

The zeros of $H(x) = -x^2 - 2x + 5$ are $-1-\sqrt{6}$

and $-1+\sqrt{6}$. The x -intercepts of the graph of H are $-1-\sqrt{6}$ and $-1+\sqrt{6}$.

23. $f(x) = g(x)$

$$(x-3)^2 = 16$$

$$x-3 = \pm\sqrt{16} = \pm 4$$

$$x = 3 \pm 4$$


$$x = 3-4 = -1 \quad \text{or} \quad x = 3+4 = 7$$

The solution set is $\{-1, 7\}$.

The x -coordinates of the points of intersection are -1 and 7 . The y -coordinates are $g(-1) = 16$ and $g(7) = 16$. The graphs of the f and g intersect at

Chapter 2: Linear and Quadratic Functions

the points $(-1, 16)$ and $(7, 16)$.



24. $f(x) = g(x)$

$$x^2 + 6x = 16$$

$$x^2 + 6x - 16 = 0$$


$$(x+8)(x-2) = 0$$

$$x+8=0 \quad \text{or} \quad x-2=0$$

$$x=-8 \quad \quad \quad x=2$$

The solution set is $\{-8, 2\}$.

The x -coordinates of the points of intersection are -8 and 2 . The y -coordinates are $g(-8) = 16$ and $g(2) = 16$. The graphs of the f and g intersect at the points $(-8, 16)$ and $(2, 16)$.



25. $f(x) = g(x)$

$$x^2 + 4x - 5 = 4x - 1$$

$$x^2 - 4 = 0$$

$$(x+2)(x-2) = 0$$


$$x+2=0 \quad \text{or} \quad x-2=0$$

$$x=-2 \quad \quad \quad x=2$$

The solution set is $\{-2, 2\}$.

The x -coordinates of the points of intersection are -2 and 2 . The y -coordinates are $g(-2) = 4(-2) - 1 = -8 - 1 = -9$ and

$g(2) = 4(2) - 1 = 8 - 1 = 7$. The graphs of the f and g intersect at the points $(-2, -9)$ and $(2, 7)$.



26. $f(x) = g(x)$

$$-2x^2 + 3x + 20 = 3x + 12$$

$$-2x^2 + 8 = 0$$

$$x^2 - 4 = 0$$


$$(x+2)(x-2) = 0$$

$$x+2=0 \quad \text{or} \quad x-2=0$$

$$x=-2 \quad \quad \quad x=2$$

The solution set is $\{-2, 2\}$.

The x -coordinates of the points of intersection are -2 and 2 . The y -coordinates are $g(-2) = 3(-2) + 12 = -6 + 12 = 6$ and $g(2) = 3(2) + 12 = 6 + 12 = 18$. The graphs of the f and g intersect at the points $(-2, 6)$ and $(2, 18)$.



27. $f(x) = 0$

$$x^4 - 5x^2 + 4 = 0$$

$$(x^2 - 4)(x^2 - 1) = 0$$

$$x^2 - 4 = 0 \quad \text{or} \quad x^2 - 1 = 0$$

$$x = \pm 2 \quad \text{or} \quad x = \pm 1$$

The zeros of $f(x) = x^4 - 5x^2 + 4$ are -2 , -1 , 1 , and 2 . The x -intercepts of the graph of f are -2 , -1 , 1 , and 2 .

28. $g(x) = 0$

$$3x^4 + 4x^2 + 1 = 0$$

$$(3x^2 + 1)(x^2 + 1) = 0$$

$$3x^2 + 1 = 0 \quad \text{or} \quad x^2 + 1 = 0$$

$$x^2 = -\frac{1}{3} \quad x^2 = -1$$

$$x = \pm\sqrt{-\frac{1}{3}} \quad x = \pm\sqrt{-1}$$

$x = \text{not real}$

The function $g(x) = 3x^4 + 4x^2 + 1$ has no real solution. The graph of g has no x -intercepts.

29. $F(x) = 0$

$$(x-3)^2 - 2(x-3) - 48 = 0$$

$$\text{Let } u = x-3 \rightarrow u^2 = (x-3)^2$$

$$u^2 - 2u - 48 = 0$$

$$(u+6)(u-8) = 0$$

$$u+6 = 0 \quad \text{or} \quad u-8 = 0$$

$$u = -6 \quad u = 8$$

$$x-3 = -6 \quad x-3 = 8$$

$$x = -3 \quad x = 11$$

The zeros of $F(x) = (x-3)^2 - 2(x-3) - 48$ are -3 and 11 . The x -intercepts of the graph of F are -3 and 11 .

30. $G(x) = 0$

$$2(x+4)^2 + 3(x+4) - 14 = 0$$

$$\text{Let } u = x+4 \rightarrow u^2 = (x+4)^2$$

$$2u^2 + 3u - 14 = 0$$

$$(2u+7)(u-2) = 0$$

$$2u+7 = 0 \quad \text{or} \quad u-2 = 0$$

$$u = -\frac{7}{2} \quad u = 2$$

$$x+4 = -\frac{7}{2} \quad x = -2$$

$$x = -\frac{15}{2}$$

The zeros of $G(x) = 2(x+4)^2 + 3(x+4) - 14$ are $-\frac{15}{2}$ and -2 . The x -intercepts of the graph of G are $-\frac{15}{2}$ and -2 .

31. $h(x) = 0$

$$3x - 13\sqrt{x} - 10 = 0$$

$$\text{Let } u = \sqrt{x} \rightarrow u^2 = x$$

$$3u^2 - 13u - 10 = 0$$

$$(3u+2)(u-5) = 0$$

$$3u+2 = 0 \quad \text{or} \quad u-5 = 0$$

$$u = -\frac{2}{3} \quad u = 5$$

$$\sqrt{x} = -\frac{2}{3} \quad x = 5^2 = 25$$

$x = \text{not real}$

$$\begin{aligned} \text{Check: } h(25) &= 3(25) - 13\sqrt{25} - 10 \\ &= 3(25) - 13(5) - 10 \\ &= 75 - 65 - 10 = 0 \end{aligned}$$

The only zero of $h(x) = 3x - 13\sqrt{x} - 10$ is 25 .
The only x -intercept of the graph of h is 25 .

32. $f(x) = 0$

$$\left(\frac{1}{x}\right)^2 - 4\left(\frac{1}{x}\right) - 12 = 0$$

$$\text{Let } u = \frac{1}{x} \rightarrow u^2 = \left(\frac{1}{x}\right)^2$$

$$u^2 - 4u - 12 = 0$$

$$(u+2)(u-6) = 0$$

$$u+2 = 0 \quad \text{or} \quad u-6 = 0$$

$$u = -2 \quad u = 6$$

$$\frac{1}{x} = -2 \quad \frac{1}{x} = 6$$

$$x = -\frac{1}{2} \quad x = \frac{1}{6}$$


The zeros of $f(x) = \left(\frac{1}{x}\right)^2 - 4\left(\frac{1}{x}\right) - 12$ are $-\frac{1}{2}$ and $\frac{1}{6}$. The x -intercepts of the graph of f are $-\frac{1}{2}$ and $\frac{1}{6}$.

33. $f(x) = (x-2)^2 + 2$

Using the graph of $y = x^2$, shift right 2 units,


Chapter 2: Linear and Quadratic Functions

then shift up 2 units.




34. $f(x) = (x+1)^2 - 4$

Using the graph of $y = x^2$, shift left 1 unit, then shift down 4 units.




35. $f(x) = -(x-4)^2$

Using the graph of $y = x^2$, shift the graph 4 units right, then reflect about the x -axis.




36. $f(x) = (x-1)^2 - 3$

Using the graph of $y = x^2$, shift the graph 1 unit right and shift 3 units down.




37. $f(x) = 2(x+1)^2 + 4$

Using the graph of $y = x^2$, stretch vertically by a factor of 2, then shift 1 unit left, then shift 4 units up.



38. $f(x) = -3(x+2)^2 + 1$

Using the graph of $y = x^2$, stretch vertically by a factor of 3, then shift 2 units left, then reflect about the x -axis, then shift 1 unit up.



39. a. $f(x) = (x-2)^2 + 2$

$$= x^2 - 4x + 4 + 2$$

$$= x^2 - 4x + 6$$

$a = 1, b = -4, c = 6$. Since $a = 1 > 0$, the graph opens up. The x -coordinate of the

$$\text{vertex is } x = -\frac{b}{2a} = -\frac{-4}{2(1)} = \frac{4}{2} = 2.$$

The y -coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f(2) = (2)^2 - 4(2) + 6 = 2.$$

Thus, the vertex is $(2, 2)$.


The axis of symmetry is the line $x = 2$.

The discriminant is:

$$b^2 - 4ac = (-4)^2 - 4(1)(6) = -8 < 0, \text{ so the}$$

graph has no x -intercepts.

The y -intercept is $f(0) = 6$.



b. Domain: $(-\infty, \infty)$. Range: $[2, \infty)$.

c. Decreasing on $(-\infty, 2)$; increasing on $(2, \infty)$.

40. a. $f(x) = (x+1)^2 - 4$

$$= x^2 + 2x + 1 - 4$$

$$= x^2 + 2x - 3$$

$a = 1, b = 2, c = -3$. Since $a = 1 > 0$, the graph opens up. The x -coordinate of the

vertex is $x = -\frac{b}{2a} = -\frac{2}{2(1)} = -1$.

The y -coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f(-1) = (-1)^2 + 2(-1) - 3 = -4.$$

Thus, the vertex is $(-1, -4)$.

The axis of symmetry is the line $x = -1$.

The discriminant is:

$$b^2 - 4ac = (2)^2 - 4(1)(-3) = 16 > 0, \text{ so the}$$


graph has two x -intercepts.

The x -intercepts are found by solving:

$$x^2 + 2x - 3 = 0$$

$$(x+3)(x-1) = 0$$

$$x = -3 \text{ or } x = 1$$



b. Domain: $(-\infty, \infty)$. Range: $[-4, \infty)$.

c. Decreasing on $(-\infty, -1)$; increasing on $(-1, \infty)$.

41. a. $f(x) = \frac{1}{4}x^2 - 16$

$a = \frac{1}{4}, b = 0, c = -16$. Since $a = \frac{1}{4} > 0$, the graph opens up. The x -coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{0}{2\left(\frac{1}{4}\right)} = -\frac{0}{\frac{1}{2}} = 0$.

The y -coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f(0) = \frac{1}{4}(0)^2 - 16 = -16.$$

Thus, the vertex is $(0, -16)$.

The axis of symmetry is the line $x = 0$.

The discriminant is:

$$b^2 - 4ac = (0)^2 - 4\left(\frac{1}{4}\right)(-16) = 16 > 0, \text{ so}$$

the graph has two x -intercepts.

The x -intercepts are found by solving:

$$\frac{1}{4}x^2 - 16 = 0$$

$$x^2 - 64 = 0$$


$$x^2 = 64$$

$$x = 8 \text{ or } x = -8$$

The x -intercepts are -8 and 8 .

Chapter 2: Linear and Quadratic Functions

The y-intercept is $f(0) = -16$.



- b. Domain: $(-\infty, \infty)$. Range: $[-16, \infty)$.
 c. Decreasing on $(-\infty, 0)$; increasing on $(0, \infty)$.

42. a. $f(x) = -\frac{1}{2}x^2 + 2$

$a = -\frac{1}{2}$, $b = 0$, $c = 2$. Since $a = -\frac{1}{2} < 0$, the graph opens down. The x-coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{0}{2(-\frac{1}{2})} = -\frac{0}{-1} = 0$.

The y-coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f(0) = -\frac{1}{2}(0)^2 + 2 = 2.$$

The axis of symmetry is the line $x = 0$.
 The discriminant is:

$$b^2 - 4ac = (0)^2 - 4\left(-\frac{1}{2}\right)(2) = 4 > 0, \text{ so the}$$

graph has two x-intercepts.

The x-intercepts are found by solving:

$$-\frac{1}{2}x^2 + 2 = 0$$


$$x^2 - 4 = 0$$

$$x^2 = 4$$

$$x = -2 \text{ or } x = 2$$

The x-intercepts are -2 and 2.

The y-intercept is $f(0) = 2$.



- b. Domain: $(-\infty, \infty)$. Range: $(-\infty, 2]$.
 c. Increasing on $(-\infty, 0)$; decreasing on $(0, \infty)$.

43. a. $f(x) = -4x^2 + 4x$

$a = -4$, $b = 4$, $c = 0$. Since $a = -4 < 0$, the graph opens down. The x-coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{4}{2(-4)} = -\frac{4}{-8} = \frac{1}{2}$.

The y-coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f\left(\frac{1}{2}\right) = -4\left(\frac{1}{2}\right)^2 + 4\left(\frac{1}{2}\right) = -1 + 2 = 1$$

Thus, the vertex is $\left(\frac{1}{2}, 1\right)$.

The axis of symmetry is the line $x = \frac{1}{2}$.

The discriminant is:

$$b^2 - 4ac = 4^2 - 4(-4)(0) = 16 > 0, \text{ so the}$$

graph has two x-intercepts.

The x-intercepts are found by solving:


$$-4x^2 + 4x = 0$$

$$-4x(x-1) = 0$$

$$x = 0 \text{ or } x = 1$$

The x-intercepts are 0 and 1.

The y-intercept is $f(0) = -4(0)^2 + 4(0) = 0$.



b. Domain: $(-\infty, \infty)$. Range: $(-\infty, 1]$.

c. Increasing on $(-\infty, \frac{1}{2})$; decreasing on $(\frac{1}{2}, \infty)$.

44. a. $f(x) = 9x^2 - 6x + 3$
 $a = 9, b = -6, c = 3$. Since $a = 9 > 0$, the graph opens up. The x -coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{-6}{2(9)} = \frac{6}{18} = \frac{1}{3}$. The y -coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f\left(\frac{1}{3}\right) = 9\left(\frac{1}{3}\right)^2 - 6\left(\frac{1}{3}\right) + 3 = 1 - 2 + 3 = 2$$

Thus, the vertex is $\left(\frac{1}{3}, 2\right)$.


The axis of symmetry is the line $x = \frac{1}{3}$.

The discriminant is:

$$b^2 - 4ac = (-6)^2 - 4(9)(3) = -72 < 0, \text{ so the graph has no } x\text{-intercepts.}$$

The y -intercept is

$$f(0) = 9(0)^2 - 6(0) + 3 = 3.$$



b. Domain: $(-\infty, \infty)$. Range: $[2, \infty)$.

c. Decreasing on $(-\infty, \frac{1}{3})$; increasing on $(\frac{1}{3}, \infty)$.

45. a. $f(x) = \frac{9}{2}x^2 + 3x + 1$

$a = \frac{9}{2}, b = 3, c = 1$. Since $a = \frac{9}{2} > 0$, the graph opens up. The x -coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{3}{2\left(\frac{9}{2}\right)} = -\frac{3}{9} = -\frac{1}{3}$.

The y -coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f\left(-\frac{1}{3}\right) = \frac{9}{2}\left(-\frac{1}{3}\right)^2 + 3\left(-\frac{1}{3}\right) + 1 = \frac{1}{2} - 1 + 1 = \frac{1}{2}$$

Thus, the vertex is $\left(-\frac{1}{3}, \frac{1}{2}\right)$.


The axis of symmetry is the line $x = -\frac{1}{3}$.

The discriminant is:

$$b^2 - 4ac = 3^2 - 4\left(\frac{9}{2}\right)(1) = 9 - 18 = -9 < 0,$$

so the graph has no x -intercepts. The y -

intercept is $f(0) = \frac{9}{2}(0)^2 + 3(0) + 1 = 1$.



b. Domain: $(-\infty, \infty)$. Range: $\left[\frac{1}{2}, \infty\right)$.

c. Decreasing on $(-\infty, -\frac{1}{3})$; increasing on $(-\frac{1}{3}, \infty)$.

Chapter 2: Linear and Quadratic Functions

46. a. $f(x) = -x^2 + x + \frac{1}{2}$
 $a = -1, b = 1, c = \frac{1}{2}$. Since $a = -1 < 0$, the graph opens down. The x -coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{1}{2(-1)} = -\frac{1}{-2} = \frac{1}{2}$.
 The y -coordinate of the vertex is


$$f\left(-\frac{b}{2a}\right) = f\left(\frac{1}{2}\right) = -\left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right) + \frac{1}{2}$$

$$= -\frac{1}{4} + 1 = \frac{3}{4}$$

 Thus, the vertex is $\left(\frac{1}{2}, \frac{3}{4}\right)$. The axis of symmetry is the line $x = \frac{1}{2}$. The discriminant is: $b^2 - 4ac = 1^2 - 4(-1)\left(\frac{1}{2}\right) = 3 > 0$, so the graph has two x -intercepts. The x -intercepts are found by solving: $-x^2 + x + \frac{1}{2} = 0$.

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-1 \pm \sqrt{3}}{2(-1)} = \frac{-1 \pm \sqrt{3}}{-2} = \frac{1 \pm \sqrt{3}}{2}$$

 The x -intercepts are $\frac{1 - \sqrt{3}}{2} \approx -0.4$ and $\frac{1 + \sqrt{3}}{2} \approx 1.4$.
 The y -intercept is $f(0) = -(0)^2 + (0) + \frac{1}{2} = \frac{1}{2}$.



- b. Domain: $(-\infty, \infty)$. Range: $\left(-\infty, \frac{3}{4}\right]$.
 c. Increasing on $\left(-\infty, \frac{1}{2}\right)$; decreasing on $\left(\frac{1}{2}, \infty\right)$.

47. a. $f(x) = 3x^2 + 4x - 1$
 $a = 3, b = 4, c = -1$. Since $a = 3 > 0$, the graph opens up. The x -coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{4}{2(3)} = -\frac{4}{6} = -\frac{2}{3}$.
 The y -coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f\left(-\frac{2}{3}\right) = 3\left(-\frac{2}{3}\right)^2 + 4\left(-\frac{2}{3}\right) - 1$$

$$= \frac{4}{3} - \frac{8}{3} - 1 = -\frac{7}{3}$$

 Thus, the vertex is $\left(-\frac{2}{3}, -\frac{7}{3}\right)$.

The axis of symmetry is the line $x = -\frac{2}{3}$.

The discriminant is:

$b^2 - 4ac = (4)^2 - 4(3)(-1) = 28 > 0$, so the graph has two x -intercepts.

The x -intercepts are found by solving:

$$3x^2 + 4x - 1 = 0.$$


$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-4 \pm \sqrt{28}}{2(3)}$$

$$= \frac{-4 \pm 2\sqrt{7}}{6} = \frac{-2 \pm \sqrt{7}}{3}$$

The x -intercepts are $\frac{-2 - \sqrt{7}}{3} \approx -1.55$ and

$$\frac{-2 + \sqrt{7}}{3} \approx 0.22.$$

The y -intercept is $f(0) = 3(0)^2 + 4(0) - 1 = -1$.



- b. Domain: $(-\infty, \infty)$. Range: $\left[-\frac{7}{3}, \infty\right)$.

- c. Decreasing on $\left(-\infty, -\frac{2}{3}\right)$; increasing on $\left(-\frac{2}{3}, \infty\right)$.

48. a. $f(x) = -2x^2 - x + 4$
 $a = -2, b = -1, c = 4$. Since $a = -2 < 0$, the graph opens down. The x -coordinate of the vertex is $x = -\frac{b}{2a} = -\frac{-1}{2(-2)} = \frac{1}{-4} = -\frac{1}{4}$.

The y -coordinate of the vertex is

$$\begin{aligned} f\left(-\frac{b}{2a}\right) &= f\left(-\frac{1}{4}\right) = -2\left(-\frac{1}{4}\right)^2 - \left(-\frac{1}{4}\right) + 4 \\ &= -\frac{1}{8} + \frac{1}{4} + 4 = \frac{33}{8} \end{aligned}$$

Thus, the vertex is $\left(-\frac{1}{4}, \frac{33}{8}\right)$.

The axis of symmetry is the line $x = -\frac{1}{4}$.


The discriminant is:

$b^2 - 4ac = (-1)^2 - 4(-2)(4) = 33 > 0$, so the graph has two x -intercepts. The x -intercepts are found by solving: $-2x^2 - x + 4 = 0$.

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-1) \pm \sqrt{33}}{2(-2)} \\ &= \frac{1 \pm \sqrt{33}}{-4} = \frac{-1 \pm \sqrt{33}}{4} \end{aligned}$$

The x -intercepts are $\frac{-1 - \sqrt{33}}{4} \approx -1.7$ and $\frac{-1 + \sqrt{33}}{4} \approx 1.2$.

The y -intercept is $f(0) = -2(0)^2 - 0 + 4 = 4$.



- b. Domain: $(-\infty, \infty)$. Range: $\left(-\infty, \frac{33}{8}\right]$.

- c. Increasing on $\left(-\infty, -\frac{1}{4}\right)$; decreasing on $\left(-\frac{1}{4}, \infty\right)$.

49. $f(x) = 3x^2 - 6x + 4$
 $a = 3, b = -6, c = 4$. Since $a = 3 > 0$, the graph opens up, so the vertex is a minimum point.

The minimum occurs at

$$x = -\frac{b}{2a} = -\frac{-6}{2(3)} = \frac{6}{6} = 1.$$

The minimum value is

$$\begin{aligned} f\left(-\frac{b}{2a}\right) &= f(1) = 3(1)^2 - 6(1) + 4 \\ &= 3 - 6 + 4 = 1 \end{aligned}$$

50. $f(x) = 2x^2 + 8x + 5$
 $a = 2, b = 8, c = 5$. Since $a = 2 > 0$, the graph opens up, so the vertex is a minimum point.

The minimum occurs at

$$x = -\frac{b}{2a} = -\frac{8}{2(2)} = -\frac{8}{4} = -2.$$

The minimum value is

$$\begin{aligned} f\left(-\frac{b}{2a}\right) &= f(-2) = 2(-2)^2 + 8(-2) + 5 \\ &= 8 - 16 + 5 = -3 \end{aligned}$$

51. $f(x) = -x^2 + 8x - 4$
 $a = -1, b = 8, c = -4$. Since $a = -1 < 0$, the graph opens down, so the vertex is a maximum point. The maximum occurs at

$$x = -\frac{b}{2a} = -\frac{8}{2(-1)} = -\frac{8}{-2} = 4.$$

The maximum value is

$$\begin{aligned} f\left(-\frac{b}{2a}\right) &= f(4) = -(4)^2 + 8(4) - 4 \\ &= -16 + 32 - 4 = 12 \end{aligned}$$

52. $f(x) = -x^2 - 10x - 3$
 $a = -1, b = -10, c = -3$. Since $a = -1 < 0$, the graph opens down, so the vertex is a maximum point. The maximum occurs at

$$x = -\frac{b}{2a} = -\frac{-10}{2(-1)} = \frac{10}{-2} = -5.$$

Chapter 2: Linear and Quadratic Functions

The maximum value is

$$f\left(-\frac{b}{2a}\right) = f(-5) = -(-5)^2 - 10(-5) - 3 \\ = -25 + 50 - 3 = 22$$

53. $f(x) = -3x^2 + 12x + 4$

$a = -3, b = 12, c = 4$. Since $a = -3 < 0$, the graph opens down, so the vertex is a maximum point. The maximum occurs at

$$x = -\frac{b}{2a} = -\frac{12}{2(-3)} = -\frac{12}{-6} = 2.$$

The maximum value is

$$f\left(-\frac{b}{2a}\right) = f(2) = -3(2)^2 + 12(2) + 4 \\ = -12 + 24 + 4 = 16$$

54. $f(x) = -2x^2 + 4$

$a = -2, b = 0, c = 4$. Since $a = -2 < 0$, the graph opens down, so the vertex is a maximum point. The maximum occurs at

$$x = -\frac{b}{2a} = -\frac{0}{2(-2)} = 0.$$

The maximum value is

$$f\left(-\frac{b}{2a}\right) = f(0) = -2(0)^2 + 4 = 4.$$

55. $x^2 + 6x - 16 < 0$

$$f(x) = x^2 + 6x - 16$$

$$x^2 + 6x - 16 = 0$$

$$(x+8)(x-2) = 0$$

$x = -8, x = 2$ are the zeros of f .

Interval	$(-\infty, -8)$	$(-8, 2)$	$(2, \infty)$
Test Number	-9	0	3
Value of f	11	-16	11
Conclusion	Positive	Negative	Positive

The solution set is $\{x \mid -8 < x < 2\}$ or, using interval notation, $(-8, 2)$.

56. $3x^2 - 2x - 1 \geq 0$

$$f(x) = 3x^2 - 2x - 1$$

$$3x^2 - 2x - 1 = 0$$

$$(3x+1)(x-1) = 0$$

$x = -\frac{1}{3}, x = 1$ are the zeros of f .

Interval	$\left(-\infty, -\frac{1}{3}\right)$	$\left(-\frac{1}{3}, 1\right)$	$(1, \infty)$
Test Number	-1	0	2
Value of f	4	-1	7
Conclusion	Positive	Negative	Positive

The solution set is $\left\{x \mid x \leq -\frac{1}{3} \text{ or } x \geq 1\right\}$ or, using interval notation, $\left(-\infty, -\frac{1}{3}\right] \cup [1, \infty)$.

57. $3x^2 \geq 14x + 5$

$$3x^2 - 14x - 5 \geq 0$$

$$f(x) = 3x^2 - 14x - 5$$

$$3x^2 - 14x - 5 = 0$$

$$(3x+1)(x-5) = 0$$

$x = -\frac{1}{3}, x = 5$ are the zeros of f .

Interval	$\left(-\infty, -\frac{1}{3}\right)$	$\left(-\frac{1}{3}, 5\right)$	$(5, \infty)$
Test Number	-1	0	2
Value of f	12	-5	19
Conclusion	Positive	Negative	Positive

The solution set is $\left\{x \mid x \leq -\frac{1}{3} \text{ or } x \geq 5\right\}$ or, using interval notation, $\left(-\infty, -\frac{1}{3}\right] \cup [5, \infty)$.

58. $4x^2 < 13x - 3$

$$4x^2 - 13x + 3 < 0$$

$$f(x) = 4x^2 - 13x + 3$$

$$4x^2 - 13x + 3 = 0$$

$$(4x-1)(x-3) = 0$$

$x = \frac{1}{4}, x = 3$ are the zeros of f .

Interval	$\left(-\infty, \frac{1}{4}\right)$	$\left(\frac{1}{4}, 3\right)$	$(3, \infty)$
Test Number	0	1	4
Value of f	0	-6	15
Conclusion	Positive	Negative	Positive

The solution set is $\left\{x \mid \frac{1}{4} < x < 3\right\}$ or, using interval notation, $\left(\frac{1}{4}, 3\right)$.


59. $f(x) = 0$

$$x^2 + 8 = 0$$

$$x^2 = -8$$

$$x = \pm\sqrt{-8} = \pm 2\sqrt{2}i$$

The zeros are $-2\sqrt{2}i$ and $2\sqrt{2}i$.




60. $f(x) = 0$

$$x^2 - 8 = 0$$

$$x^2 = 8$$

$$x = \pm\sqrt{8} = \pm 2\sqrt{2}$$

The zeros are $-2\sqrt{2}$ and $2\sqrt{2}$.



61. $g(x) = 0$


$$x^2 + 2x - 4 = 0$$

$$a = 1, b = 2, c = -4$$

$$b^2 - 4ac = 2^2 - 4(1)(-4) = 4 + 16 = 20$$

$$x = \frac{-2 \pm \sqrt{20}}{2(1)} = \frac{-2 \pm 2\sqrt{5}}{2} = -1 \pm \sqrt{5}$$

The zeros are $-1 - \sqrt{5}$ and $-1 + \sqrt{5}$.



62. $h(x) = 0$


$$x^2 + 4x + 5 = 0$$

$$a = 1, b = 4, c = 5$$

$$b^2 - 4ac = 4^2 - 4(1)(5) = 16 - 20 = -4$$

$$x = \frac{-4 \pm \sqrt{-4}}{2(1)} = \frac{-4 \pm 2i}{2} = -2 \pm i$$

The zeros are $-2 - i$ and $-2 + i$.



63. $p(x) = 0$


$$-2x^2 + 4x - 3 = 0$$

$$a = -2, b = 4, c = -3$$

$$b^2 - 4ac = 4^2 - 4(-2)(-3) = 16 - 24 = -8$$

$$x = \frac{-4 \pm \sqrt{-8}}{2(-2)} = \frac{-4 \pm 2\sqrt{2}i}{-4} = 1 \pm \frac{\sqrt{2}}{2}i$$

The zeros are $1 - \frac{\sqrt{2}}{2}i$ and $1 + \frac{\sqrt{2}}{2}i$.



Chapter 2: Linear and Quadratic Functions

64. $F(x) = 0$


$$-3x^2 + 6x - 5 = 0$$

$$a = -3, b = 6, c = -5$$

$$b^2 - 4ac = 6^2 - 4(-3)(-5) = 36 - 60 = -24$$

$$x = \frac{-6 \pm \sqrt{-24}}{2(-3)} = \frac{-6 \pm 2\sqrt{6}i}{-6} = 1 \pm \frac{\sqrt{6}}{3}i$$

The zeros are $1 - \frac{\sqrt{6}}{3}i$ and $1 + \frac{\sqrt{6}}{3}i$.



65. $f(x) = 0$


$$4x^2 + 4x + 3 = 0$$

$$a = 4, b = 4, c = 3$$

$$b^2 - 4ac = 4^2 - 4(4)(3) = 16 - 48 = -32$$

$$x = \frac{-4 \pm \sqrt{-32}}{2(4)} = \frac{-4 \pm 4\sqrt{2}i}{8} = -\frac{1}{2} \pm \frac{\sqrt{2}}{2}i$$

The zeros are $-\frac{1}{2} - \frac{\sqrt{2}}{2}i$ and $-\frac{1}{2} + \frac{\sqrt{2}}{2}i$.



66. $G(x) = 0$


$$5x^2 + 5x + 2 = 0$$

$$a = 5, b = 5, c = 2$$

$$b^2 - 4ac = 5^2 - 4(5)(2) = 25 - 40 = -15$$

$$x = \frac{-5 \pm \sqrt{-15}}{2(5)} = \frac{-5 \pm \sqrt{15}i}{10} = -\frac{1}{2} \pm \frac{\sqrt{15}}{10}i$$

The zeros are $-\frac{1}{2} - \frac{\sqrt{15}}{10}i$ and $-\frac{1}{2} + \frac{\sqrt{15}}{10}i$.



67. $|2x + 3| = 7$

$$2x + 3 = 7 \text{ or } 2x + 3 = -7$$

$$2x = 4 \text{ or } 2x = -10$$

$$x = 2 \text{ or } x = -5$$

The solution set is $\{-5, 2\}$.

68. $|3x - 1| = 5$

$$3x - 1 = 5 \text{ or } 3x - 1 = -5$$

$$3x = 6 \text{ or } 3x = -4$$

$$x = 2 \text{ or } x = -\frac{4}{3}$$

The solution set is $\{-\frac{4}{3}, 2\}$.

69. $|2 - 3x| + 2 = 9$

$$|2 - 3x| = 7$$

$$2 - 3x = 7 \text{ or } 2 - 3x = -7$$

$$-3x = 5 \text{ or } -3x = -9$$

$$x = -\frac{5}{3} \text{ or } x = 3$$

The solution set is $\{-\frac{5}{3}, 3\}$.

70. $|1-2x|+1=4$

$$|1-2x|=3$$

$$1-2x=3 \quad \text{or} \quad 1-2x=-3$$

$$-2x=2 \quad \text{or} \quad -2x=-4$$

$$x=-1 \quad \text{or} \quad x=2$$

The solution set is $\{-1, 2\}$.


71. $|3x+4|<\frac{1}{2}$

$$-\frac{1}{2}<3x+4<\frac{1}{2}$$

$$-\frac{9}{2}<3x<-\frac{7}{2}$$

$$-\frac{3}{2}<x<-\frac{7}{6}$$

$$\left\{x \mid -\frac{3}{2}<x<-\frac{7}{6}\right\} \text{ or } \left(-\frac{3}{2}, -\frac{7}{6}\right)$$




72. $|1-2x|<\frac{1}{3}$

$$-\frac{1}{3}<1-2x<\frac{1}{3}$$

$$-\frac{4}{3}<-2x<-\frac{2}{3}$$

$$\frac{2}{3}>x>\frac{1}{3}$$

$$\left\{x \mid \frac{1}{3}<x<\frac{2}{3}\right\} \text{ or } \left(\frac{1}{3}, \frac{2}{3}\right)$$




73. $|2x-5|\geq 9$

$$2x-5\leq -9 \quad \text{or} \quad 2x-5\geq 9$$

$$2x\leq -4 \quad \text{or} \quad 2x\geq 14$$

$$x\leq -2 \quad \text{or} \quad x\geq 7$$

$$\{x \mid x\leq -2 \text{ or } x\geq 7\} \text{ or } (-\infty, -2] \cup [7, \infty)$$




74. $|3x+1|\geq 10$

$$3x+1\leq -10 \quad \text{or} \quad 3x+1\geq 10$$

$$3x\leq -11 \quad \text{or} \quad 3x\geq 9$$

$$x\leq -\frac{11}{3} \quad \text{or} \quad x\geq 3$$

$$\left\{x \mid x\leq -\frac{11}{3} \text{ or } x\geq 3\right\} \text{ or } \left(-\infty, -\frac{11}{3}\right] \cup [3, \infty)$$



75. $2+|2-3x|\leq 4$


$$|2-3x|\leq 2$$

$$-2\leq 2-3x\leq 2$$

$$-4\leq -3x\leq 0$$

$$\frac{4}{3}\geq x\geq 0$$

$$\left\{x \mid 0\leq x\leq \frac{4}{3}\right\} \text{ or } \left[0, \frac{4}{3}\right]$$



76. $\frac{1}{2}+\left|\frac{2x-1}{3}\right|\leq 1$

$$\left|\frac{2x-1}{3}\right|\leq \frac{1}{2}$$

$$-\frac{1}{2}\leq \frac{2x-1}{3}\leq \frac{1}{2}$$

$$-\frac{3}{2}\leq 2x-1\leq \frac{3}{2}$$

$$-\frac{1}{2}\leq 2x\leq \frac{5}{2}$$

$$-\frac{1}{4}\leq x\leq \frac{5}{4}$$

$$\left\{x \mid -\frac{1}{4}\leq x\leq \frac{5}{4}\right\} \text{ or } \left[-\frac{1}{4}, \frac{5}{4}\right]$$



Chapter 2: Linear and Quadratic Functions

77. $1 - |2 - 3x| < -4$

$$-|2 - 3x| < -5$$

$$|2 - 3x| > 5$$


$$2 - 3x < -5 \quad \text{or} \quad 2 - 3x > 5$$

$$7 < 3x \quad \text{or} \quad -3 > 3x$$

$$\frac{7}{3} < x \quad \text{or} \quad -1 > x$$

$$x < -1 \quad \text{or} \quad x > \frac{7}{3}$$

$$\left\{ x \mid x < -1 \text{ or } x > \frac{7}{3} \right\} \quad \text{or} \quad (-\infty, -1) \cup \left(\frac{7}{3}, \infty \right)$$



78. $1 - \left| \frac{2x-1}{3} \right| < -2$

$$-\left| \frac{2x-1}{3} \right| < -3$$

$$\left| \frac{2x-1}{3} \right| > 3$$


$$\frac{2x-1}{3} < -3 \quad \text{or} \quad \frac{2x-1}{3} > 3$$

$$2x-1 < -9 \quad \text{or} \quad 2x-1 > 9$$

$$2x < -8 \quad \text{or} \quad 2x > 10$$

$$x < -4 \quad \text{or} \quad x > 5$$

$$\{x \mid x < -4 \text{ or } x > 5\} \quad \text{or} \quad (-\infty, -4) \cup (5, \infty)$$



79. a. Company A: $C(x) = 0.06x + 7.00$

Company B: $C(x) = 0.08x$

b. $0.06x + 7.00 = 0.08x$

$$7.00 = 0.02x$$

$$350 = x$$

The bill from Company A will equal the bill from Company B if 350 minutes are used.

c. $0.08x < 0.06x + 7.00$

$$0.02x < 7.00$$

$$x < 350$$

The bill from Company B will be less than the bill from Company A if fewer than 350 minutes are used. That is, $0 \leq x < 350$.

80. a. $S(x) = 0.01x + 15,000$

b. $S(1,000,000) = 0.01(1,000,000) + 15,000$
 $= 10,000 + 15,000 = 25,000$

In 2005, Bill's salary was \$25,000.

c. $0.01x + 15,000 = 100,000$

$$0.01x = 85,000$$

$$x = 8,500,000$$

Bill's sales would have to be \$8,500,000 in order to earn \$100,000.

d. $0.01x + 15,000 > 150,000$

$$0.01x > 135,000$$

$$x > 13,500,000$$

Bill's sales would have to be more than \$13,500,000 in order for his salary to exceed \$150,000.

81. a. If $x = 1500 - 10p$, then $p = \frac{1500 - x}{10}$.

$$R(p) = px = p(1500 - 10p) = -10p^2 + 1500p$$

b. Domain: $\{p \mid 0 < p \leq 150\}$

c. $p = \frac{-b}{2a} = \frac{-1500}{2(-10)} = \frac{-1500}{-20} = 75$


d. The maximum revenue is

$$R(75) = -10(75)^2 + 1500(75)$$

$$= -56250 + 112500 = \$56,250$$

e. $x = 1500 - 10(75) = 1500 - 750 = 750$

f. Graph $R = -10p^2 + 1500p$ and $R = 56000$.



Find where the graphs intersect by solving

$$56000 = -10p^2 + 1500p$$

$$10p^2 - 1500p + 56000 = 0$$

$$p^2 - 150p + 5600 = 0$$

$$(p - 70)(p - 80) = 0$$

$$p = 70, p = 80$$

The company should charge between \$70 and \$80.

82. Let x represent the number of passengers over 20. Then $20 + x$ represents the total number of passengers. $15 - 0.1x$ represents the fare for each passenger. Solving the equation for total cost (\$482.40), we have:

$$(20 + x)(15 - 0.1x) = 482.40$$

$$300 + 13x - 0.1x^2 = 482.40$$

$$-0.1x^2 + 13x - 182.40 = 0$$

$$x^2 - 130x + 1824 = 0$$

$$(x - 114)(x - 16) = 0$$

$$x = 114 \text{ or } x = 16$$

Since the capacity of the bus is 44, we discard the answer 114. The total number of passengers is $20 + 16 = 36$, and the ticket price per passenger is $15 - 0.1(16) = \$13.40$. So 36 senior citizens went on the trip. Each person paid \$13.40.

83. Let $w = 4$. Then,

$$l^2 = 4(l + 4)$$

$$l^2 = 4l + 16$$

$$l^2 - 4l - 16 = 0$$

$$l = \frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(-16)}}{2(1)}$$

$$= \frac{4 \pm \sqrt{80}}{2} = \frac{4 \pm 4\sqrt{5}}{2} = 2 \pm 2\sqrt{5}$$

$$\approx 6.47 \text{ or } -2.47$$

Disregard the negative answer since length must be positive. Thus, the length of the plasterboard should be cut down to approximately 6.47 feet.

84. length of $\text{leg}_1 = x$, length of $\text{leg}_2 = 17 - x$

By the Pythagorean Theorem, we have:

$$x^2 + (17 - x)^2 = (13)^2$$

$$x^2 + x^2 - 34x + 289 = 169$$

$$2x^2 - 34x + 120 = 0$$


$$x^2 - 17x + 60 = 0$$

$$(x - 12)(x - 5) = 0$$

$$x = 12 \text{ or } x = 5$$

Thus, the legs are 5 cm and 12 cm long.

85. Let w = the width. Then $w + 2$ = the length.



By the Pythagorean Theorem we have:

$$w^2 + (w + 2)^2 = (10)^2$$

$$w^2 + w^2 + 4w + 4 = 100$$

$$2w^2 + 4w - 96 = 0$$

$$w^2 + 2w - 48 = 0$$

$$(w + 8)(w - 6) = 0$$

$$w = -8 \text{ or } w = 6$$

Disregard the negative answer because the width of a rectangle must be positive. Thus, the width is 6 inches, and the length is 8 inches

86. $C(x) = 4.9x^2 - 617.4x + 19,600$;

$$a = 4.9, b = -617.4, c = 19,600. \text{ Since}$$

$a = 4.9 > 0$, the graph opens up, so the vertex is a minimum point.

- a. The minimum marginal cost occurs at

$$x = -\frac{b}{2a} = -\frac{-617.40}{2(4.9)} = \frac{617.40}{9.8} = 63.$$

Thus, 63 golf clubs should be manufactured in order to minimize the marginal cost.

- b. The minimum marginal cost is

$$\begin{aligned} C\left(-\frac{b}{2a}\right) &= C(63) \\ &= 4.9(63)^2 - (617.40)(63) + 19600 \\ &= \$151.90 \end{aligned}$$

87. Since there are 200 feet of border, we know that $2x + 2y = 200$. The area is to be maximized, so $A = x \cdot y$. Solving the perimeter formula for y :

$$2x + 2y = 200$$

$$2y = 200 - 2x$$

$$y = 100 - x$$

The area function is:

$$A(x) = x(100 - x) = -x^2 + 100x$$

The maximum value occurs at the vertex:

$$x = -\frac{b}{2a} = -\frac{100}{2(-1)} = -\frac{100}{-2} = 50$$

The pond should be 50 feet by 50 feet for maximum area.

88. Let x represent the length and y represent the width of the rectangle.

$$2x + 2y = 20$$

$$2y = 20 - 2x$$

$$y = 10 - x$$

Chapter 2: Linear and Quadratic Functions

$$A = x \cdot y = 16$$

$$x(10 - x) = 16$$

Solving the area equation:

$$10x - x^2 = 16$$

$$x^2 - 10x + 16 = 0$$

$$(x - 8)(x - 2) = 0$$

$$x = 8 \text{ or } x = 2$$

The length and width of the rectangle are 8 feet by 2 feet.

89. The area function is:


$$A(x) = x(10 - x) = -x^2 + 10x$$

The maximum value occurs at the vertex:

$$x = -\frac{b}{2a} = -\frac{10}{2(-1)} = -\frac{10}{-2} = 5$$

The maximum area is:

$$\begin{aligned} A(5) &= -(5)^2 + 10(5) \\ &= -25 + 50 = 25 \text{ square units} \end{aligned}$$



90. Locate the origin at the point directly under the highest point of the arch. Then the equation is in the form: $y = -ax^2 + k$, where $a > 0$. Since the maximum height is 10 feet, when $x = 0$, $y = k = 10$. Since the point $(10, 0)$ is on the parabola, we can find the constant:

$$0 = -a(10)^2 + 10$$

$$a = \frac{10}{10^2} = \frac{1}{10} = 0.10$$


The equation of the parabola is:

$$y = -\frac{1}{10}x^2 + 10$$


At $x = 8$:

$$y = -\frac{1}{10}(8)^2 + 10 = -6.4 + 10 = 3.6 \text{ feet}$$

91. a.




- b. Yes, the two variables appear to have a linear relationship.
c. Using the LINear REGression program, the line of best fit is:
 $y = 1.390171918x + 1.113952697$




- d. $y = 1.390171918(26.5) + 1.113952697$
 $\approx 38.0 \text{ mm}$

92. a.



The data appear to be quadratic with $a < 0$.

- b. Using the QUADratic REGression program, the quadratic function of best fit is:
 $y = -7.76x^2 + 411.88x + 942.72$



The maximum revenue occurs at

$$A = \frac{-b}{2a} = \frac{-(411.88)}{2(-7.76)}$$

$$= \frac{-411.88}{-15.52} \approx \$26.5 \text{ thousand}$$


- c. The maximum revenue is

$$R\left(\frac{-b}{2a}\right) = R(26.53866)$$

$$= -7.76(26.5)^2 + (411.88)(26.5) + 942.72$$

$$\approx \$6408 \text{ thousand}$$


- d. 6500



Chapter 2 Test

1. $f(x) = -4x + 3$

- a. The slope f is -4 .
- b. The slope is negative, so the graph is decreasing.
- c. Plot the point $(0, 3)$. Use the slope to find an additional point by moving 1 unit to the right and 4 units down.



2.

x	y	Avg. rate of change $= \frac{\Delta y}{\Delta x}$
-2	12	
-1	7	$\frac{7-12}{-1-(-2)} = \frac{-5}{1} = -5$
0	2	$\frac{2-7}{0-(-1)} = \frac{-5}{1} = -5$

1	-3	$\frac{-3-2}{1-0} = \frac{-5}{1} = -5$
2	-8	$\frac{-8-(-3)}{2-1} = \frac{-5}{1} = -5$

Since the average rate of change is constant at -5 , this is a linear function with slope $= -5$. To find the equation of the line, we use the point-slope formula and one of the points.

$$y - y_1 = m(x - x_1)$$

$$y - 2 = -5(x - 0)$$

$$y = -5x + 2$$

3. $f(x) = 0$

$$3x^2 - 2x - 8 = 0$$

$$(3x+4)(x-2) = 0$$

$$3x+4=0 \quad \text{or} \quad x-2=0$$

$$x = -\frac{4}{3} \quad x = 2$$

The zeros of f are $-\frac{4}{3}$ and 2 .

4. $G(x) = 0$

$$-2x^2 + 4x + 1 = 0$$

$$a = -2, b = 4, c = 1$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-4 \pm \sqrt{4^2 - 4(-2)(1)}}{2(-2)}$$

$$= \frac{-4 \pm \sqrt{24}}{-4} = \frac{-4 \pm 2\sqrt{6}}{-4} = \frac{2 \pm \sqrt{6}}{2}$$

The zeros of G are $\frac{2-\sqrt{6}}{2}$ and $\frac{2+\sqrt{6}}{2}$.

5. $f(x) = g(x)$

$$x^2 + 3x = 5x + 3$$

$$x^2 - 2x - 3 = 0$$


$$(x+1)(x-3) = 0$$

$$x+1=0 \quad \text{or} \quad x-3=0$$

$$x = -1 \quad x = 3$$

The solution set is $\{-1, 3\}$.

Chapter 2: Linear and Quadratic Functions



6. $f(x) = 0$

$$(x-1)^2 + 5(x-1) + 4 = 0$$

Let $u = x-1 \rightarrow u^2 = (x-1)^2$

$$u^2 + 5u + 4 = 0$$

$$(u+4)(u+1) = 0$$

$$u+4=0 \quad \text{or} \quad u+1=0$$

$$u = -4 \quad u = -1$$


$$x-1 = -4 \quad x-1 = -1$$

$$x = -3 \quad x = 0$$

The zeros of G are -3 and 0 .

7. $f(x) = (x-3)^2 - 2$

Using the graph of $y = x^2$, shift right 3 units, then shift down 2 units.



8. a. $f(x) = 3x^2 - 12x + 4$

$a = 3, b = -12, c = 4$. Since $a = 3 > 0$, the graph opens up.

b. The x -coordinate of the vertex is

$$x = -\frac{b}{2a} = -\frac{-12}{2(3)} = \frac{12}{6} = 2.$$

The y -coordinate of the vertex is

$$f\left(-\frac{b}{2a}\right) = f(2) = 3(2)^2 - 12(2) + 4$$

$$= 12 - 24 + 4 = -8$$

Thus, the vertex is $(2, -8)$.

c. The axis of symmetry is the line $x = 2$.

d. The discriminant is:

$b^2 - 4ac = (-12)^2 - 4(3)(4) = 96 > 0$, so the graph has two x -intercepts. The x -intercepts are found by solving: $3x^2 - 12x + 4 = 0$.

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-12) \pm \sqrt{96}}{2(3)}$$


$$= \frac{12 \pm 4\sqrt{6}}{6} = \frac{6 \pm 2\sqrt{6}}{3}$$

The x -intercepts are $\frac{6-2\sqrt{6}}{3} \approx 0.37$ and

$\frac{6+2\sqrt{6}}{3} \approx 3.63$. The y -intercept is

$$f(0) = 3(0)^2 - 12(0) + 4 = 4.$$

e.



f. The domain is $(-\infty, \infty)$.

The range is $[-8, \infty)$.

g. Decreasing on $(-\infty, 2)$.

Increasing on $(2, \infty)$.

9. a. $g(x) = -2x^2 + 4x - 5$

$a = -2, b = 4, c = -5$. Since $a = -2 < 0$, the graph opens down.

b. The x -coordinate of the vertex is

$$x = -\frac{b}{2a} = -\frac{4}{2(-2)} = \frac{4}{4} = 1.$$

The y -coordinate of the vertex is

$$g\left(-\frac{b}{2a}\right) = g(1) = -2(1)^2 + 4(1) - 5$$

$$= -2 + 4 - 5 = -3$$


Thus, the vertex is $(1, -3)$.

c. The axis of symmetry is the line $x = 1$.

d. The discriminant is:

$$b^2 - 4ac = (4)^2 - 4(-2)(-5) = -24 < 0, \text{ so the graph has no } x\text{-intercepts. The } y\text{-intercept is } g(0) = -2(0)^2 + 4(0) - 5 = -5.$$

e.



f. The domain is $(-\infty, \infty)$.

The range is $(-\infty, -3]$.

g. Increasing on $(-\infty, 1)$.

Decreasing on $(1, \infty)$.

10. Consider the form $y = a(x-h)^2 + k$. From the graph we know that the vertex is $(1, -32)$ so we have $h = 1$ and $k = -32$. The graph also passes through the point $(x, y) = (0, -30)$. Substituting these values for x, y, h , and k , we can solve for a :
 $-30 = a(0-1)^2 + (-32)$ The quadratic function is
 $-30 = a(-1)^2 - 32$
 $-30 = a - 32$
 $2 = a$
 $f(x) = 2(x-1)^2 - 32 = 2x^2 - 4x - 30$.

11. $f(x) = -2x^2 + 12x + 3$
 $a = -2, b = 12, c = 3$. Since $a = -2 < 0$, the graph opens down, so the vertex is a maximum point. The maximum occurs at
 $x = -\frac{b}{2a} = -\frac{12}{2(-2)} = -\frac{12}{-4} = 3$.
The maximum value is
 $f(3) = -2(3)^2 + 12(3) + 3 = -18 + 36 + 3 = 21$.

12. $x^2 - 10x + 24 \geq 0$
 $f(x) = x^2 - 10x + 24$
 $x^2 - 10x + 24 = 0$
 $(x-4)(x-6) = 0$

$x = 4, x = 6$ are the zeros of f .

Interval	$(-\infty, 4)$	$(4, 6)$	$(6, \infty)$
Test Number	0	5	7
Value of f	24	-1	3
Conclusion	Positive	Negative	Positive

The solution set is $\{x \mid x \leq 4 \text{ or } x \geq 6\}$ or, using interval notation, $(-\infty, 4] \cup [6, \infty)$.

13. $f(x) = 0$
 $2x^2 + 4x + 5 = 0$
 $a = 2, b = 4, c = 5$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-4 \pm \sqrt{4^2 - 4(2)(5)}}{2(2)}$$


$$= \frac{-4 \pm \sqrt{-24}}{4} = \frac{-4 \pm 2\sqrt{6}i}{4} = -1 \pm \frac{\sqrt{6}}{2}i$$

The complex zeros of f are $-1 - \frac{\sqrt{6}}{2}i$ and $-1 + \frac{\sqrt{6}}{2}i$.

14. $|3x+1| = 8$
 $3x+1 = 8$ or $3x+1 = -8$
 $3x = 7$ or $3x = -9$
 $x = \frac{7}{3}$ or $x = -3$

The solution set is $\left\{-3, \frac{7}{3}\right\}$.

15. $\left|\frac{x+3}{4}\right| < 2$
 $-2 < \frac{x+3}{4} < 2$
 $-8 < x+3 < 8$
 $-11 < x < 5$
 $\{x \mid -11 < x < 5\}$ or $(-11, 5)$



Chapter 2: Linear and Quadratic Functions

16. $|2x+3|-4 \geq 3$


$$|2x+3| \geq 7$$

$$2x+3 \leq -7 \quad \text{or} \quad 2x+3 \geq 7$$

$$2x \leq -10 \quad \text{or} \quad 2x \geq 4$$

$$x \leq -5 \quad \text{or} \quad x \geq 2$$

$$\{x|x \leq -5 \text{ or } x \geq 2\} \quad \text{or} \quad (-\infty, -5] \cup [2, \infty)$$



17. a. $C(m) = 0.15m + 129.50$

b. $C(860) = 0.15(860) + 129.50$

$$= 129 + 129.50 = 258.50$$

If 860 miles are driven, the rental cost is \$258.50.

c. $C(m) = 213.80$

$$0.15m + 129.50 = 213.80$$

$$0.15m = 84.30$$

$$m = 562$$

The rental cost is \$213.80 if 562 miles were driven.

18. a. $R(x) = x\left(-\frac{1}{10}x + 1000\right) = -\frac{1}{10}x^2 + 1000x$

b. $R(400) = -\frac{1}{10}(400)^2 + 1000(400)$

$$= -16,000 + 400,000$$

$$= \$384,000$$

c. $x = \frac{-b}{2a} = \frac{-1000}{2\left(-\frac{1}{10}\right)} = \frac{-1000}{\left(-\frac{1}{5}\right)} = 5000$

The maximum revenue is

$$R(5000) = -\frac{1}{10}(5000)^2 + 1000(5000)$$

$$= -250,000 + 5,000,000$$

$$= \$2,500,000$$


Thus, 5000 units maximizes revenue at \$2,500,000.

d. $p = -\frac{1}{10}(5000) + 1000$

$$= -500 + 1000$$


$$= \$500$$

19. a. Set A:



The data appear to be linear with a negative slope.

Set B:



The data appear to be quadratic and opens up.

b. Using the LINear REGression program, the linear function of best fit is:

$$y = -4.234x - 2.362$$

```
LinReg
y=ax+b
a=-4.234042553
b=-2.361702128
r^2=.9341180356
r=-.9664978197
```

c. Using the QUADratic REGression program, the quadratic function of best fit is:

$$y = 1.993x^2 + 0.289x + 2.503$$

```
QuadReg
y=ax^2+bx+c
a=1.992842536
b=.2893660532
c=2.503067485
```

Chapter 2 Cumulative Review

1. $P = (-1, 3); Q = (4, -2)$

Distance between P and Q :

$$\begin{aligned} d(P, Q) &= \sqrt{(4 - (-1))^2 + (-2 - 3)^2} \\ &= \sqrt{(5)^2 + (-5)^2} \\ &= \sqrt{25 + 25} \\ &= \sqrt{50} = 5\sqrt{2} \end{aligned}$$

Midpoint between P and Q :

$$\left(\frac{-1 + 4}{2}, \frac{3 - 2}{2} \right) = \left(\frac{3}{2}, \frac{1}{2} \right) = (1.5, 0.5)$$

2. $y = x^3 - 3x + 1$

a. $(-2, -1): -1 = (-2)^3 - 3(-2) + 1$
 $-1 = -8 + 6 + 1$
 $-1 = -1$

Yes, $(-2, -1)$ is on the graph.

b. $(2, 3): 3 = (2)^3 - 3(2) + 1$
 $3 = 8 - 6 + 1$
 $3 = 3$

Yes, $(2, 3)$ is on the graph.

c. $(3, 1): 1 = (3)^3 - 3(3) + 1$
 $1 = -27 - 9 + 1$
 $1 \neq -35$


No, $(3, 1)$ is not on the graph.

3. $5x + 3 \geq 0$

$$5x \geq -3$$

$$x \geq -\frac{3}{5}$$

The solution set is $\left\{ x \mid x \geq -\frac{3}{5} \right\}$ or $\left[-\frac{3}{5}, +\infty \right)$.



4. $(-1, 4)$ and $(2, -2)$ are points on the line.

$$\text{Slope} = \frac{-2 - 4}{2 - (-1)} = \frac{-6}{3} = -2$$


$$y - y_1 = m(x - x_1)$$

$$y - 4 = -2(x - (-1))$$

$$y - 4 = -2(x + 1)$$

$$y - 4 = -2x - 2$$

$$y = -2x + 2$$



5. Perpendicular to $y = 2x + 1$;
 Containing $(3, 5)$


$$\text{Slope of perpendicular} = -\frac{1}{2}$$

$$y - y_1 = m(x - x_1)$$

$$y - 5 = -\frac{1}{2}(x - 3)$$

$$y - 5 = -\frac{1}{2}x + \frac{3}{2}$$

$$y = -\frac{1}{2}x + \frac{13}{2}$$



6. $x^2 + y^2 - 4x + 8y - 5 = 0$


$$x^2 - 4x + y^2 + 8y = 5$$

$$(x^2 - 4x + 4) + (y^2 + 8y + 16) = 5 + 4 + 16$$

$$(x - 2)^2 + (y + 4)^2 = 25$$

$$(x - 2)^2 + (y + 4)^2 = 5^2$$

Center: $(2, -4)$ Radius = 5



7. Yes, this is a function since each x -value is paired with exactly one y -value.

8. $f(x) = x^2 - 4x + 1$

a. $f(2) = 2^2 - 4(2) + 1 = 4 - 8 + 1 = -3$

b. $f(x) + f(2) = x^2 - 4x + 1 + (-3)$
 $= x^2 - 4x - 2$

c. $f(-x) = (-x)^2 - 4(-x) + 1 = x^2 + 4x + 1$

d. $-f(x) = -(x^2 - 4x + 1) = -x^2 + 4x - 1$

e. $f(x+2) = (x+2)^2 - 4(x+2) + 1$
 $= x^2 + 4x + 4 - 4x - 8 + 1$
 $= x^2 - 3$

f. $\frac{f(x+h) - f(x)}{h}$
 $= \frac{(x+h)^2 - 4(x+h) + 1 - (x^2 - 4x + 1)}{h}$
 $= \frac{x^2 + 2xh + h^2 - 4x - 4h + 1 - x^2 + 4x - 1}{h}$
 $= \frac{2xh + h^2 - 4h}{h}$
 $= \frac{h(2x + h - 4)}{h} = 2x + h - 4$

9. $h(z) = \frac{3z-1}{6z-7}$

The denominator cannot be zero:

$$6z - 7 \neq 0$$

$$6z \neq 7$$

$$z \neq \frac{7}{6}$$

Domain: $\left\{ z \mid z \neq \frac{7}{6} \right\}$

10. Yes, the graph represents a function since it passes the Vertical Line Test.

11. $f(x) = \frac{x}{x+4}$

a. $f(1) = \frac{1}{1+4} = \frac{1}{5} \neq \frac{1}{4}$, so $\left(1, \frac{1}{4}\right)$ is not on the graph of f .

b. $f(-2) = \frac{-2}{-2+4} = \frac{-2}{2} = -1$, so $(-2, -1)$ is a point on the graph of f .

c. Solve for x :

$$2 = \frac{x}{x+4}$$

$$2x + 8 = x$$

$$x = -8$$

So, $(-8, 2)$ is a point on the graph of f .


12. $f(x) = \frac{x^2}{2x+1}$

$$f(-x) = \frac{(-x)^2}{2(-x)+1} = \frac{x^2}{-2x+1} \neq f(x) \text{ or } -f(x)$$

Therefore, f is neither even nor odd.

13. $f(x) = x^3 - 5x + 4$ on the interval $(-4, 4)$

Use MAXIMUM and MINIMUM on the graph of $y_1 = x^3 - 5x + 4$.



Local maximum is 5.30 and occurs at $x \approx -1.29$;

Local minimum is -3.30 and occurs at $x \approx 1.29$;

f is increasing on $(-4, -1.29)$ or $(1.29, 4)$;

f is decreasing on $(-1.29, 1.29)$.

14. $f(x) = 3x + 5$; $g(x) = 2x + 1$

a. $f(x) = g(x)$

$$3x + 5 = 2x + 1$$

$$3x + 5 = 2x + 1$$

$$x = -4$$

b. $f(x) > g(x)$

$$3x + 5 > 2x + 1$$

$$3x + 5 > 2x + 1$$

$$x > -4$$

The solution set is $\{x \mid x > -4\}$ or $(-4, \infty)$.

15. a. Domain: $\{x \mid -4 \leq x \leq 4\}$ or $[-4, 4]$

Range: $\{y \mid -1 \leq y \leq 3\}$ or $[-1, 3]$

b. Intercepts: $(-1, 0)$, $(0, -1)$, $(1, 0)$

x -intercepts: $-1, 1$

y -intercept: -1


c. The graph is symmetric with respect to the y -axis.

d. When $x = 2$, the function takes on a value of 1. Therefore, $f(2) = 1$.


e. The function takes on the value 3 at $x = -4$ and $x = 4$.

f. $f(x) < 0$ means that the graph lies below the x -axis. This happens for x values between -1 and 1 . Thus, the solution set is $\{x \mid -1 < x < 1\}$ or $(-1, 1)$.


g. The graph of $y = f(x) + 2$ is the graph of $y = f(x)$ but shifted up 2 units.



h. The graph of $y = f(-x)$ is the graph of $y = f(x)$ but reflected about the y -axis.



i. The graph of $y = 2f(x)$ is the graph of $y = f(x)$ but stretched vertically by a factor of 2. That is, the coordinate of each point is multiplied by 2.



j. Since the graph is symmetric about the y -axis, the function is even.

k. The function is increasing on the open interval $(0, 4)$.

Chapter 2 Projects


Project I

Answers will vary depending on the stock selected and the time period.

Chapter 2: Linear and Quadratic Functions

Project II


a.



b. The data would be best fit by a quadratic function.

```
QuadReg
y=ax^2+bx+c
a=.0851846811
b=-14.46460932
c=1069.518992
```

$$y = 0.085x^2 - 14.46x + 1069.52$$



These results seem reasonable since the function fits the data well.

c. $s_0 = 0\text{m}$

Type	Weight kg	Velocity m/sec	Equation in the form: $s(t) = -4.9t^2 + \frac{\sqrt{2}}{2}v_0t + s_0$
MG 17	10.2	905	$s(t) = -4.9t^2 + 639.93t$ Best. (It goes the highest)
MG 131	19.7	710	$s(t) = -4.9t^2 + 502.05t$
MG 151	41.5	850	$s(t) = -4.9t^2 + 601.04t$
MG 151/20	42.3	695	$s(t) = -4.9t^2 + 491.44t$
MG/FF	35.7	575	$s(t) = -4.9t^2 + 406.59t$
MK 103	145	860	$s(t) = -4.9t^2 + 608.11t$
MK 108	58	520	$s(t) = -4.9t^2 + 367.70t$
WGr 21	111	315	$s(t) = -4.9t^2 + 222.74t$

$s_0 = 200\text{m}$

Type	Weight kg	Velocity m/sec	Equation in the form: $s(t) = -4.9t^2 + \frac{\sqrt{2}}{2}v_0t + s_0$
MG 17	10.2	905	$s(t) = -4.9t^2 + 639.93t + 200$ Best. (It goes the highest)
MG 131	19.7	710	$s(t) = -4.9t^2 + 502.05t + 200$

MG 151	41.5	850	$s(t) = -4.9t^2 + 601.04t + 200$
MG 151/20	42.3	695	$s(t) = -4.9t^2 + 491.44t + 200$
MG/FF	35.7	575	$s(t) = -4.9t^2 + 406.59t + 200$
MK 103	145	860	$s(t) = -4.9t^2 + 608.11t + 200$
MK 108	58	520	$s(t) = -4.9t^2 + 367.70t + 200$
WGr 21	111	315	$s(t) = -4.9t^2 + 222.74t + 200$

$$s_0 = 30\text{m}$$

Type	Weight kg	Velocity m/sec	Equation in the form: $s(t) = -4.9t^2 + \frac{\sqrt{2}}{2}v_0t + s_0$
MG 17	10.2	905	$s(t) = -4.9t^2 + 639.93t + 30$ Best. (It goes the highest)
MG 131	19.7	710	$s(t) = -4.9t^2 + 502.05t + 30$
MG 151	41.5	850	$s(t) = -4.9t^2 + 601.04t + 30$
MG 151/20	42.3	695	$s(t) = -4.9t^2 + 491.44t + 30$
MG/FF	35.7	575	$s(t) = -4.9t^2 + 406.59t + 30$
MK 103	145	860	$s(t) = -4.9t^2 + 608.11t + 30$
MK 108	58	520	$s(t) = -4.9t^2 + 367.70t + 30$
WGr 21	111	315	$s(t) = -4.9t^2 + 222.74t + 30$

Notice that the gun is what makes the difference, not how high it is mounted necessarily. The only way to change the true maximum height that the projectile can go is to change the angle at which it fires.

Project III

a.

x	1	2	3	4	5
$y = -2x + 5$	3	1	-1	-3	-5

b.


$$\frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - 3}{2 - 1} = -2$$

$$\frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-1 - 1}{3 - 2} = -2$$

$$\frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 - (-1)}{4 - 3} = -2$$

$$\frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-5 - (-3)}{5 - 4} = -2$$

All of the values of $\frac{\Delta y}{\Delta x}$ are the same.



d.

$$\frac{\Delta I}{\Delta x} = \frac{30633 - 9548}{10} = 2108.50$$

$$\frac{\Delta I}{\Delta x} = \frac{37088 - 30633}{10} = 645.50$$

$$\frac{\Delta I}{\Delta x} = \frac{41072 - 37088}{10} = 398.40$$

$$\frac{\Delta I}{\Delta x} = \frac{34414 - 41072}{10} = -665.80$$

$$\frac{\Delta I}{\Delta x} = \frac{19167 - 34414}{10} = -1524.70$$

Chapter 2: Linear and Quadratic Functions

These $\frac{\Delta I}{\Delta x}$ values are not all equal. The data are not linearly related.

e.

x	-2	-1	0	1	2	3	4
y	23	9	3	5	15	33	59
$\frac{\Delta y}{\Delta x}$		-14	-6	2	10	18	26

As x increases, $\frac{\Delta y}{\Delta x}$ increases. This makes sense because the parabola is increasing (going up) steeply as x increases.

f.

x	-2	-1	0	1	2	3	4
y	23	9	3	5	15	33	59
$\frac{\Delta^2 y}{\Delta x^2}$			8	8	8	8	8

The second differences are all the same.

- g. The paragraph should mention at least two observations:
1. The first differences for a linear function are all the same.
 2. The second differences for a quadratic function are the same.

Project IV

- a. – i. Answers will vary, depending on where the CBL is located above the bouncing ball.
- j. The ratio of the heights between bounces will be the same.