


# SOLUTIONS MANUAL

LIAL  
HORNSBY  
SCHNEIDER

## Trigonometry

9<sup>TH</sup> EDITION



# Chapter 2

## Acute Angles and Right Triangles

### Section 2.1: Trigonometric Functions of Acute Angles

$$1. \sin A = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{21}{29}$$

$$\cos A = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{20}{29}$$

$$\tan A = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{21}{20}$$

$$2. \sin A = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{45}{53}$$

$$\cos A = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{28}{53}$$

$$\tan A = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{45}{28}$$

$$3. \sin A = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{n}{p}$$

$$\cos A = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{m}{p}$$

$$\tan A = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{n}{m}$$

$$4. \sin A = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{k}{z}$$

$$\cos A = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{y}{z}$$

$$\tan A = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{k}{y}$$

For Exercises 5–10, refer to the Function Values of Special Angles chart on page 54 of the text.

$$5. \text{ C; } \sin 30^\circ = \frac{1}{2}$$

$$6. \text{ H; } \cos 45^\circ = \frac{\sqrt{2}}{2}$$

$$7. \text{ B; } \tan 45^\circ = 1$$

$$8. \text{ G; } \sec 60^\circ = \frac{1}{\cos 60^\circ} = \frac{1}{\frac{1}{2}} = 2$$

$$9. \text{ E; } \csc 60^\circ = \frac{1}{\sin 60^\circ} = \frac{1}{\frac{\sqrt{3}}{2}} = \frac{2}{\sqrt{3}} \\ = \frac{2}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{2\sqrt{3}}{3}$$

$$10. \text{ A; } \cot 30^\circ = \frac{\cos 30^\circ}{\sin 30^\circ} = \frac{\frac{\sqrt{3}}{2}}{\frac{1}{2}} = \frac{\sqrt{3}}{2} \cdot \frac{2}{1} = \sqrt{3}$$

$$11. a = 5, b = 12$$

$$c^2 = a^2 + b^2 \Rightarrow c^2 = 5^2 + 12^2 \Rightarrow c^2 = 169 \Rightarrow c = 13$$

$$\sin B = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{b}{c} = \frac{12}{13}$$

$$\cos B = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{a}{c} = \frac{5}{13}$$

$$\tan B = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{b}{a} = \frac{12}{5}$$

$$\cot B = \frac{\text{side adjacent}}{\text{side opposite}} = \frac{a}{b} = \frac{5}{12}$$

$$\sec B = \frac{\text{hypotenuse}}{\text{side adjacent}} = \frac{c}{a} = \frac{13}{5}$$

$$\csc B = \frac{\text{hypotenuse}}{\text{side opposite}} = \frac{c}{b} = \frac{13}{12}$$

$$12. a = 3, b = 5$$

$$c^2 = a^2 + b^2 \Rightarrow c^2 = 3^2 + 5^2 \Rightarrow c^2 = 34 \Rightarrow c = \sqrt{34}$$

$$\sin B = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{b}{c} = \frac{5}{\sqrt{34}}$$

$$= \frac{5}{\sqrt{34}} \cdot \frac{\sqrt{34}}{\sqrt{34}} = \frac{5\sqrt{34}}{34}$$

$$\cos B = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{a}{c} = \frac{3}{\sqrt{34}}$$

$$= \frac{3}{\sqrt{34}} \cdot \frac{\sqrt{34}}{\sqrt{34}} = \frac{3\sqrt{34}}{34}$$

$$\tan B = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{b}{a} = \frac{5}{3}$$

(continued on next page)

(continued from page 45)

$$\cot B = \frac{\text{side adjacent}}{\text{side opposite}} = \frac{a}{b} = \frac{3}{5}$$

$$\sec B = \frac{\text{hypotenuse}}{\text{side adjacent}} = \frac{c}{a} = \frac{\sqrt{34}}{3}$$

$$\csc B = \frac{\text{hypotenuse}}{\text{side opposite}} = \frac{c}{b} = \frac{\sqrt{34}}{5}$$

13.  $a = 6, c = 7$

$$c^2 = a^2 + b^2 \Rightarrow 7^2 = 6^2 + b^2 \Rightarrow$$

$$49 = 36 + b^2 \Rightarrow 13 = b^2 \Rightarrow \sqrt{13} = b$$

$$\sin B = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{b}{c} = \frac{\sqrt{13}}{7}$$

$$\cos B = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{a}{c} = \frac{6}{7}$$

$$\tan B = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{b}{a} = \frac{\sqrt{13}}{6}$$

$$\cot B = \frac{\text{side adjacent}}{\text{side opposite}} = \frac{a}{b} = \frac{6}{\sqrt{13}}$$

$$= \frac{6}{\sqrt{13}} \cdot \frac{\sqrt{13}}{\sqrt{13}} = \frac{6\sqrt{13}}{13}$$

$$\sec B = \frac{\text{hypotenuse}}{\text{side adjacent}} = \frac{c}{a} = \frac{7}{6}$$

$$\csc B = \frac{\text{hypotenuse}}{\text{side opposite}} = \frac{c}{b} = \frac{7}{\sqrt{13}}$$

$$= \frac{7}{\sqrt{13}} \cdot \frac{\sqrt{13}}{\sqrt{13}} = \frac{7\sqrt{13}}{13}$$

14.  $b = 7, c = 12$

$$c^2 = a^2 + b^2 \Rightarrow 12^2 = a^2 + 7^2 \Rightarrow$$

$$144 = a^2 + 49 \Rightarrow 95 = a^2 \Rightarrow \sqrt{95} = a$$

$$\sin B = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{b}{c} = \frac{7}{12}$$

$$\cos B = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{a}{c} = \frac{\sqrt{95}}{12}$$

$$\tan B = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{b}{a} = \frac{7}{\sqrt{95}}$$

$$= \frac{7}{\sqrt{95}} \cdot \frac{\sqrt{95}}{\sqrt{95}} = \frac{7\sqrt{95}}{95}$$

$$\cot B = \frac{\text{side adjacent}}{\text{side opposite}} = \frac{a}{b} = \frac{\sqrt{95}}{7}$$

$$\sec B = \frac{\text{hypotenuse}}{\text{side adjacent}} = \frac{c}{a} = \frac{12}{\sqrt{95}}$$

$$= \frac{12}{\sqrt{95}} \cdot \frac{\sqrt{95}}{\sqrt{95}} = \frac{12\sqrt{95}}{95}$$

$$\csc B = \frac{\text{hypotenuse}}{\text{side opposite}} = \frac{c}{b} = \frac{12}{7}$$

15.  $a = 3, c = 5$

$$c^2 = a^2 + b^2 \Rightarrow 5^2 = 3^2 + b^2 \Rightarrow$$

$$25 = 9 + b^2 \Rightarrow 16 = b^2 \Rightarrow 4 = b$$

$$\sin B = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{b}{c} = \frac{4}{5}$$

$$\cos B = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{a}{c} = \frac{3}{5}$$

$$\tan B = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{b}{a} = \frac{4}{3}$$

$$\cot B = \frac{\text{side adjacent}}{\text{side opposite}} = \frac{a}{b} = \frac{3}{4}$$

$$\sec B = \frac{\text{hypotenuse}}{\text{side adjacent}} = \frac{c}{a} = \frac{5}{3}$$

$$\csc B = \frac{\text{hypotenuse}}{\text{side opposite}} = \frac{c}{b} = \frac{5}{4}$$

16.  $b = 8, c = 11$

$$c^2 = a^2 + b^2 \Rightarrow 11^2 = a^2 + 8^2 \Rightarrow$$

$$121 = a^2 + 64 \Rightarrow 57 = a^2 \Rightarrow \sqrt{57} = a$$

$$\sin B = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{b}{c} = \frac{8}{11}$$

$$\cos B = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{a}{c} = \frac{\sqrt{57}}{11}$$

$$\tan B = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{b}{a} = \frac{8}{\sqrt{57}}$$

$$= \frac{8}{\sqrt{57}} \cdot \frac{\sqrt{57}}{\sqrt{57}} = \frac{8\sqrt{57}}{57}$$

$$\cot B = \frac{\text{side adjacent}}{\text{side opposite}} = \frac{a}{b} = \frac{\sqrt{57}}{8}$$

$$\sec B = \frac{\text{hypotenuse}}{\text{side adjacent}} = \frac{c}{a} = \frac{11}{\sqrt{57}}$$

$$= \frac{11}{\sqrt{57}} \cdot \frac{\sqrt{57}}{\sqrt{57}} = \frac{11\sqrt{57}}{57}$$

$$\csc B = \frac{\text{hypotenuse}}{\text{side opposite}} = \frac{c}{b} = \frac{11}{8}$$

17.  $\sin \theta = \cos(90^\circ - \theta); \cos \theta = \sin(90^\circ - \theta);$   
 $\tan \theta = \cot(90^\circ - \theta); \cot \theta = \tan(90^\circ - \theta);$   
 $\sec \theta = \csc(90^\circ - \theta); \csc \theta = \sec(90^\circ - \theta)$

18.  $\cot 73^\circ = \tan(90^\circ - 73^\circ) = \tan 17^\circ$

19.  $\sec 39^\circ = \csc(90^\circ - 39^\circ) = \csc 51^\circ$

20.  $\sin 27^\circ = \cos(90^\circ - 27^\circ) = \cos 63^\circ$

21.  $\sec(\theta + 15^\circ) = \csc[90^\circ - (\theta + 15^\circ)]$   
 $= \csc(75^\circ - \theta)$

$$22. \cos(\alpha + 20^\circ) = \sin[90^\circ - (\alpha + 20^\circ)] \\ = \sin(70^\circ - \alpha)$$

$$23. \cot(\theta - 10^\circ) = \tan[90^\circ - (\theta - 10^\circ)] \\ = \tan(100^\circ - \theta)$$

$$24. \tan 25.4^\circ = \cot(90^\circ - 25.4^\circ) = \cot 64.6^\circ$$

$$25. \sin 38.7^\circ = \cos(90^\circ - 38.7^\circ) = \cos 51.3^\circ$$

26. Using  $A = 50^\circ, 102^\circ, 248^\circ,$  and  $-26^\circ,$  we see that  $\sin(90^\circ - A)$  and  $\cos A$  yield the same values.

<pre>sin(90-50) .6427876097 cos(50) .6427876097</pre>	<pre>sin(90-102) -.2079116908 cos(102) -.2079116908</pre>
<pre>sin(90-248) -.3746065934 cos(248) -.3746065934</pre>	<pre>sin(90--26) .8987940463 cos(-26) .8987940463</pre>

For exercises 27–36, if the functions in the equations are cofunctions, then the equations are true if the sum of the angles is  $90^\circ$ .

$$27. \tan \alpha = \cot(\alpha + 10^\circ) \Rightarrow \\ \alpha + (\alpha + 10^\circ) = 90^\circ \Rightarrow 2\alpha + 10^\circ = 90^\circ \Rightarrow \\ 2\alpha = 80^\circ \Rightarrow \alpha = 40^\circ$$

$$28. \cos \theta = \sin 2\theta \Rightarrow \theta + 2\theta = 90^\circ \Rightarrow \\ 3\theta = 90^\circ \Rightarrow \theta = 30^\circ$$

$$29. \sin(2\theta + 10^\circ) = \cos(3\theta - 20^\circ) \\ (2\theta + 10^\circ) + (3\theta - 20^\circ) = 90^\circ \\ 5\theta - 10^\circ = 90^\circ \\ 5\theta = 100^\circ \Rightarrow \theta = 20^\circ$$

$$30. \sec(\beta + 10^\circ) = \csc(2\beta + 20^\circ) \\ (\beta + 10^\circ) + (2\beta + 20^\circ) = 90^\circ \\ 3\beta + 30^\circ = 90^\circ \\ 3\beta = 60^\circ \Rightarrow \beta = 20^\circ$$

$$31. \tan(3\beta + 4^\circ) = \cot(5\beta - 10^\circ) \\ (3\beta + 4^\circ) + (5\beta - 10^\circ) = 90^\circ \\ 8\beta - 6^\circ = 90^\circ \\ 8\beta = 96^\circ \Rightarrow \beta = 12^\circ$$

$$32. \cot(5\theta + 2^\circ) = \tan(2\theta + 4^\circ) \\ (5\theta + 2^\circ) + (2\theta + 4^\circ) = 90^\circ \\ 7\theta + 6^\circ = 90^\circ \\ 7\theta = 84^\circ \Rightarrow \theta = 12^\circ$$

$$33. \sin(\theta - 20^\circ) = \cos(2\theta + 5^\circ) \\ (\theta - 20^\circ) + (2\theta + 5^\circ) = 90^\circ \\ 3\theta - 15^\circ = 90^\circ \\ 3\theta = 105^\circ \Rightarrow \theta = 35^\circ$$

$$34. \cos(2\theta + 50^\circ) = \sin(2\theta - 20^\circ) \\ (2\theta + 50^\circ) + (2\theta - 20^\circ) = 90^\circ \\ 4\theta + 30^\circ = 90^\circ \\ 4\theta = 60^\circ \Rightarrow \theta = 15^\circ$$

$$35. \sec(3\beta + 10^\circ) = \csc(\beta + 8^\circ) \\ (3\beta + 10^\circ) + (\beta + 8^\circ) = 90^\circ \\ 4\beta + 18^\circ = 90^\circ \\ 4\beta = 72^\circ \Rightarrow \beta = 18^\circ$$

$$36. \csc(\beta + 40^\circ) = \sec(\beta - 20^\circ) \\ (\beta + 40^\circ) + (\beta - 20^\circ) = 90^\circ \\ 2\beta + 20^\circ = 90^\circ \\ 2\beta = 70^\circ \Rightarrow \beta = 35^\circ$$

$$37. \sin 50^\circ > \sin 40^\circ$$

In the interval from  $0^\circ$  to  $90^\circ$ , as the angle increases, so does the sine of the angle, so  $\sin 50^\circ > \sin 40^\circ$  is true.

$$38. \tan 28^\circ \leq \tan 40^\circ$$

In the interval from  $0^\circ$  to  $90^\circ$ , as the angle increases, the tangent of the angle increases, so  $\tan 40^\circ > \tan 28^\circ \Rightarrow \tan 28^\circ \leq \tan 40^\circ$  is true.

$$39. \sin 46^\circ < \cos 46^\circ$$

Using the cofunction identity,  $\cos 46^\circ = \sin(90^\circ - 46^\circ) = \sin 44^\circ$ . In the interval from  $0^\circ$  to  $90^\circ$ , as the angle increases, so does the sine of the angle, so  $\sin 46^\circ < \sin 44^\circ \Rightarrow \sin 46^\circ < \cos 44^\circ$  is false.

$$40. \cos 28^\circ < \sin 28^\circ$$

Using the cofunction identity,  $\sin 28^\circ = \cos(90^\circ - 28^\circ) = \cos 62^\circ$ . In the interval from  $0^\circ$  to  $90^\circ$ , as the angle increases, the cosine of the angle decreases, so  $\cos 28^\circ < \cos 62^\circ \Rightarrow \cos 28^\circ < \sin 28^\circ$  is false.

41.  $\tan 41^\circ < \cot 41^\circ$

Using the cofunction identity,  
 $\cot 41^\circ = \tan(90^\circ - 41^\circ) = \tan 49^\circ$ . In the interval from  $0^\circ$  to  $90^\circ$ , as the angle increases, the tangent of the angle increases, so  $\tan 41^\circ < \tan 49^\circ \Rightarrow \tan 41^\circ < \cot 41^\circ$  is true.

42.  $\cot 30^\circ < \tan 40^\circ$

Using the cofunction identity,  
 $\cot 30^\circ = \tan(90^\circ - 30^\circ) = \tan 60^\circ$ . In the interval from  $0^\circ$  to  $90^\circ$ , as the angle increases, the tangent of the angle increases, so  $\tan 60^\circ < \tan 40^\circ \Rightarrow \cot 30^\circ < \cot 40^\circ$  is false.


43.  $\sec 60^\circ > \sec 30^\circ$

In the interval from  $0^\circ$  to  $90^\circ$ , as the angle increases, the cosine of the angle decreases, so the secant of the angle increases. Thus,  $\sec 60^\circ > \sec 30^\circ$  is true.

44.  $\csc 20^\circ < \csc 30^\circ$

In the interval from  $0^\circ$  to  $90^\circ$ , as the angle increases, sine of the angle increases, so cosecant of the angle decreases. Thus  $\csc 20^\circ < \csc 30^\circ$  is false.

Use the following figures for exercises 45–60.



45.  $\tan 30^\circ = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{1}{\sqrt{3}}$   
 $= \frac{1}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{\sqrt{3}}{3}$

46.  $\cot 30^\circ = \frac{\text{side adjacent}}{\text{side opposite}} = \frac{\sqrt{3}}{1} = \sqrt{3}$

47.  $\sin 30^\circ = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{1}{2}$

48.  $\cos 30^\circ = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{\sqrt{3}}{2}$

49.  $\sec 30^\circ = \frac{\text{hypotenuse}}{\text{side adjacent}} = \frac{2}{\sqrt{3}}$   
 $= \frac{2}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{2\sqrt{3}}{3}$

50.  $\csc 30^\circ = \frac{\text{hypotenuse}}{\text{side opposite}} = \frac{2}{1} = 2$

51.  $\csc 45^\circ = \frac{\text{hypotenuse}}{\text{side opposite}} = \frac{\sqrt{2}}{1} = \sqrt{2}$

52.  $\sec 45^\circ = \frac{\text{hypotenuse}}{\text{side adjacent}} = \frac{\sqrt{2}}{1} = \sqrt{2}$

53.  $\cos 45^\circ = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{1}{\sqrt{2}}$   
 $= \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{2}$

54.  $\cot 45^\circ = \frac{\text{side adjacent}}{\text{side opposite}} = \frac{1}{1} = 1$

55.  $\tan 45^\circ = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{1}{1} = 1$

56.  $\sin 45^\circ = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{1}{\sqrt{2}}$   
 $= \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{2}$


57.  $\sin 60^\circ = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{\sqrt{3}}{2}$

58.  $\cos 60^\circ = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{1}{2}$

59.  $\tan 60^\circ = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{\sqrt{3}}{1} = \sqrt{3}$


60.  $\csc 60^\circ = \frac{\text{hypotenuse}}{\text{side opposite}} = \frac{2}{\sqrt{3}}$   
 $= \frac{2}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{2\sqrt{3}}{3}$

61.




62.  $\sin 45^\circ = \frac{y}{4} \Rightarrow y = 4 \sin 45^\circ = 4 \cdot \frac{\sqrt{2}}{2} = 2\sqrt{2}$   
and

$\cos 45^\circ = \frac{x}{4} \Rightarrow x = 4 \cos 45^\circ = 4 \cdot \frac{\sqrt{2}}{2} = 2\sqrt{2}$



63. The legs of the right triangle provide the coordinates of  $P$ ,  $(2\sqrt{2}, 2\sqrt{2})$ .



$\sin 60^\circ = \frac{y}{2} \Rightarrow y = 2 \sin 60^\circ = 2 \cdot \frac{\sqrt{3}}{2} = \sqrt{3}$

and  $\cos 60^\circ = \frac{x}{2} \Rightarrow x = 2 \cos 60^\circ = 2 \cdot \frac{1}{2} = 1$

The legs of the right triangle provide the coordinates of  $P$ .  $P$  is  $(1, \sqrt{3})$ .

65.  $Y_1$  is  $\sin x$  and  $Y_2$  is  $\tan x$ .

$\sin 0^\circ = 0$	$\tan 0^\circ = 0$
$\sin 30^\circ = .5$	$\tan 30^\circ \approx .57735$
$\sin 45^\circ \approx .70711$	$\tan 45^\circ = 1$
$\sin 60^\circ \approx .86603$	$\tan 60^\circ = 1.7321$
$\sin 90^\circ = 1$	$\tan 90^\circ$ : undefined

66.  $Y_1$  is  $\cos x$  and  $Y_2$  is  $\csc x$ .


$\cos 0^\circ = 1$	$\csc 0^\circ$ : undefined
$\cos 30^\circ \approx .86603$	$\csc 30^\circ = 2$
$\cos 45^\circ \approx .70711$	$\csc 45^\circ \approx 1.4142$
$\cos 60^\circ = .5$	$\csc 60^\circ \approx 1.1547$
$\cos 90^\circ = 0$	$\csc 90^\circ = 1$

67. Since  $\sin 60^\circ = \frac{\sqrt{3}}{2}$  and  $60^\circ$  is between  $0^\circ$  and  $90^\circ$ ,  $A = 60^\circ$ .


68. .7071067812 is a rational approximation for the exact value  $\frac{\sqrt{2}}{2}$  (an irrational value).

69. The point of intersection is  $(.70710678, .70710678)$ . This corresponds to the point  $\left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$ .


Plot1 Plot2 Plot3	WINDOW
$Y_1 = X$	Xmin=-1.8
$Y_2 = \sqrt{1-X^2}$	Xmax=1.8
$Y_3 =$	Xsc1=1
$Y_4 =$	Ymin=-1.2
$Y_5 =$	Ymax=1.2
$Y_6 =$	Ysc1=1
$Y_7 =$	Xres=1



These coordinates are the sine and cosine of  $45^\circ$ .



The line passes through  $(0, 0)$  and  $(1, \sqrt{3})$ . The slope is change in  $y$  over the change in  $x$ . Thus,  $m = \frac{\sqrt{3}}{1} = \sqrt{3}$  and the equation of the line is  $y = \sqrt{3}x$ .



The line passes through  $(0, 0)$  and  $(\sqrt{3}, 1)$ . The slope is change in  $y$  over the change in  $x$ . Thus,  $m = \frac{1}{\sqrt{3}} = \frac{1}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{\sqrt{3}}{3}$  and the equation of the line is  $y = \frac{\sqrt{3}}{3}x$ .

72. One point on the line  $y = \frac{\sqrt{3}}{3}x$ , is the origin  $(0,0)$ . Let  $(x, y)$  be any other point on this line. Then, by the definition of slope,

$$m = \frac{y-0}{x-0} = \frac{y}{x} = \frac{\sqrt{3}}{3},$$


but also, by the definition of tangent,  $\tan \theta = \frac{\sqrt{3}}{3}$ . Because  $\tan 30^\circ = \frac{\sqrt{3}}{3}$ , the line  $y = \frac{\sqrt{3}}{3}x$  makes a  $30^\circ$  angle with the positive  $x$ -axis. (See Exercise 71).

73. One point on the line  $y = \sqrt{3}x$  is the origin  $(0,0)$ . Let  $(x, y)$  be any other point on this line. Then, by the definition of slope,

$$m = \frac{y-0}{x-0} = \frac{y}{x} = \sqrt{3},$$

but also, by the definition of tangent,  $\tan \theta = \sqrt{3}$ . Because  $\tan 60^\circ = \sqrt{3}$ , the line  $y = \sqrt{3}x$  makes a  $60^\circ$  angle with the positive  $x$ -axis (See exercise 70).


74. (a) The diagonal forms two isosceles right triangles. Each angle formed by a side of the square and the diagonal measures  $45^\circ$ .



(b) By the Pythagorean theorem,


$$k^2 + k^2 = c^2 \Rightarrow 2k^2 = c^2 \Rightarrow c = \sqrt{2}k.$$

Thus, the length of the diagonal is  $\sqrt{2}k$ .




(c) In a  $45^\circ$ - $45^\circ$  right triangle, the hypotenuse has a length that is  $\sqrt{2}$  times as long as either leg.

75. (a) Each of the angles of the equilateral triangle has measure  $\frac{1}{3}(180^\circ) = 60^\circ$ .



(b) The perpendicular bisects the opposite side so the length of each side opposite each  $30^\circ$  angle is  $k$ .




(c) Let  $x$  = the length of the perpendicular. Then apply the Pythagorean theorem.

$$x^2 + k^2 = (2k)^2 \Rightarrow x^2 + k^2 = 4k^2 \Rightarrow x^2 = 3k^2 \Rightarrow x = \sqrt{3}k$$

The length of the perpendicular is  $\sqrt{3}k$ .

(d) In a  $30^\circ$ - $60^\circ$  right triangle, the hypotenuse is always 2 times as long as the shorter leg, and the longer leg has a length that is  $\sqrt{3}$  times as long as that of the shorter leg. Also, the shorter leg is opposite the  $30^\circ$  angle, and the longer leg is opposite the  $60^\circ$  angle.

76. Apply the relationships between the lengths of the sides of a  $30^\circ$ - $60^\circ$  right triangle first to the triangle on the left to find the values of  $a$  and  $b$ . In the  $30^\circ$ - $60^\circ$  right triangle, the side opposite the  $30^\circ$  angle is  $\frac{1}{2}$  the length of the hypotenuse. The longer leg is  $\sqrt{3}$  times the shorter leg.




$$a = \frac{1}{2}(24) = 12 \text{ and } b = a\sqrt{3} = 12\sqrt{3}$$

Apply the relationships between the lengths of the sides of a  $45^\circ - 45^\circ$  right triangle next to the triangle on the right to find the values of  $d$  and  $c$ . In the  $45^\circ - 45^\circ$  right triangle, the sides opposite the  $45^\circ$  angles measure the same.

The hypotenuse is  $\sqrt{2}$  times the measure of a leg.  $d = b = 12\sqrt{3}$  and

$$c = d\sqrt{2} = (12\sqrt{3})(\sqrt{2}) = 12\sqrt{6}$$

77. Apply the relationships between the lengths of the sides of a  $30^\circ - 60^\circ$  right triangle first to the triangle on the left to find the values of  $y$  and  $x$ , and then to the triangle on the right to find the values of  $z$  and  $w$ . In the  $30^\circ - 60^\circ$  right triangle, the side opposite the  $30^\circ$  angle is  $\frac{1}{2}$  the length of the hypotenuse. The longer leg is  $\sqrt{3}$  times the shorter leg.




Thus, we have

$$y = \frac{1}{2}(9) = \frac{9}{2} \text{ and } x = y\sqrt{3} = \frac{9\sqrt{3}}{2}$$

$$y = z\sqrt{3}, \text{ so } z = \frac{y}{\sqrt{3}} = \frac{\frac{9}{2}}{\sqrt{3}} = \frac{9\sqrt{3}}{6} = \frac{3\sqrt{3}}{2},$$

$$\text{and } w = 2z, \text{ so } w = 2\left(\frac{3\sqrt{3}}{2}\right) = 3\sqrt{3}$$

78. Apply the relationships between the lengths of the sides of a  $30^\circ - 60^\circ$  right triangle first to the triangle on the right to find the values of  $m$  and  $a$ . In the  $30^\circ - 60^\circ$  right triangle, the side opposite the  $60^\circ$  angle is  $\sqrt{3}$  times as long as the side opposite to the  $30^\circ$  angle. The length of the hypotenuse is 2 times as long as the shorter leg (opposite the  $30^\circ$  angle).



Thus, we have

$$7 = m\sqrt{3} \Rightarrow m = \frac{7}{\sqrt{3}} = \frac{7}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{7\sqrt{3}}{3} \text{ and}$$

$$a = 2m \Rightarrow a = 2\left(\frac{7\sqrt{3}}{3}\right) = \frac{14\sqrt{3}}{3}$$


Apply the relationships between the lengths of the sides of a  $45^\circ - 45^\circ$  right triangle next to the triangle on the left to find the values of  $n$  and  $q$ . In the  $45^\circ - 45^\circ$  right triangle, the sides opposite the  $45^\circ$  angles measure the same.

The hypotenuse is  $\sqrt{2}$  times the measure of a

leg. Thus, we have  $n = a = \frac{14\sqrt{3}}{3}$  and

$$q = n\sqrt{2} = \left(\frac{14\sqrt{3}}{3}\right)\sqrt{2} = \frac{14\sqrt{6}}{3}.$$

79. Apply the relationships between the lengths of the sides of a  $45^\circ - 45^\circ$  right triangle to the triangle on the left to find the values of  $p$  and  $r$ . In the  $45^\circ - 45^\circ$  right triangle, the sides opposite the  $45^\circ$  angles measure the same. The hypotenuse is  $\sqrt{2}$  times the measure of a leg.



Thus, we have  $p = 15$  and  $r = p\sqrt{2} = 15\sqrt{2}$


Apply the relationships between the lengths of the sides of a  $30^\circ - 60^\circ$  right triangle next to the triangle on the right to find the values of  $q$  and  $t$ . In the  $30^\circ - 60^\circ$  right triangle, the side opposite the  $60^\circ$  angle is  $\sqrt{3}$  times as long as the side opposite to the  $30^\circ$  angle. The length of the hypotenuse is 2 times as long as the shorter leg (opposite the  $30^\circ$  angle). Thus, we have  $r = q\sqrt{3} \Rightarrow$

$$q = \frac{r}{\sqrt{3}} = \frac{15\sqrt{2}}{\sqrt{3}} = \frac{15\sqrt{2}}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = 5\sqrt{6} \text{ and}$$

$$t = 2q = 2(5\sqrt{6}) = 10\sqrt{6}$$



80. Let  $h$  be the height of the equilateral triangle.  $h$  bisects the base,  $s$ , and forms two  $30^\circ$ – $60^\circ$  right triangles.



The formula for the area of a triangle is

$$A = \frac{1}{2}bh. \text{ In this triangle, } b = s. \text{ The height } h$$

of the triangle is the side opposite the  $60^\circ$  angle in either  $30^\circ$ – $60^\circ$  right triangle. The side

opposite the  $30^\circ$  angle is  $\frac{s}{2}$ . The height

is  $\sqrt{3} \cdot \frac{s}{2} = \frac{s\sqrt{3}}{2}$ . So the area of the entire

$$\text{triangle is } A = \frac{1}{2}s \left( \frac{s\sqrt{3}}{2} \right) = \frac{s^2\sqrt{3}}{4}.$$

81. Since  $A = \frac{1}{2}bh$ , we have

$$A = \frac{1}{2} \cdot s \cdot s = \frac{1}{2}s^2 \text{ or } A = \frac{s^2}{2}.$$

82. Yes, the third angle can be found by subtracting the given acute angle from  $90^\circ$ , and the remaining two sides can be found using a trigonometric function involving the known angle and side.

83. Answers will vary.

### Section 2.2: Trigonometric Functions of Non-Acute Angles


- C;  $180^\circ - 98^\circ = 82^\circ$   
( $98^\circ$  is in quadrant II)
- F;  $212^\circ - 180^\circ = 32^\circ$   
( $212^\circ$  is in quadrant III)
- A;  $-135^\circ + 360^\circ = 225^\circ$  and  
 $225^\circ - 180^\circ = 45^\circ$   
( $225^\circ$  is in quadrant III)
- B;  $-60^\circ + 360^\circ = 300^\circ$  and  
 $360^\circ - 300^\circ = 60^\circ$   
( $300^\circ$  is in quadrant IV)
- D;  $750^\circ - 2 \cdot 360^\circ = 30^\circ$   
( $30^\circ$  is in quadrant I)
- B;  $480^\circ - 360^\circ = 120^\circ$  and  $180^\circ - 120^\circ = 60^\circ$   
( $120^\circ$  is in quadrant II)
- 2 is a good choice for  $r$  because in a  $30^\circ$ – $60^\circ$  right triangle, the hypotenuse is twice the length of the shorter side (the side opposite to the  $30^\circ$  angle). By choosing 2, one avoids introducing a fraction (or decimal) when determining the length of the shorter side. Choosing any even positive integer for  $r$  would have this result; however, 2 is the most convenient value.

- 8.–9. Answers will vary.

	$\theta$	$\sin \theta$	$\cos \theta$	$\tan \theta$	$\cot \theta$	$\sec \theta$	$\csc \theta$
10.	$30^\circ$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	$\sqrt{3}$	$\frac{2\sqrt{3}}{3}$	2
11.	$45^\circ$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	1	$\sqrt{2}$	$\sqrt{2}$
12.	$60^\circ$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{\sqrt{3}}{3}$	2	$\frac{2\sqrt{3}}{3}$
13.	$120^\circ$	$\frac{\sqrt{3}}{2}$	$\cos 120^\circ$ $= -\cos 60^\circ$ $= -\frac{1}{2}$	$-\sqrt{3}$	$\cot 120^\circ$ $= -\cot 60^\circ$ $= -\frac{\sqrt{3}}{3}$	$\sec 120^\circ$ $= -\sec 60^\circ$ $= -2$	$\frac{2\sqrt{3}}{3}$
14.	$135^\circ$	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\tan 135^\circ$ $= -\tan 45^\circ$ $= -1$	$\cot 135^\circ$ $= -\cot 45^\circ$ $= -1$	$-\sqrt{2}$	$\sqrt{2}$

$\theta$	$\sin \theta$	$\cos \theta$	$\tan \theta$	$\cot \theta$	$\sec \theta$	$\csc \theta$
15. $150^\circ$	$\sin 150^\circ$ $= \sin 30^\circ$ $= \frac{1}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{3}$	$\cot 150^\circ$ $= -\cot 30^\circ$ $= -\sqrt{3}$	$\sec 150^\circ$ $= -\sec 30^\circ$ $= -\frac{2\sqrt{3}}{3}$	2
16. $210^\circ$	$-\frac{1}{2}$	$\cos 210^\circ$ $= -\cos 30^\circ$ $= -\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	$\sqrt{3}$	$\sec 210^\circ$ $= -\sec 30^\circ$ $= -\frac{2\sqrt{3}}{3}$	-2
17. $240^\circ$	$-\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$	$\tan 240^\circ$ $= \tan 60^\circ$ $= \sqrt{3}$	$\cot 240^\circ$ $= \cot 60^\circ$ $= \frac{\sqrt{3}}{3}$	-2	$-\frac{2\sqrt{3}}{3}$


18. To find the reference angle for  $300^\circ$ , sketch this angle in standard position.



The reference angle is  $360^\circ - 300^\circ = 60^\circ$ .  
Since  $300^\circ$  lies in quadrant IV, the sine, tangent, cotangent, and cosecant are negative.

$$\begin{aligned}\sin 300^\circ &= -\sin 60^\circ = -\frac{\sqrt{3}}{2} \\ \cos 300^\circ &= \cos 60^\circ = \frac{1}{2} \\ \tan 300^\circ &= -\tan 60^\circ = -\sqrt{3} \\ \cot 300^\circ &= -\cot 60^\circ = -\frac{\sqrt{3}}{3} \\ \sec 300^\circ &= \sec 60^\circ = 2 \\ \csc 300^\circ &= -\csc 60^\circ = -\frac{2\sqrt{3}}{3}\end{aligned}$$


19. To find the reference angle for  $315^\circ$ , sketch this angle in standard position.



The reference angle is  $360^\circ - 315^\circ = 45^\circ$ .  
Since  $315^\circ$  lies in quadrant IV, the sine, tangent, cotangent, and cosecant are negative.

$$\begin{aligned}\sin 315^\circ &= -\sin 45^\circ = -\frac{\sqrt{2}}{2} \\ \cos 315^\circ &= \cos 45^\circ = \frac{\sqrt{2}}{2} \\ \tan 315^\circ &= -\tan 45^\circ = -1 \\ \cot 315^\circ &= -\cot 45^\circ = -1 \\ \sec 315^\circ &= \sec 45^\circ = \sqrt{2} \\ \csc 315^\circ &= -\csc 45^\circ = -\sqrt{2}\end{aligned}$$


20. To find the reference angle for  $405^\circ$ , sketch this angle in standard position.



The reference angle for  $405^\circ$  is  $405^\circ - 360^\circ = 45^\circ$ . Because  $405^\circ$  lies in quadrant I, the values of all of its trigonometric functions will be positive, so these values will be identical to the trigonometric function values for  $45^\circ$ . (See the Function Values of Special Angles table on page 54.)

$$\begin{aligned}\sin 405^\circ &= \sin 45^\circ = \frac{\sqrt{2}}{2} \\ \cos 405^\circ &= \cos 45^\circ = \frac{\sqrt{2}}{2} \\ \tan 405^\circ &= \tan 45^\circ = 1 \\ \cot 405^\circ &= \cot 45^\circ = 1 \\ \sec 405^\circ &= \sec 45^\circ = \sqrt{2} \\ \csc 405^\circ &= \csc 45^\circ = \sqrt{2}\end{aligned}$$


21. To find the reference angle for  $-300^\circ$ , sketch this angle in standard position.



The reference angle for  $-300^\circ$  is  $-300^\circ + 360^\circ = 60^\circ$ . Because  $-300^\circ$  lies in quadrant I, the values of all of its trigonometric functions will be positive, so these values will be identical to the trigonometric function values for  $60^\circ$ . See the Function Values of Special Angles table on page 54.)

$$\begin{aligned}\sin(-300^\circ) &= \sin 60^\circ = \frac{\sqrt{3}}{2} \\ \cos(-300^\circ) &= \cos 60^\circ = \frac{1}{2} \\ \tan(-300^\circ) &= \tan 60^\circ = \sqrt{3} \\ \cot(-300^\circ) &= \cot 60^\circ = \frac{\sqrt{3}}{3} \\ \sec(-300^\circ) &= \sec 60^\circ = 2 \\ \csc(-300^\circ) &= \csc 60^\circ = \frac{2\sqrt{3}}{3}\end{aligned}$$

22. To find the reference angle for  $420^\circ$ , sketch this angle in standard position.




The reference angle for  $420^\circ$  is  $420^\circ - 360^\circ = 60^\circ$ . Because  $420^\circ$  lies in quadrant I, the values of all of its trigonometric functions will be positive, so these values will be identical to the trigonometric function values for  $60^\circ$ . See the Function Values of Special Angles table on page 54.)

$$\begin{aligned}\sin(420^\circ) &= \sin 60^\circ = \frac{\sqrt{3}}{2} \\ \cos(420^\circ) &= \cos 60^\circ = \frac{1}{2} \\ \tan(420^\circ) &= \tan 60^\circ = \sqrt{3}\end{aligned}$$

$$\begin{aligned}\cot(420^\circ) &= \cot 60^\circ = \frac{\sqrt{3}}{3} \\ \sec(420^\circ) &= \sec 60^\circ = 2 \\ \csc(420^\circ) &= \csc 60^\circ = \frac{2\sqrt{3}}{3}\end{aligned}$$


23. To find the reference angle for  $480^\circ$ , sketch this angle in standard position.



$480^\circ$  is coterminal with  $480^\circ - 360^\circ = 120^\circ$ . The reference angle is  $180^\circ - 120^\circ = 60^\circ$ . Because  $480^\circ$  lies in quadrant II, the cosine, tangent, cotangent, and secant are negative.

$$\begin{aligned}\sin(480^\circ) &= \sin 60^\circ = \frac{\sqrt{3}}{2} \\ \cos(480^\circ) &= -\cos 60^\circ = -\frac{1}{2} \\ \tan(480^\circ) &= -\tan 60^\circ = -\sqrt{3} \\ \cot(480^\circ) &= -\cot 60^\circ = -\frac{\sqrt{3}}{3} \\ \sec(480^\circ) &= -\sec 60^\circ = -2 \\ \csc(480^\circ) &= \csc 60^\circ = \frac{2\sqrt{3}}{3}\end{aligned}$$


24. To find the reference angle for  $495^\circ$ , sketch this angle in standard position.



$495^\circ$  is coterminal with  $495^\circ - 360^\circ = 135^\circ$ . The reference angle is  $180^\circ - 135^\circ = 45^\circ$ . Since  $495^\circ$  lies in quadrant II, the cosine, tangent, cotangent, and secant are negative.

$$\begin{aligned}\sin 495^\circ &= \sin 45^\circ = \frac{\sqrt{2}}{2} \\ \cos 495^\circ &= -\cos 45^\circ = -\frac{\sqrt{2}}{2} \\ \tan 495^\circ &= -\tan 45^\circ = -1 \\ \cot 495^\circ &= -\cot 45^\circ = -1 \\ \sec 495^\circ &= -\sec 45^\circ = -\sqrt{2} \\ \csc 495^\circ &= \csc 45^\circ = \sqrt{2}\end{aligned}$$


25. To find the reference angle for  $570^\circ$  sketch this angle in standard position.



$570^\circ$  is coterminal with  $570^\circ - 360^\circ = 210^\circ$ . The reference angle is  $210^\circ - 180^\circ = 30^\circ$ . Since  $570^\circ$  lies in quadrant III, the sine, cosine, secant, and cosecant are negative.

$$\begin{aligned}\sin 570^\circ &= -\sin 30^\circ = -\frac{1}{2} \\ \cos 570^\circ &= -\cos 30^\circ = -\frac{\sqrt{3}}{2} \\ \tan 570^\circ &= \tan 30^\circ = \frac{\sqrt{3}}{3} \\ \cot 570^\circ &= \cot 30^\circ = \sqrt{3} \\ \sec 570^\circ &= -\sec 30^\circ = -\frac{2\sqrt{3}}{3} \\ \csc 570^\circ &= -\csc 30^\circ = -2\end{aligned}$$

26. To find the reference angle for  $750^\circ$ , sketch this angle in standard position.



$750^\circ$  is coterminal with  $30^\circ$  because  $750^\circ - 2 \cdot 360^\circ = 750^\circ - 720^\circ = 30^\circ$ . Since  $750^\circ$  lies in quadrant I, the values of all of its trigonometric functions will be positive, so these values will be identical to the trigonometric function values for  $30^\circ$ .

$$\begin{aligned}\sin 750^\circ &= \sin 30^\circ = \frac{1}{2} \\ \cos 750^\circ &= \cos 30^\circ = \frac{\sqrt{3}}{2} \\ \tan 750^\circ &= \tan 30^\circ = \frac{\sqrt{3}}{3} \\ \cot 750^\circ &= \cot 30^\circ = \sqrt{3} \\ \sec 750^\circ &= \sec 30^\circ = \frac{2\sqrt{3}}{3} \\ \csc 750^\circ &= \csc 30^\circ = 2\end{aligned}$$

27.  $1305^\circ$  is coterminal with  $1305^\circ - 3 \cdot 360^\circ = 1305^\circ - 1080^\circ = 225^\circ$ . The reference angle is  $225^\circ - 180^\circ = 45^\circ$ . Since  $1305^\circ$  lies in quadrant III, the sine, cosine, and secant and cosecant are negative.

$$\begin{aligned}\sin 1305^\circ &= -\sin 45^\circ = -\frac{\sqrt{2}}{2} \\ \cos 1305^\circ &= -\cos 45^\circ = -\frac{\sqrt{2}}{2} \\ \tan 1305^\circ &= \tan 45^\circ = 1 \\ \cot 1305^\circ &= \cot 45^\circ = 1 \\ \sec 1305^\circ &= -\sec 45^\circ = -\sqrt{2} \\ \csc 1305^\circ &= -\csc 45^\circ = -\sqrt{2}\end{aligned}$$

28.  $1500^\circ$  is coterminal with  $1500^\circ - 4 \cdot 360^\circ = 1500^\circ - 1440^\circ = 60^\circ$ . Because  $1500^\circ$  lies in quadrant I, the values of all of its trigonometric functions will be positive, so these values will be identical to the trigonometric function values for  $60^\circ$ .

$$\begin{aligned}\sin(420^\circ) &= \sin 60^\circ = \frac{\sqrt{3}}{2} \\ \cos(420^\circ) &= \cos 60^\circ = \frac{1}{2} \\ \tan(420^\circ) &= \tan 60^\circ = \sqrt{3} \\ \cot(420^\circ) &= \cot 60^\circ = \frac{\sqrt{3}}{3} \\ \sec(420^\circ) &= \sec 60^\circ = 2 \\ \csc(420^\circ) &= \csc 60^\circ = \frac{2\sqrt{3}}{3}\end{aligned}$$

29.  $2670^\circ$  is coterminal with  $2670^\circ - 7 \cdot 360^\circ = 2670^\circ - 2520^\circ = 150^\circ$ . The reference angle is  $180^\circ - 150^\circ = 30^\circ$ . Since  $2670^\circ$  lies in quadrant II, the cosine, tangent, cotangent, and secant are negative.

$$\begin{aligned}\sin 2670^\circ &= \sin 30^\circ = \frac{1}{2} \\ \cos 2670^\circ &= -\cos 30^\circ = -\frac{\sqrt{3}}{2} \\ \tan 2670^\circ &= -\tan 30^\circ = -\frac{\sqrt{3}}{3} \\ \cot 2670^\circ &= -\cot 30^\circ = -\sqrt{3} \\ \sec 2670^\circ &= -\sec 30^\circ = -\frac{2\sqrt{3}}{3} \\ \csc 2670^\circ &= \csc 30^\circ = 2\end{aligned}$$

- 30.**  $-390^\circ$  is coterminal with  $-390^\circ + 2 \cdot 360^\circ = -390^\circ + 720^\circ = 330^\circ$ . The reference angle is  $360^\circ - 330^\circ = 30^\circ$ . Since  $-390^\circ$  lies in quadrant IV, the sine, tangent, cotangent, and cosecant are negative.

$$\sin(-390^\circ) = -\sin 30^\circ = -\frac{1}{2}$$

$$\cos(-390^\circ) = \cos 30^\circ = \frac{\sqrt{3}}{2}$$

$$\tan(-390^\circ) = -\tan 30^\circ = -\frac{\sqrt{3}}{3}$$

$$\cot(-390^\circ) = -\cot 30^\circ = -\sqrt{3}$$

$$\sec(-390^\circ) = \sec 30^\circ = \frac{2\sqrt{3}}{3}$$

$$\csc(-390^\circ) = -\csc 30^\circ = -2$$

- 31.**  $-510^\circ$  is coterminal with  $-510^\circ + 2 \cdot 360^\circ = -510^\circ + 720^\circ = 210^\circ$ . The reference angle is  $210^\circ - 180^\circ = 30^\circ$ . Since  $-510^\circ$  lies in quadrant III, the sine, cosine, and secant and cosecant are negative.

$$\sin(-510^\circ) = -\sin 30^\circ = -\frac{1}{2}$$

$$\cos(-510^\circ) = -\cos 30^\circ = -\frac{\sqrt{3}}{2}$$

$$\tan(-510^\circ) = \tan 30^\circ = \frac{\sqrt{3}}{3}$$

$$\cot(-510^\circ) = \cot 30^\circ = \sqrt{3}$$

$$\sec(-510^\circ) = -\sec 30^\circ = -\frac{2\sqrt{3}}{3}$$

$$\csc(-510^\circ) = -\csc 30^\circ = -2$$

- 32.**  $-1020^\circ$  is coterminal with  $-1020^\circ + 3 \cdot 360^\circ = -1020^\circ + 1080^\circ = 60^\circ$ . Because  $-1020^\circ$  lies in quadrant I, the values of all of its trigonometric functions will be positive, so these values will be identical to the trigonometric function values for  $60^\circ$ .

$$\sin(420^\circ) = \sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos(420^\circ) = \cos 60^\circ = \frac{1}{2}$$

$$\tan(420^\circ) = \tan 60^\circ = \sqrt{3}$$

$$\cot(420^\circ) = \cot 60^\circ = \frac{\sqrt{3}}{3}$$

$$\sec(420^\circ) = \sec 60^\circ = 2$$

$$\csc(420^\circ) = \csc 60^\circ = \frac{2\sqrt{3}}{3}$$

- 33.**  $-1290^\circ$  is coterminal with  $-1290^\circ + 4 \cdot 360^\circ = -1290^\circ + 1440^\circ = 150^\circ$ . The reference angle is  $180^\circ - 150^\circ = 30^\circ$ . Since  $-1290^\circ$  lies in quadrant II, the cosine, tangent, cotangent, and secant are negative.

$$\sin 2670^\circ = \sin 30^\circ = \frac{1}{2}$$

$$\cos 2670^\circ = -\cos 30^\circ = -\frac{\sqrt{3}}{2}$$

$$\tan 2670^\circ = -\tan 30^\circ = -\frac{\sqrt{3}}{3}$$

$$\cot 2670^\circ = -\cot 30^\circ = -\sqrt{3}$$

$$\sec 2670^\circ = -\sec 30^\circ = -\frac{2\sqrt{3}}{3}$$

$$\csc 2670^\circ = \csc 30^\circ = 2$$

- 34.**  $-855^\circ$  is coterminal with  $-855^\circ + 3 \cdot 360^\circ = -855^\circ + 1080^\circ = 225^\circ$ . The reference angle is  $225^\circ - 180^\circ = 45^\circ$ . Since  $-855^\circ$  lies in quadrant III, the sine, cosine, and secant and cosecant are negative.

$$\sin(-855^\circ) = -\sin 45^\circ = -\frac{\sqrt{2}}{2}$$

$$\cos(-855^\circ) = -\cos 45^\circ = -\frac{\sqrt{2}}{2}$$

$$\tan(-855^\circ) = \tan 45^\circ = 1$$

$$\cot(-855^\circ) = \cot 45^\circ = 1$$

$$\sec(-855^\circ) = -\sec 45^\circ = -\sqrt{2}$$

$$\csc(-855^\circ) = -\csc 45^\circ = -\sqrt{2}$$

- 35.**  $-1860^\circ$  is coterminal with  $-1860^\circ + 6 \cdot 360^\circ = -1860^\circ + 2160^\circ = 300^\circ$ . The reference angle is  $360^\circ - 300^\circ = 60^\circ$ . Since  $-1860^\circ$  lies in quadrant IV, the sine, tangent, cotangent, and cosecant are negative.

$$\sin(-1860^\circ) = -\sin 60^\circ = -\frac{\sqrt{3}}{2}$$

$$\cos(-1860^\circ) = \cos 60^\circ = \frac{1}{2}$$

$$\tan(-1860^\circ) = -\tan 60^\circ = -\sqrt{3}$$

$$\cot(-1860^\circ) = -\cot 60^\circ = -\frac{\sqrt{3}}{3}$$

$$\sec(-1860^\circ) = \sec 60^\circ = 2$$

$$\csc(-1860^\circ) = -\csc 60^\circ = -\frac{2\sqrt{3}}{3}$$

36. Since  $1305^\circ$  is coterminal with an angle of  $1305^\circ - 3 \cdot 360^\circ = 1305^\circ - 1080^\circ = 225^\circ$ , it lies in quadrant III. Its reference angle is  $225^\circ - 180^\circ = 45^\circ$ . Since the sine is negative in quadrant III, we have

$$\sin 1305^\circ = -\sin 45^\circ = -\frac{\sqrt{2}}{2}.$$

37. Since  $-510^\circ$  is coterminal with an angle of  $-510^\circ + 2 \cdot 360^\circ = -510^\circ + 720^\circ = 210^\circ$ , it lies in quadrant III. Its reference angle is  $210^\circ - 180^\circ = 30^\circ$ . Since the cosine is negative in quadrant III, we have

$$\cos(-510^\circ) = -\cos 30^\circ = -\frac{\sqrt{3}}{2}.$$

38. Since  $-1020^\circ$  is coterminal with an angle of  $-1020^\circ + 3 \cdot 360^\circ = -1020^\circ + 1080^\circ = 60^\circ$ , it lies in quadrant I. Because  $-1020^\circ$  lies in quadrant I, the values of all of its trigonometric functions will be positive, so  $\tan(-1020^\circ) = \tan 60^\circ = \sqrt{3}$ .

39. Since  $1500^\circ$  is coterminal with an angle of  $1500^\circ - 4 \cdot 360^\circ = 1500^\circ - 1440^\circ = 60^\circ$ , it lies in quadrant I. Because  $1500^\circ$  lies in quadrant I, the values of all of its trigonometric functions will be positive, so

$$\sin 1500^\circ = \sin 60^\circ = \frac{\sqrt{3}}{2}.$$

40. Since  $-495^\circ$  is coterminal with an angle of  $-495^\circ + 2 \cdot 360^\circ = -495^\circ + 720^\circ = 225^\circ$ , it lies in quadrant III. Its reference angle is  $225^\circ - 180^\circ = 45^\circ$ . Since the secant is negative in quadrant III, we have

$$\sec(-495^\circ) = -\sec 45^\circ = -\sqrt{2}.$$

41. Since  $-855^\circ$  is coterminal with  $-855^\circ + 3 \cdot 360^\circ = -855^\circ + 1080^\circ = 225^\circ$ , it lies in quadrant III. Its reference angle is  $225^\circ - 180^\circ = 45^\circ$ . Since the cosecant is negative in quadrant III, we have.

$$\csc(-855^\circ) = -\csc 45^\circ = -\sqrt{2}$$

42. Since  $2280^\circ$  is coterminal with  $2280^\circ - 6 \cdot 360^\circ = 2280^\circ - 2160^\circ = 120^\circ$ , it lies in quadrant II. Its reference angle is  $180^\circ - 120^\circ = 60^\circ$ . Since the cotangent is negative in quadrant II, we have

$$\cot 2280^\circ = -\cot 60^\circ = -\frac{\sqrt{3}}{3}.$$

43. Since  $3015^\circ$  is coterminal with  $3015^\circ - 8 \cdot 360^\circ = 3015^\circ - 2880^\circ = 135^\circ$ , it lies in quadrant II. Its reference angle is  $180^\circ - 135^\circ = 45^\circ$ . Since the tangent is negative in quadrant II, we have  $\tan 3015^\circ = -\tan 45^\circ = -1$ .

44.  $\sin 30^\circ + \sin 60^\circ \stackrel{?}{=} \sin(30^\circ + 60^\circ)$

Evaluate each side to determine whether this statement is true or false.

$$\sin 30^\circ + \sin 60^\circ = \frac{1}{2} + \frac{\sqrt{3}}{2} = \frac{1 + \sqrt{3}}{2} \text{ and}$$

$$\sin(30^\circ + 60^\circ) = \sin 90^\circ = 1$$

Since  $\frac{1 + \sqrt{3}}{2} \neq 1$ , the given statement is false.

45.  $\sin(30^\circ + 60^\circ) \stackrel{?}{=} \sin 30^\circ \cdot \cos 60^\circ + \sin 60^\circ \cdot \cos 30^\circ$

Evaluate each side to determine whether this equation is true or false.

$$\sin(30^\circ + 60^\circ) = \sin 90^\circ = 1 \text{ and}$$

$$\begin{aligned} \sin 30^\circ \cdot \cos 60^\circ + \sin 60^\circ \cdot \cos 30^\circ \\ = \frac{1}{2} \cdot \frac{1}{2} + \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{3}}{2} = \frac{1}{4} + \frac{3}{4} = 1 \end{aligned}$$

Since,  $1 = 1$ , the statement is true.

46.  $\cos 60^\circ \stackrel{?}{=} 2 \cos^2 30^\circ - 1$

Evaluate each side to determine whether this statement is true or false.

$$\cos 60^\circ = \frac{1}{2} \text{ and}$$

$$\begin{aligned} 2 \cos^2 30^\circ - 1 &= 2 \left( \frac{\sqrt{3}}{2} \right)^2 - 1 = 2 \left( \frac{3}{4} \right) - 1 \\ &= \frac{3}{2} - 1 = \frac{1}{2} \end{aligned}$$

Since  $\frac{1}{2} = \frac{1}{2}$ , the statement is true.

47.  $\cos 60^\circ \stackrel{?}{=} 2 \cos 30^\circ$

Evaluate each side to determine whether this statement is true or false.

$$\cos 60^\circ = \frac{1}{2} \text{ and } 2 \cos 30^\circ = 2 \left( \frac{\sqrt{3}}{2} \right) = \sqrt{3}$$

Since  $\frac{1}{2} \neq \sqrt{3}$ , the statement is false.

48.  $\sin 120^\circ \stackrel{?}{=} \sin 150^\circ - \sin 30^\circ$

Evaluate each side to determine whether this statement is true or false.

$$\sin 120^\circ = \frac{\sqrt{3}}{2} \text{ and}$$

$$\sin 150^\circ - \sin 30^\circ = \frac{1}{2} - \frac{1}{2} = 0$$

Since  $\frac{\sqrt{3}}{2} \neq 0$ , the statement is false.

49.  $\sin 120^\circ \stackrel{?}{=} \sin 180^\circ \cdot \cos 60^\circ - \sin 60^\circ \cdot \cos 180^\circ$

Evaluate each side to determine whether this statement is true or false.

$$\sin 120^\circ = \frac{\sqrt{3}}{2} \text{ and}$$

$$\begin{aligned} \sin 180^\circ \cdot \cos 60^\circ - \sin 60^\circ \cdot \cos 180^\circ \\ = 0\left(\frac{1}{2}\right) - \left(\frac{\sqrt{3}}{2}\right)(-1) = 0 - \left(-\frac{\sqrt{3}}{2}\right) = \frac{\sqrt{3}}{2} \end{aligned}$$

Since  $\frac{\sqrt{3}}{2} = \frac{\sqrt{3}}{2}$ , the statement is true.

50.  $\sin(2 \cdot 30^\circ) \stackrel{?}{=} 2 \sin 30^\circ \cdot \cos 30^\circ$

Evaluate each side to determine whether this statement is true or false.

$$\sin(2 \cdot 30^\circ) = \sin 60^\circ = \frac{\sqrt{3}}{2} \text{ and}$$

$$2 \sin 30^\circ \cdot \cos 30^\circ = 2\left(\frac{1}{2}\right)\left(\frac{\sqrt{3}}{2}\right) = \frac{\sqrt{3}}{2}$$

Since  $\frac{\sqrt{3}}{2} = \frac{\sqrt{3}}{2}$ , the statement is true.

51.  $\sin^2 45^\circ + \cos^2 45^\circ \stackrel{?}{=} 1$

$$\left(\frac{\sqrt{2}}{2}\right)^2 + \left(\frac{\sqrt{2}}{2}\right)^2 = \frac{2}{4} + \frac{2}{4} = 1$$

Since  $1 = 1$ , the statement is true.

52.  $\tan^2 60^\circ + 1 = \sec^2 60^\circ$

Evaluate each side to determine whether this statement is true or false.

$$\tan^2 60^\circ + 1 = (\sqrt{3})^2 + 1 = 3 + 1 = 4 \text{ and}$$

$\sec^2 60^\circ = 2^2 = 4$ . Since  $4 = 4$ , the statement is true.

53.  $\cos(30^\circ + 60^\circ) \stackrel{?}{=} \cos 30^\circ + \cos 60^\circ$

Evaluate each side to determine whether this statement is true or false.

$$\cos(30^\circ + 60^\circ) = \cos 90^\circ = 0 \text{ and}$$

$$\cos 30^\circ + \cos 60^\circ = \frac{\sqrt{3}}{2} + \frac{1}{2} = \frac{\sqrt{3}+1}{2}$$

Since  $0 \neq \frac{\sqrt{3}+1}{2}$ , the statement is false.

54.  $225^\circ$  is in quadrant III, so the reference angle is  $225^\circ - 180^\circ = 45^\circ$ .

$$\cos 45^\circ = \frac{x}{r} \Rightarrow x = r \cos 45^\circ = 10 \cdot \frac{\sqrt{2}}{2} = 5\sqrt{2}$$

and

$$\sin 45^\circ = \frac{y}{r} \Rightarrow y = r \sin 45^\circ = 10 \cdot \frac{\sqrt{2}}{2} = 5\sqrt{2}$$

Since  $225^\circ$  is in quadrant III, both the  $x$ - and  $y$ -coordinates will be negative. The coordinates of  $P$  are  $(-5\sqrt{2}, -5\sqrt{2})$ .

55.  $150^\circ$  is in quadrant II, so the reference angle is  $180^\circ - 150^\circ = 30^\circ$ .

$$\cos 30^\circ = \frac{x}{r} \Rightarrow x = r \cos 30^\circ = 6 \cdot \frac{\sqrt{3}}{2} = 3\sqrt{3}$$

$$\text{and } \sin 30^\circ = \frac{y}{r} \Rightarrow y = r \sin 30^\circ = 6 \cdot \frac{1}{2} = 3$$

Since  $150^\circ$  is in quadrant II, the  $x$ -coordinate will be negative. The coordinates of  $P$  are  $(-3\sqrt{3}, 3)$ .

56. For every angle  $\theta$ ,  $\sin^2 \theta + \cos^2 \theta = 1$ . Since  $(-.8)^2 + (.6)^2 = .64 + .36 = 1$ , there is an angle  $\theta$  for which  $\cos \theta = .6$  and  $\sin \theta = -.8$ . Since  $\cos \theta > 0$  and  $\sin \theta < 0$ ,  $\theta$  lies in quadrant IV.

57. For every angle  $\theta$ ,  $\sin^2 \theta + \cos^2 \theta = 1$ . Since  $\left(\frac{3}{4}\right)^2 + \left(\frac{2}{3}\right)^2 = \frac{9}{16} + \frac{4}{9} = \frac{145}{144} \neq 1$ , there is no angle  $\theta$  for which  $\cos \theta = \frac{2}{3}$  and  $\sin \theta = \frac{3}{4}$ .

58. If  $\theta$  is in the interval  $(90^\circ, 180^\circ)$ , then  $90^\circ < \theta < 180^\circ \Rightarrow 45^\circ < \frac{\theta}{2} < 90^\circ$ . Thus  $\frac{\theta}{2}$  lies in quadrant I, and  $\sin \frac{\theta}{2}$  is positive.

59. If  $\theta$  is in the interval  $(90^\circ, 180^\circ)$ , then  
 $90^\circ < \theta < 180^\circ \Rightarrow 45^\circ < \frac{\theta}{2} < 90^\circ$ . Thus  $\frac{\theta}{2}$   
 lies in quadrant I, and  $\cos \frac{\theta}{2}$  is positive.
60. If  $\theta$  is in the interval  $(90^\circ, 180^\circ)$ , then  
 $90^\circ < \theta < 180^\circ \Rightarrow 270^\circ < \theta + 180^\circ < 360^\circ$ .  
 Thus  $\theta + 180^\circ$  lies in quadrant IV, and  
 $\cot(\theta + 180^\circ)$  is negative.
61. If  $\theta$  is in the interval  $(90^\circ, 180^\circ)$ , then  
 $90^\circ < \theta < 180^\circ \Rightarrow 270^\circ < \theta + 180^\circ < 360^\circ$ .  
 Thus  $\theta + 180^\circ$  lies in quadrant IV, and  
 $\sec(\theta + 180^\circ)$  is positive.
62. If  $\theta$  is in the interval  $(90^\circ, 180^\circ)$ , then  
 $90^\circ < \theta < 180^\circ \Rightarrow -90^\circ > -\theta > -180^\circ \Rightarrow$   
 $-180^\circ < \theta < -90^\circ$   
 Since  $180^\circ$  is coterminal with  
 $-180^\circ + 360^\circ = 180^\circ$  and  $-90^\circ$  is coterminal  
 with  $-90^\circ + 360^\circ = 270^\circ$ ,  $-\theta$  lies in quadrant  
 III, and  $\cos(-\theta)$  is negative.
63. If  $\theta$  is in the interval  $(90^\circ, 180^\circ)$ , then  
 $90^\circ < \theta < 180^\circ \Rightarrow -90^\circ > -\theta > -180^\circ \Rightarrow$   
 $-180^\circ < \theta < -90^\circ$   
 Since  $180^\circ$  is coterminal with  
 $-180^\circ + 360^\circ = 180^\circ$  and  $-90^\circ$  is coterminal  
 with  $-90^\circ + 360^\circ = 270^\circ$ ,  $-\theta$  lies in quadrant  
 III, and  $\sin(-\theta)$  is negative.
64.  $\theta$  and  $\theta + n \cdot 360^\circ$  are coterminal angles, so  
 the sine of each of these will result in the same  
 value.
65.  $\theta$  and  $\theta + n \cdot 360^\circ$  are coterminal angles, so  
 the cosine of each of these will result in the  
 same value.
66. The reference angle for  $115^\circ$  is  
 $180^\circ - 115^\circ = 65^\circ$ . Since  $115^\circ$  is in quadrant II  
 the cosine is negative. Cos  $\theta$  decreases on the  
 interval  $(90^\circ, 180^\circ)$  from 0 to  $-1$ . Therefore,  
 cos  $115^\circ$  is closest to  $-4$ .
67. The reference angle for  $115^\circ$  is  
 $180^\circ - 115^\circ = 65^\circ$ . Since  $115^\circ$  is in quadrant II  
 the cosine is negative. Sin  $\theta$  decreases on the  
 interval  $(90^\circ, 180^\circ)$  from 1 to 0. Therefore,  
 sin  $115^\circ$  is closest to .9.
68. When  $\theta = 45^\circ$ ,  $\sin \theta = \cos \theta = \frac{\sqrt{2}}{2}$ . Sine and  
 cosine are opposites in quadrants II and IV.  
 Thus,  $180^\circ - \theta = 180^\circ - 45^\circ = 135^\circ$  in  
 quadrant II, and  $360^\circ - \theta = 360^\circ - 45^\circ = 315^\circ$   
 in quadrant IV.
69. When  $\theta = 45^\circ$ ,  $\sin \theta = \cos \theta = \frac{\sqrt{2}}{2}$ . Sine and  
 cosine are both positive in quadrant I and both  
 negative in quadrant III. Since  
 $\theta + 180^\circ = 45^\circ + 180^\circ = 225^\circ$ ,  $45^\circ$  is the  
 quadrant I angle, and  $225^\circ$  is the quadrant III  
 angle.
70.  $L = \frac{(\theta_2 - \theta_1)S^2}{200(h + S \tan \alpha)}$
- (a) Substitute  $h = 1.9$  ft,  $\alpha = .9^\circ$ ,  $\theta_1 = -3^\circ$ ,  
 $\theta_2 = 4^\circ$ , and  $S = 336$  ft:  

$$L = \frac{[4 - (-3)]336^2}{200(1.9 + 336 \tan .9^\circ)} = 550$$
 ft
- (b) Substitute  $h = 1.9$  ft,  $\alpha = 1.5^\circ$ ,  $\theta_1 = -3^\circ$ ,  
 $\theta_2 = 4^\circ$ , and  $S = 336$  ft:  

$$L = \frac{[4 - (-3)]336^2}{200(1.9 + 336 \tan 1.5^\circ)} = 369$$
 ft
- (c) Answers will vary.
71.  $\sin \theta = \frac{1}{2}$   
 Since  $\sin \theta$  is positive,  $\theta$  must lie in  
 quadrants I or II. Since one angle, namely  
 $30^\circ$ , lies in quadrant I, that angle is also the  
 reference angle,  $\theta'$ . The angle in quadrant II  
 will be  $180^\circ - \theta' = 180^\circ - 30^\circ = 150^\circ$ .
72.  $\cos \theta = \frac{\sqrt{3}}{2}$   
 Since  $\cos \theta$  is positive,  $\theta$  must lie in  
 quadrants I or IV. Since one angle, namely  
 $30^\circ$ , lies in quadrant I, that angle is also the  
 reference angle,  $\theta'$ . The angle in quadrant IV  
 will be  $360^\circ - \theta' = 360^\circ - 30^\circ = 330^\circ$ .



73.  $\tan \theta = -\sqrt{3}$

Since  $\tan \theta$  is negative,  $\theta$  must lie in quadrants II or IV. Since the absolute value of  $\tan \theta$  is  $\sqrt{3}$ , the reference angle,  $\theta'$  must be  $60^\circ$ . The quadrant II angle  $\theta$  equals  $180^\circ - \theta' = 180^\circ - 60^\circ = 120^\circ$ , and the quadrant IV angle  $\theta$  equals  $360^\circ - \theta' = 360^\circ - 60^\circ = 300^\circ$ .

74.  $\sec \theta = -\sqrt{2}$

Since  $\sec \theta$  is negative,  $\theta$  must lie in quadrants II or III. Since the absolute value of  $\sec \theta$  is  $\sqrt{2}$ , the reference angle,  $\theta'$  must be  $45^\circ$ . The quadrant II angle  $\theta$  equals  $180^\circ - \theta' = 180^\circ - 45^\circ = 135^\circ$ , and the quadrant III angle  $\theta$  equals  $180^\circ + \theta' = 180^\circ + 45^\circ = 225^\circ$ .

75.  $\cos \theta = \frac{\sqrt{2}}{2}$

Since  $\cos \theta$  is positive,  $\theta$  must lie in quadrants I or IV. Since one angle, namely  $45^\circ$ , lies in quadrant I, that angle is also the reference angle,  $\theta'$ . The angle in quadrant IV will be  $360^\circ - \theta' = 360^\circ - 45^\circ = 315^\circ$ .

76.  $\cot \theta = -\frac{\sqrt{3}}{3}$

Since  $\cot \theta$  is negative,  $\theta$  must lie in quadrants II or IV. Since the absolute value of  $\cot \theta$  is  $\frac{\sqrt{3}}{3}$ , the reference angle,  $\theta'$  must be  $60^\circ$ . The quadrant II angle  $\theta$  equals  $180^\circ - \theta' = 180^\circ - 60^\circ = 120^\circ$ , and the quadrant IV angle  $\theta$  equals  $360^\circ - \theta' = 360^\circ - 60^\circ = 300^\circ$ .

77.  $\csc \theta = -2$

Since  $\csc \theta$  is negative,  $\theta$  must lie in quadrants III or IV. The absolute value of  $\csc \theta$  is 2, so the reference angle,  $\theta'$ , is  $30^\circ$ . The angle in quadrant III will be  $180^\circ + \theta' = 180^\circ + 30^\circ = 210^\circ$ , and the quadrant IV angle is  $360^\circ - \theta' = 360^\circ - 30^\circ = 330^\circ$ .

78.  $\sin \theta = -\frac{\sqrt{3}}{2}$

Since  $\sin \theta$  is negative,  $\theta$  must lie in quadrants III or IV. The absolute value of  $\sin \theta$  is  $\frac{\sqrt{3}}{2}$ , so the reference angle,  $\theta'$ , is  $60^\circ$ . The angle in quadrant III will be  $180^\circ + \theta' = 180^\circ + 60^\circ = 240^\circ$ , and the quadrant IV angle is  $360^\circ - \theta' = 360^\circ - 60^\circ = 300^\circ$ .

79.  $\tan \theta = \frac{\sqrt{3}}{3}$

Since  $\tan \theta$  is positive,  $\theta$  must lie in quadrants I or III. Since one angle, namely  $30^\circ$ , lies in quadrant I, that angle is also the reference angle,  $\theta'$ . The angle in quadrant III will be  $180^\circ + \theta' = 180^\circ + 30^\circ = 210^\circ$ .

80.  $\cos \theta = -\frac{1}{2}$

Since  $\cos \theta$  is negative,  $\theta$  must lie in quadrants II or III. Since the absolute value of  $\cos \theta$  is  $\frac{1}{2}$ , the reference angle,  $\theta'$  must be  $60^\circ$ . The quadrant II angle  $\theta$  equals  $180^\circ - \theta' = 180^\circ - 60^\circ = 120^\circ$ , and the quadrant III angle  $\theta$  equals  $180^\circ + \theta' = 180^\circ + 60^\circ = 240^\circ$ .

81.  $\csc \theta = -\sqrt{2}$

Since  $\csc \theta$  is negative,  $\theta$  must lie in quadrants III or IV. Since the absolute value of  $\csc \theta$  is  $\sqrt{2}$ , the reference angle,  $\theta'$  must be  $45^\circ$ . The quadrant III angle  $\theta$  equals  $180^\circ + \theta' = 180^\circ + 45^\circ = 225^\circ$ , and the quadrant IV angle  $\theta$  equals  $360^\circ - \theta' = 360^\circ - 45^\circ = 315^\circ$ .

82.  $\cot \theta = -1$


Since  $\cot \theta$  is negative,  $\theta$  must lie in quadrants II or IV. Since the absolute value of  $\cot \theta$  is 1 the reference angle,  $\theta'$  must be  $45^\circ$ . The quadrant II angle  $\theta$  equals  $180^\circ - \theta' = 180^\circ - 45^\circ = 135^\circ$ , and the quadrant IV angle  $\theta$  equals  $360^\circ - \theta' = 360^\circ - 45^\circ = 315^\circ$ .

### Section 2.3: Finding Trigonometric Function Values Using a Calculator

1. The CAUTION at the beginning of this section suggests verifying that a calculator is in degree mode by finding  $\sin 90^\circ$ . If the calculator is in degree mode, the display should be 1.
2. When a scientific or graphing calculator is used to find a trigonometric function value, in most cases the result is an approximate value.
3. To find values of the cotangent, secant, and cosecant functions with a calculator, it is necessary to find the reciprocal of the reciprocal function value.
4. The reciprocal is used before the inverse function key when finding the angle, but after the function key when finding the trigonometric function value.


For Exercises 5–21, be sure your calculator is in degree mode. If your calculator accepts angles in degrees, minutes, and seconds, it is not necessary to change angles to decimal degrees.

Keystroke sequences may vary on the type and/or model of calculator being used. Screens shown will be from a TI-83 Plus calculator. To obtain the degree ( $^\circ$ ) and ( $'$ ) symbols, go to the ANGLE menu (2nd APPS).




For Exercises 5–15, the calculation for decimal degrees is indicated for calculators that do not accept degree, minutes, and seconds.

5.  $\sin 38^\circ 42' \approx .6252427$




$$38^\circ 42' = \left(38 + \frac{42}{60}\right)^\circ = 38.7^\circ$$

6.  $\cot 41^\circ 24' \approx 1.1342773$




$$42^\circ 24' = \left(42 + \frac{24}{60}\right)^\circ = 41.4^\circ$$

7.  $\sec 13^\circ 15' \approx 1.0273488$




$$13^\circ 15' = \left(13 + \frac{15}{60}\right)^\circ = 13.25^\circ$$

8.  $\csc 145^\circ 45' \approx 1.7768146$




$$145^\circ 45' = \left(145 + \frac{45}{60}\right)^\circ = 145.75^\circ$$

9.  $\cot 183^\circ 48' \approx 15.055723$




$$183^\circ 48' = \left(183 + \frac{48}{60}\right)^\circ = 183.8^\circ$$

10.  $\cos 421^\circ 30' \approx .4771588$



$$421^\circ 30' = \left(421 + \frac{30}{60}\right)^\circ = 421.5^\circ$$

11.  $\sec 312^\circ 12' \approx 1.4887142$



$$312^\circ 12' = \left(312 + \frac{12}{60}\right)^\circ = 312.2^\circ$$

12.  $\tan(-80^\circ 6') \approx -5.7297416$

```
tan(-80°6')
-5.729741647
```

$$(-80^\circ 6') = -\left(80 + \frac{6}{60}\right)^\circ = -80.1^\circ$$

13.  $\sin(-317^\circ 36') \approx .6743024$

```
sin(-317°36')
.6743023876
```

$$-317^\circ 36' = -\left(317 + \frac{36}{60}\right)^\circ = -317.6^\circ$$

14.  $\cot(-512^\circ 20') \approx 1.9074147$

```
tan(-512°20')
.5242698282
Ans-1
1.907414744
```

$$-512^\circ 20' = -\left(512 + \frac{20}{60}\right)^\circ = -512.333333^\circ$$

15.  $\cos(-15') \approx .9999905$

```
cos(-0°15')
.9999904807
```

$$-15' = -\left(\frac{15}{60}\right)^\circ = -.25^\circ$$

16.  $\frac{1}{\sec 14.8^\circ} = \cos 14.8^\circ \approx .9668234$

```
cos(14.8°)
.9668233886
```

17.  $\frac{1}{\cot 23.4^\circ} = \tan 23.4^\circ \approx .4327386$

```
tan(23.4°)
.4327386422
```

18.  $\frac{\sin 33^\circ}{\cos 33^\circ} = \tan 33^\circ \approx .6494076$

```
tan(33°)
.6494075932
```

19.  $\frac{\cos 77^\circ}{\sin 77^\circ} = \cot 77^\circ \approx .2308682$

```
tan(77°)
4.331475874
Ans-1
.2308681911
```

20.  $\cos(90^\circ - 3.69^\circ) = \sin 3.69^\circ \approx .0643581$

```
sin(3.69)
.0643581381
```

21.  $\cot(90^\circ - 4.72^\circ) = \tan 4.72^\circ \approx .0825664$

```
tan(4.72)
.0825664011
```

22.  $\frac{1}{\tan(90^\circ - 22^\circ)} = \frac{1}{\cot 22^\circ} = \tan 22^\circ \approx .4040262$

```
tan(22)
.4040262258
```

23.  $\tan \theta = 1.4739716 \Rightarrow \theta \approx 55.845496^\circ$

```
tan-1(1.4739716)
55.84549629
```

24.  $\tan \theta = 6.4358841 \Rightarrow \theta \approx 81.168073^\circ$

```
tan-1(6.4358841)
81.16807334
```

25.  $\sin \theta = .27843196 \Rightarrow \theta \approx 16.166641^\circ$

```
sin-1(.27843196)
16.16664145
```

26.  $\sec \theta = 1.1606249 \Rightarrow \theta \approx 30.502748^\circ$

```
1/1.1606249
.8616048131
cos-1(Ans)
30.50274845
```

27.  $\cot \theta = 1.2575516 \Rightarrow \theta \approx 38.491580^\circ$

```
1/1.2575516
.7951959983
tan-1(Ans)
38.49157974
```

28.  $\csc \theta = 1.3861147 \Rightarrow \theta \approx 46.173582^\circ$

```
1/1.3861147
.7214410178
sin-1(Ans)
46.17358205
```

29.  $\sec \theta = 2.7496222 \Rightarrow \theta \approx 68.673241^\circ$

```
1/2.7496222
.3636863275
cos-1(Ans)
68.6732406
```

30.  $\sin \theta = .84802194 \Rightarrow \theta \approx 57.997172^\circ$

```
sin-1(.84802194)
57.99717206
```

31.  $\cos \theta = .70058013 \Rightarrow \theta \approx 45.526434^\circ$

```
cos-1(.70058013)
45.52643354
```

32. A common mistake is to have the calculator in radian mode, when it should be in degree mode (and vice versa).

```
sin(30)
-.9880316241
```

```
Normal Sci Eng
Float 0123456789
Radian Degree
Func Par Pol Seq
Connected Dot
Sequential Simul
Real a+bi re^θi
Full Horiz G-T
```

33. If the calculator allowed an angle
- $\theta$
- where
- $0^\circ \leq \theta < 360^\circ$
- , then we would need to find an angle within this interval that is coterminal with
- $2000^\circ$
- by subtracting a multiple of
- $360^\circ$
- :
- $2000^\circ - 5 \cdot 360^\circ = 2000^\circ - 1800^\circ = 200^\circ$
- . If the calculator was more restrictive on evaluating angles (such as
- $0^\circ \leq \theta < 90^\circ$
- ), then reference angles would need to be used.

34. Find the arctan of 1.482560969.

```
tan-1(1.482560969)
56.00000001
tan(56)
1.482560969
```

$A = 56^\circ.$

35. Find the sine of
- $22^\circ$
- .

```
sin(22)
.3746065934
sin-1(.3746065934)
22
```

$A = .3746065934^\circ$

- 36.
- $\sin 35^\circ \cos 55^\circ + \cos 35^\circ \sin 55^\circ = 1$

```
sin(35)cos(55)+c
os(35)sin(55)
1
```

37.  $\cos 100^\circ \cos 80^\circ - \sin 100^\circ \sin 80^\circ = -1$

A calculator screen showing the expression  $\cos(100)\cos(80) - \sin(100)\sin(80)$  and the result  $-1$ .

38.  $\cos 75^\circ 29' \cos 14^\circ 31' - \sin 75^\circ 29' \sin 14^\circ 31' = 0$

A calculator screen showing the expression  $\cos(75^\circ 29')\cos(14^\circ 31') - \sin(75^\circ 29')\sin(14^\circ 31')$  and the result  $0$ .

39.  $\cos 28^\circ 14' \cos 61^\circ 46' - \sin 28^\circ 14' \sin 61^\circ 46' = 1$

A calculator screen showing the expression  $\cos(28^\circ 14')\cos(61^\circ 46') - \sin(28^\circ 14')\sin(61^\circ 46')$  and the result  $1$ .

40.  $\sin^2 36^\circ + \cos^2 36^\circ = 1$

A calculator screen showing the expression  $(\sin(36))^2 + (\cos(36))^2$  and the result  $1$ .

41.  $2 \sin 25^\circ 13' \cos 25^\circ 13' - \sin 50^\circ 26' = 0$

A calculator screen showing the expression  $2\sin(25^\circ 13')\cos(25^\circ 13') - \sin(50^\circ 26')$  and the result  $0$ .

42. For Auto A, calculate  $70 \cdot \cos 10^\circ \approx 68.94$ .  
Auto A's reading is approximately 68.94 mph.  
For Auto B, calculate  $70 \cdot \cos 20^\circ \approx 65.78$ .  
Auto B's reading is approximately 65.78 mph.

43. The figure for this exercise indicates a right triangle. Because we are not considering the time involved in detecting the speed of the car, we will consider the speeds as sides of the right triangle. Given angle  $\theta$ ,  $\cos \theta = \frac{r}{a}$ .  
Thus, the speed that the radar detects is  $r = a \cos \theta$ .

44.  $\cos 40^\circ \stackrel{?}{=} 2 \cos 20^\circ$   
Using a calculator gives  $\cos 40^\circ \approx .76604444$  and  $2 \cos 20^\circ \approx 1.87938524$ . Thus, the statement is false.

45.  $\sin 10^\circ + \sin 10^\circ \stackrel{?}{=} \sin 20^\circ$   
Using a calculator gives  $\sin 10^\circ \approx .34729636$  and  $\sin 20^\circ \approx .34202014$ . Thus, the statement is false.

46.  $\cos 70^\circ \stackrel{?}{=} 2 \cos^2 35^\circ - 1$   
Using a calculator gives  $\cos 70^\circ \approx .34202014$  and  $2 \cos^2 35^\circ - 1 \approx .34202014$ . Thus, the statement is true.

47.  $\sin 50^\circ \stackrel{?}{=} 2 \sin 25^\circ \cos 25^\circ$   
Using a calculator gives  $\sin 50^\circ \approx .76604444$  and  $2 \sin 25^\circ \cos 25^\circ \approx .76604444$ . Thus, the statement is true.

48.  $2 \cos 38^\circ 22' \stackrel{?}{=} \cos 76^\circ 44'$   
Using a calculator gives  $2 \cos 38^\circ 22' \approx 1.56810939$  and  $\cos 76^\circ 44' \approx .22948353$ . Thus, the statement is false.

49.  $\cos 40^\circ \stackrel{?}{=} 1 - 2 \sin^2 80^\circ$   
Using a calculator gives  $\cos 40^\circ \approx .76604444$  and  $1 - 2 \sin^2 80^\circ \approx -.93969262$ . Thus, the statement is false.

50.  $\frac{1}{2} \sin 40^\circ \stackrel{?}{=} \sin \frac{1}{2}(40^\circ)$   
Using a calculator gives  $\frac{1}{2} \sin 40^\circ \approx .32139380$  and  $\sin \frac{1}{2}(40^\circ) \approx .342002014$ . Thus, the statement is false.

51.  $\sin 39^\circ 48' + \cos 39^\circ 48' \stackrel{?}{=} 1$   
Using a calculator gives  $\sin 39^\circ 48' + \cos 39^\circ 48' \approx 1.40839322 \neq 1$ .  
Thus, the statement is false.

52.  $\cos(30^\circ + 20^\circ) \stackrel{?}{=} \cos 30^\circ + \cos 20^\circ$   
Using a calculator gives  $\cos(30^\circ + 20^\circ) \approx .64278761$  and  $\cos 30^\circ + \cos 20^\circ \approx 1.8057180$ . Thus, the statement is false.

$$53. \cos(30^\circ + 20^\circ) \stackrel{?}{=} \cos 30^\circ \cos 20^\circ - \sin 30^\circ \sin 20^\circ$$

Using a calculator gives

$$\cos(30^\circ + 20^\circ) \approx .64278761 \text{ and}$$

$$\cos 30^\circ \cos 20^\circ - \sin 30^\circ \sin 20^\circ \approx .64278761.$$

Thus, the statement is true.

$$54. \tan^2 72^\circ 25' + 1 \stackrel{?}{=} \sec^2 72^\circ 25'$$

Using a calculator gives

$$\tan^2 72^\circ 25' + 1 \approx 9.9577102 \text{ and}$$

$\sec^2 72^\circ 25' \approx 9.9577102$ . Thus, the statement is true.

$$55. 1 + \cot^2 42.5^\circ \stackrel{?}{=} \csc^2 42.5^\circ$$

Using a calculator gives

$$1 + \cot^2 42.5^\circ \approx 2.1909542 \text{ and}$$

$\csc^2 42.5^\circ \approx 2.1909542$ . Thus, the statement is true.

$$56. F = W \sin \theta$$

$$F = 2400 \sin(-2.4^\circ) \approx -100.5 \text{ lb}$$

$F$  is negative because the car is traveling downhill.

$$57. F = W \sin \theta$$

$$F = 2100 \sin 1.8^\circ \approx 65.96 \text{ lb}$$

$$58. F = W \sin \theta$$

$$-145 = W \sin(-3^\circ) \Rightarrow \frac{-145}{\sin(-3^\circ)} = W \Rightarrow$$

$$W \approx 2771 \text{ lb}$$

$$59. F = W \sin \theta$$

$$-130 = 2600 \sin \theta \Rightarrow \frac{-130}{2600} = \sin \theta \Rightarrow$$

$$-.05 = \sin \theta \Rightarrow \theta = \sin^{-1}(-.05) \approx -2.87^\circ$$

$$60. F = W \sin \theta$$

$$F = 2200 \sin 2^\circ \approx 76.77889275 \text{ lb}$$

$$F = 2000 \sin 2.2^\circ \approx 76.77561818 \text{ lb}$$

The 2200-lb car on a  $2^\circ$  uphill grade has the greater grade resistance.

$$61. F = W \sin \theta$$

$$150 = 3000 \sin \theta \Rightarrow \frac{150}{3000} = \sin \theta \Rightarrow$$

$$.05 = \sin \theta \Rightarrow \theta = \sin^{-1}(.05) \approx 2.87^\circ$$

62.

$\theta$	$\sin \theta$	$\tan \theta$	$\frac{\pi\theta}{180}$
$0^\circ$	.0000	.0000	.0000
$.5^\circ$	.0087	.0087	.0087

$\theta$	$\sin \theta$	$\tan \theta$	$\frac{\pi\theta}{180}$
$1^\circ$	.0175	.0175	.0175
$1.5^\circ$	.0262	.0262	.0262
$2^\circ$	.0349	.0349	.0349
$2.5^\circ$	.0436	.0437	.0436
$3^\circ$	.0523	.0524	.0524
$3.5^\circ$	.0610	.0612	.0611
$4^\circ$	.0698	.0699	.0698

(a) From the table, we see that if  $\theta$  is small,

$$\sin \theta = \tan \theta = \frac{\pi\theta}{180}.$$

$$(b) F = W \sin \theta = W \tan \theta = \frac{W\pi\theta}{180}$$

$$(c) \tan \theta = \frac{4}{100} = .04$$

$$F \approx W \tan \theta = 2000(.04) = 80 \text{ lb}$$

$$(d) \text{ Use } F \approx \frac{W\pi\theta}{180} \text{ from part (b). Let}$$

$$\theta = 3.75 \text{ and } W = 1800.$$

$$F \approx \frac{1800\pi(3.75)}{180} \approx 117.81 \text{ lb}$$

$$63. R = \frac{V^2}{g(f + \tan \theta)}$$

(a) Since 45 mph = 66 ft/sec,

$V = 66$ ,  $\theta = 3^\circ$ ,  $g = 32.2$ , and  $f = .14$ , we have

$$R = \frac{V^2}{g(f + \tan \theta)} = \frac{66^2}{32.2(.14 + \tan 3^\circ)} \approx 703 \text{ ft}$$

(b) Since there are 5280 ft in one mile and 3600 sec in one min, we have

$$70 \text{ mph} = 70 \text{ mph} \cdot \frac{1 \text{ hr}}{3600 \text{ sec}} \cdot \frac{5280 \text{ ft}}{1 \text{ mi}}$$

$$= 102 \frac{2}{3} \text{ ft per sec}$$

$$\approx 102.67 \text{ ft per sec}$$

Since  $V = 102.67$ ,  $\theta = 3^\circ$ ,  $g = 32.2$ , and  $f = .14$ , we have

$$R = \frac{V^2}{g(f + \tan \theta)} \approx \frac{102.67^2}{32.2(.14 + \tan 3^\circ)} \approx 1701 \text{ ft}$$

- (c) Intuitively, increasing  $\theta$  would make it easier to negotiate the curve at a higher speed much like is done at a race track. Mathematically, a larger value of  $\theta$  (acute) will lead to a larger value for  $\tan \theta$ . If  $\tan \theta$  increases, then the ratio determining  $R$  will *decrease*. Thus, the radius can be smaller and the curve sharper if  $\theta$  is increased.

$$R = \frac{V^2}{g(f + \tan \theta)} = \frac{66^2}{32.2(.14 + \tan 4^\circ)} \approx 644 \text{ ft}$$

$$R = \frac{V^2}{g(f + \tan \theta)} \approx \frac{102.67^2}{32.2(.14 + \tan 4^\circ)} \approx 1559 \text{ ft}$$

As predicted, both values are less.

64. From Exercise 113,  $R = \frac{V^2}{g(f + \tan \theta)}$ .

Solving for  $V$  we have

$$R = \frac{V^2}{g(f + \tan \theta)} \Rightarrow V^2 = Rg(f + \tan \theta) \Rightarrow$$

$$V = \sqrt{Rg(f + \tan \theta)}$$

Since  $R = 1150$ ,  $\theta = 2.1^\circ$ ,  $g = 32.2$ , and

$f = .14$ , we have

$$V = \sqrt{Rg(f + \tan \theta)} = \sqrt{1150(32.2)(.14 + \tan 2.1^\circ)} \approx 80.9 \text{ ft/sec}$$

$80.9 \text{ ft/sec} \cdot 3600 \text{ sec/hr} \cdot 1 \text{ mi}/5280 \text{ ft} \approx 55 \text{ mph}$ ,

so it should have a 55 mph speed limit

65. (a)  $\theta_1 = 46^\circ$ ,  $\theta_2 = 31^\circ$ ,  $c_1 = 3 \times 10^8$  m per sec

$$\frac{c_1}{c_2} = \frac{\sin \theta_1}{\sin \theta_2} \Rightarrow c_2 = \frac{c_1 \sin \theta_2}{\sin \theta_1} \Rightarrow$$

$$c_2 = \frac{(3 \times 10^8)(\sin 31^\circ)}{\sin 46^\circ} \approx 2 \times 10^8$$

Since  $c_1$  is only given to one significant digit,  $c_2$  can only be given to one significant digit. The speed of light in the second medium is about  $2 \times 10^8$  m per sec.

(b)  $\theta_1 = 39^\circ$ ,  $\theta_2 = 28^\circ$ ,  $c_1 = 3 \times 10^8$  m per sec

$$\frac{c_1}{c_2} = \frac{\sin \theta_1}{\sin \theta_2} \Rightarrow c_2 = \frac{c_1 \sin \theta_2}{\sin \theta_1} \Rightarrow$$

$$c_2 = \frac{(3 \times 10^8)(\sin 28^\circ)}{\sin 39^\circ} \approx 2 \times 10^8$$

Since  $c_1$  is only given to one significant digit,  $c_2$  can only be given to one significant digit. The speed of light in the second medium is about  $2 \times 10^8$  m per sec.

66. (a)  $\theta_1 = 40^\circ$ ,  $c_2 = 1.5 \times 10^8$  m per sec, and  $c_1 = 3 \times 10^8$  m per sec

$$\frac{c_1}{c_2} = \frac{\sin \theta_1}{\sin \theta_2} \Rightarrow \sin \theta_2 = \frac{c_2 \sin \theta_1}{c_1} \Rightarrow$$

$$\sin \theta_2 = \frac{(1.5 \times 10^8)(\sin 40^\circ)}{3 \times 10^8} \Rightarrow$$

$$\theta_2 = \sin^{-1} \left[ \frac{(1.5 \times 10^8)(\sin 40^\circ)}{3 \times 10^8} \right] \approx 19^\circ$$

(b)  $\theta_1 = 62^\circ$ ,  $c_2 = 2.6 \times 10^8$  m per sec and  $c_1 = 3 \times 10^8$  m per sec

$$\frac{c_1}{c_2} = \frac{\sin \theta_1}{\sin \theta_2} \Rightarrow \sin \theta_2 = \frac{c_2 \sin \theta_1}{c_1} \Rightarrow$$

$$\sin \theta_2 = \frac{(2.6 \times 10^8)(\sin 62^\circ)}{3 \times 10^8} \Rightarrow$$

$$\theta_2 = \sin^{-1} \left[ \frac{(2.6 \times 10^8)(\sin 62^\circ)}{3 \times 10^8} \right] \approx 50^\circ$$

67.  $\theta_1 = 90^\circ$ ,  $c_1 = 3 \times 10^8$  m per sec, and

$$c_2 = 2.254 \times 10^8$$

$$\frac{c_1}{c_2} = \frac{\sin \theta_1}{\sin \theta_2} \Rightarrow \sin \theta_2 = \frac{c_2 \sin \theta_1}{c_1} \Rightarrow$$

$$\sin \theta_2 = \frac{(2.254 \times 10^8)(\sin 90^\circ)}{3 \times 10^8}$$

$$= \frac{2.254 \times 10^8 (1)}{3 \times 10^8} = \frac{2.254}{3} \Rightarrow$$

$$\theta_2 = \sin^{-1} \left( \frac{2.254}{3} \right) \approx 48.7^\circ$$

68.  $\theta_1 = 90^\circ - 29.6^\circ = 60.4^\circ$ ,  $c_1 = 3 \times 10^8$  m per sec, and  $c_2 = 2.254 \times 10^8$   $c_1 = 3 \times 10^8$  m per sec

$$\frac{c_1}{c_2} = \frac{\sin \theta_1}{\sin \theta_2} \Rightarrow \sin \theta_2 = \frac{c_2 \sin \theta_1}{c_1}$$

$$\sin \theta_2 = \frac{(2.254 \times 10^8)(\sin 60.4^\circ)}{3 \times 10^8}$$

$$= \frac{2.254}{3} (\sin 60.4^\circ) \Rightarrow$$

$$\theta_2 = \sin^{-1} \left( \frac{2.254}{3} (\sin 60.4^\circ) \right) \approx 40.8^\circ$$

Light from the object is refracted at an angle of  $40.8^\circ$  from the vertical. Light from the horizon is refracted at an angle of  $48.7^\circ$  from the vertical. Therefore, the fish thinks the object lies at an angle of  $48.7^\circ - 40.8^\circ = 7.9^\circ$  above the horizon.

69. (a) Let

$$V_1 = 55 \text{ mph} = 55 \text{ mph} \cdot \frac{1 \text{ hr}}{3600 \text{ sec}} \cdot \frac{5280 \text{ ft}}{1 \text{ mi}}$$

$$= 80 \frac{2}{3} \text{ ft per sec} \approx 80.67 \text{ ft per sec,}$$

and

$$V_2 = 30 \text{ mph} = 30 \text{ mph} \cdot \frac{1 \text{ hr}}{3600 \text{ sec}} \cdot \frac{5280 \text{ ft}}{1 \text{ mi}}$$

$$= 44 \text{ ft per sec}$$

Also, let  $\theta = 3.5^\circ$ ,  $K_1 = .4$ , and  $K_2 = .02$ .

$$D = \frac{1.05(V_1^2 - V_2^2)}{64.4(K_1 + K_2 + \sin \theta)}$$

$$= \frac{1.05(80.67^2 - 44^2)}{64.4(.4 + .02 + \sin 3.5^\circ)} \approx 155 \text{ ft}$$

- (b) Let  $V_1 \approx 80.67$  ft per sec,

$$V_2 = 44 \text{ ft per sec, } \theta = -2^\circ,$$

$$K_1 = .4, \text{ and } K_2 = .02.$$

$$D = \frac{1.05(V_1^2 - V_2^2)}{64.4(K_1 + K_2 + \sin \theta)}$$

$$= \frac{1.05(80.67^2 - 44^2)}{64.4[.4 + .02 + \sin(-2^\circ)]} \approx 194 \text{ ft}$$

70. Using the values for  $K_1$  and  $K_2$  from Exercise 106, determine  $V_2$  when  $D = 200$ ,  $\theta = -3.5^\circ$ ,

$$V_1 = 90 \text{ mph} = 90 \text{ mph} \cdot \frac{1 \text{ hr}}{3600 \text{ sec}} \cdot \frac{5280 \text{ ft}}{1 \text{ mi}}$$

$$= 132 \text{ ft per sec}$$

$$D = \frac{1.05(V_1^2 - V_2^2)}{64.4(K_1 + K_2 + \sin \theta)}$$

$$200 = \frac{1.05(132^2 - V_2^2)}{64.4[.4 + .02 + \sin(-3.5^\circ)]}$$

$$200 = \frac{1.05(132^2) - 1.05V_2^2}{23.12}$$

$$200(23.12) = 18,295.2 - 1.05V_2^2$$

$$4624 = 18,295.2 - 1.05V_2^2$$

$$-13,671.2 = -1.05V_2^2$$

$$V_2^2 = \frac{-13,671.2}{-1.05}$$

$$V_2^2 = 13020.19048$$

$$V_2 \approx 114.106$$

$$V_2 \approx 114 \text{ ft/sec} \cdot 3600 \text{ sec/hr} \cdot 1 \text{ mi}/5280 \text{ ft}$$

$$\approx 78 \text{ mph}$$

## Chapter 2 Quiz (Sections 2.1–2.3)

1.  $\sin A = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{24}{40} = \frac{3}{5}$

$$\cos A = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{32}{40} = \frac{4}{5}$$

$$\tan A = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{24}{32} = \frac{3}{4}$$

2.

$\theta$	$\sin \theta$	$\cos \theta$	$\tan \theta$	$\cot \theta$	$\sec \theta$	$\csc \theta$
$30^\circ$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	$\sqrt{3}$	$\frac{2\sqrt{3}}{3}$	2
$45^\circ$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	1	$\sqrt{2}$	$\sqrt{2}$
$60^\circ$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{\sqrt{3}}{3}$	2	$\frac{2\sqrt{3}}{3}$

3.  $\sin 30^\circ = \frac{w}{36} \Rightarrow w = 36 \sin 30^\circ = 36 \cdot \frac{1}{2} = 18$

$$\cos 30^\circ = \frac{x}{36} \Rightarrow x = 36 \cos 30^\circ = 36 \cdot \frac{\sqrt{3}}{2} = 18\sqrt{3}$$


$$\tan 45^\circ = \frac{w}{y} \Rightarrow 1 = \frac{18}{y} \Rightarrow y = 18$$

$$\sin 45^\circ = \frac{w}{z} \Rightarrow \frac{\sqrt{2}}{2} = \frac{18}{z} \Rightarrow z = \frac{36}{\sqrt{2}} = 18\sqrt{2}$$



4. The height of one of the six equilateral triangles from the solar cell is

$$\sin \theta = \frac{h}{x} \Rightarrow h = x \sin \theta.$$



Thus, the area of each of the triangles is

$$A = \frac{1}{2}bh = \frac{1}{2}x^2 \sin \theta. \text{ So, the area of the solar cell is } A = 6 \cdot \frac{1}{2}x^2 \sin \theta = 3x^2 \sin \theta.$$

5.  $180^\circ - 135^\circ = 45^\circ$ , so the reference angle is  $45^\circ$ . The original angle ( $135^\circ$ ) lies in quadrant II, so the sine and cosecant are positive, while the remaining trigonometric functions are negative.

$$\sin 135^\circ = \frac{\sqrt{2}}{2}; \quad \cos 135^\circ = -\frac{\sqrt{2}}{2}$$

$$\tan 135^\circ = -1; \quad \cot 135^\circ = -1$$

$$\sec 135^\circ = -\sqrt{2}; \quad \csc 135^\circ = \sqrt{2}$$

6.  $-150^\circ$  is coterminal with  $360^\circ - 150^\circ = 210^\circ$ . Since this lies in quadrant III, the reference angle is  $210^\circ - 180^\circ = 30^\circ$ . In quadrant III, the tangent and cotangent functions are positive, while the remaining trigonometric functions are negative.

$$\sin(-150^\circ) = -\sin 30^\circ = -\frac{1}{2}$$

$$\cos(-150^\circ) = -\cos 30^\circ = -\frac{\sqrt{3}}{2}$$

$$\tan(-150^\circ) = \tan 30^\circ = \frac{\sqrt{3}}{3}$$

$$\cot(-150^\circ) = \cot 30^\circ = \sqrt{3}$$

$$\sec(-150^\circ) = -\sec 30^\circ = -\frac{2\sqrt{3}}{3}$$

$$\csc(-150^\circ) = -\csc 30^\circ = -2$$

7.  $1020^\circ$  is coterminal with  $1020^\circ - 720^\circ = 300^\circ$ . Since this lies in quadrant IV, the reference angle is  $360^\circ - 300^\circ = 60^\circ$ . In quadrant IV, the cosine and secant are positive, while the remaining trigonometric functions are negative.

$$\sin 1020^\circ = -\sin 60^\circ = -\frac{\sqrt{3}}{2}$$

$$\cos 1020^\circ = \cos 60^\circ = \frac{1}{2}$$

$$\tan 1020^\circ = -\tan 60^\circ = -\sqrt{3}$$

$$\cot 1020^\circ = -\cot 60^\circ = -\frac{\sqrt{3}}{3}$$

$$\sec 1020^\circ = \sec 60^\circ = 2$$

$$\csc 1020^\circ = -\csc 60^\circ = -\frac{2\sqrt{3}}{3}$$

8.  $\sin \theta = \frac{\sqrt{3}}{2}$

Since  $\sin \theta$  is positive,  $\theta$  must lie in quadrants I or II, and the reference angle,  $\theta'$ , is  $60^\circ$ . The angle in quadrant I is  $60^\circ$ , while the angle in quadrant II is  $180^\circ - \theta' = 180^\circ - 60^\circ = 120^\circ$ .

9.  $\sec \theta = -\sqrt{2}$

Since  $\sec \theta$  is negative,  $\theta$  must lie in quadrants II or III. Since the absolute value of  $\sec \theta$  is  $\sqrt{2}$ , the reference angle,  $\theta'$  must be  $45^\circ$ . The quadrant II angle  $\theta$  equals  $180^\circ - \theta' = 180^\circ - 45^\circ = 135^\circ$ , and the quadrant III angle  $\theta$  equals  $180^\circ + \theta' = 180^\circ + 45^\circ = 225^\circ$ .

10.  $\sin 42^\circ 18' \approx .67301251$

```
sin(42°18')
.6730125135
```

11.  $\sec(-212^\circ 12') \approx -1.18176327$

```
cos(-212°12')
-.8461931661
Ans^-1
-1.181763266
```

12.  $\tan \theta = 2.6743210 \Rightarrow \theta \approx 69.497888^\circ$

```
tan^-1(2.6743210)
69.49788821
```

13.  $\csc \theta = 2.3861147 \Rightarrow \theta \approx 24.777233^\circ$

```
1/2.3861147
.4190913371
sin^-1(Ans)
24.77723309
```

14. The statement is false.


$$\sin(60^\circ + 30^\circ) = \sin 90^\circ = 1, \text{ while}$$

$$\sin 60^\circ + \sin 30^\circ = \frac{\sqrt{3}}{2} + \frac{1}{2} = \frac{\sqrt{3} + 1}{2}.$$

15. The statement is true. Using the cofunction identity,  $\tan(90^\circ - 35^\circ) = \cot 35^\circ$ .

### Section 2.4: Solving Right Triangles

- 22,894.5 to 22,895.5
- 28,999.5 to 29,000.5
- 8958.5 to 8959.5
- Answers will vary.  
No; the number of points scored will be a whole number.
- Answers will vary.  
It would be cumbersome to write 2 as 2.00 or 2.000, for example, if the measurements had 3 or 4 significant digits (depending on the problem). In the formula, it is understood that 2 is an exact value. Since the radius measurement, 54.98 cm, has four significant digits, an appropriate answer would be 345.4 cm.
- 23.0 ft indicates 3 significant digits and 23.00 ft indicates four significant digits.
- If  $h$  is the actual height of a building and the height is measured as 58.6 ft, then  $|h - 58.6| \leq .05$ .
- If  $w$  is the actual weight of a car and the weight is measured as 1542 lb, then  $|w - 1542| \leq .5$ .
- $A = 36^\circ 20'$ ,  $c = 964$  m



$$A + B = 90^\circ \Rightarrow B = 90^\circ - A \Rightarrow$$

$$B = 90^\circ - 36^\circ 20' = 89^\circ 60' - 36^\circ 20' = 53^\circ 40'$$


$$\sin A = \frac{a}{c} \Rightarrow \sin 36^\circ 20' = \frac{a}{964} \Rightarrow$$

$$a = 964 \sin 36^\circ 20' \approx 571 \text{ m (rounded to three significant digits)}$$

$$\cos A = \frac{b}{c} \Rightarrow \cos 36^\circ 20' = \frac{b}{964} \Rightarrow$$

$$b = 964 \cos 36^\circ 20' \approx 777 \text{ m (rounded to three significant digits)}$$

10.  $A = 31^\circ 40'$ ,  $a = 35.9$  km



$$A + B = 90^\circ \Rightarrow B = 90^\circ - A \Rightarrow$$

$$B = 90^\circ - 31^\circ 40' = 89^\circ 60' - 31^\circ 40' = 58^\circ 20'$$


$$\sin A = \frac{a}{c} \Rightarrow \sin 31^\circ 40' = \frac{35.9}{c} \Rightarrow$$

$$c = \frac{35.9}{\sin 31^\circ 40'} \approx 68.4 \text{ km (rounded to three significant digits)}$$

$$\tan A = \frac{a}{b} \Rightarrow \tan 31^\circ 40' = \frac{35.9}{b} \Rightarrow$$

$$b = \frac{35.9}{\tan 31^\circ 40'} \approx 58.2 \text{ km (rounded to three significant digits)}$$

11.  $N = 51.2^\circ$ ,  $m = 124$  m



$$M + N = 90^\circ \Rightarrow M = 90^\circ - N \Rightarrow$$

$$M = 90^\circ - 51.2^\circ = 38.8^\circ$$


$$\tan N = \frac{n}{m} \Rightarrow \tan 51.2^\circ = \frac{n}{124} \Rightarrow$$

$$n = 124 \tan 51.2^\circ \approx 154 \text{ m (rounded to three significant digits)}$$

$$\cos N = \frac{m}{p} \Rightarrow \cos 51.2^\circ = \frac{124}{p} \Rightarrow$$

$$p = \frac{124}{\cos 51.2^\circ} \approx 198 \text{ m (rounded to three significant digits)}$$

12.  $X = 47.8^\circ, z = 89.6$  cm




$$Y + X = 90^\circ \Rightarrow Y = 90^\circ - X \Rightarrow Y = 90^\circ - 47.8^\circ = 42.2^\circ$$

$$\sin X = \frac{x}{z} \Rightarrow \sin 47.8^\circ = \frac{x}{89.6} \Rightarrow x = 89.6 \sin 47.8^\circ \approx 66.4 \text{ cm (rounded to three significant digits)}$$

$$\cos X = \frac{y}{z} \Rightarrow \cos 47.8^\circ = \frac{y}{89.6} \Rightarrow y = 89.6 \cos 47.8^\circ \approx 60.2 \text{ cm (rounded to three significant digits)}$$

13.  $B = 42.0892^\circ, b = 56.851$




$$A + B = 90^\circ \Rightarrow A = 90^\circ - B \Rightarrow A = 90^\circ - 42.0892^\circ = 47.9108^\circ$$

$$\sin B = \frac{b}{c} \Rightarrow \sin 42.0892^\circ = \frac{56.851}{c} \Rightarrow c = \frac{56.851}{\sin 42.0892^\circ} \approx 84.816 \text{ cm (rounded to five significant digits)}$$

$$\tan B = \frac{b}{a} \Rightarrow \tan 42.0892^\circ = \frac{56.851}{a} \Rightarrow a = \frac{56.851}{\tan 42.0892^\circ} \approx 62.942 \text{ cm (rounded to five significant digits)}$$

14.  $B = 68.5142^\circ, c = 3579.42$




$$A + B = 90^\circ \Rightarrow A = 90^\circ - B \Rightarrow A = 90^\circ - 68.5142^\circ = 21.4858^\circ$$

$$\sin B = \frac{b}{c} \Rightarrow \sin 68.5142^\circ = \frac{b}{3579.42} \Rightarrow b = 3579.42 \sin 68.5142^\circ \approx 3330.68 \text{ m (rounded to six significant digits)}$$

$$\cos B = \frac{a}{c} \Rightarrow \cos 68.5142^\circ = \frac{a}{3579.42} \Rightarrow a = 3579.42 \cos 68.5142^\circ \approx 1311.04 \text{ m (rounded to six significant digits)}$$

15.  $a = 12.5, b = 16.2$



Using the Pythagorean theorem, we have  $a^2 + b^2 = c^2 \Rightarrow 12.5^2 + 16.2^2 = c^2 \Rightarrow 418.69 = c^2 \Rightarrow c \approx 20.5$  ft (rounded to three significant digits)


$$\tan A = \frac{a}{b} \Rightarrow \tan A = \frac{12.5}{16.2} \Rightarrow A = \tan^{-1} \frac{12.5}{16.2} \approx 37.6540^\circ \approx 37^\circ + (.6540 \cdot 60)' \approx 37^\circ 39' \approx 37^\circ 40'$$

(rounded to three significant digits)

$$\tan B = \frac{b}{a} \Rightarrow \tan B = \frac{16.2}{12.5} \Rightarrow B = \tan^{-1} \frac{16.2}{12.5} \approx 52.3460^\circ \approx 52^\circ + (.3460 \cdot 60)' \approx 52^\circ 21' \approx 52^\circ 20'$$

(rounded to three significant digits)

16.  $a = 4.80, c = 15.3$



Using the Pythagorean theorem, we have  $a^2 + b^2 = c^2 \Rightarrow 4.80^2 + b^2 = 15.3^2 \Rightarrow 4.80^2 + b^2 = 15.3^2 \Rightarrow b^2 = 15.3^2 - 4.80^2 = 211.05 \Rightarrow b \approx 14.5$  m (rounded to three significant digits)


$$\sin A = \frac{a}{c} \Rightarrow \sin A = \frac{4.80}{15.3} \Rightarrow A = \sin^{-1} \frac{4.80}{15.3} \approx 18.2839^\circ \approx 18^\circ + (.2839 \cdot 60)' \approx 18^\circ 17' \approx 18^\circ 20'$$

(rounded to three significant digits)

$$\begin{aligned}\cos B &= \frac{a}{c} \Rightarrow \cos B = \frac{4.80}{15.3} \Rightarrow \\ B &= \cos^{-1} \frac{4.80}{15.3} \approx 71.7161^\circ \\ &\approx 71^\circ + (.7161 \cdot 60)' \approx 71^\circ 43' \approx 71^\circ 40'\end{aligned}$$


(rounded to three significant digits)

17. No; You need to have at least one side to solve the triangle.
18. If we are given an acute angle and a side in a right triangle, the unknown part of the triangle requiring the least work to find is the other acute angle. It may be found by subtracting the given acute angle from  $90^\circ$ .
19. Answers will vary. If you know one acute angle, the other acute angle may be found by subtracting the given acute angle from  $90^\circ$ . If you know one of the sides, then choose two of the trigonometric ratios involving sine, cosine or tangent that involve the known side in order to find the two unknown sides.
20. Answers will vary. If you know the lengths of two sides, you can set up a trigonometric ratio to solve for one of the acute angles. The other acute angle may be found by subtracting the calculated acute angle from  $90^\circ$ . With either of the two acute angles that have been determined, you can set up a trigonometric ratio along with one of the known sides to solve for the missing side.
21.  $A = 28.0^\circ$ ,  $c = 17.4$  ft




$$\begin{aligned}A + B &= 90^\circ \\ B &= 90^\circ - A \\ B &= 90^\circ - 28.0^\circ = 62.0^\circ \\ \sin A &= \frac{a}{c} \Rightarrow \sin 28.0^\circ = \frac{a}{17.4} \Rightarrow \\ a &= 17.4 \sin 28.0^\circ \approx 8.17 \text{ ft} \quad (\text{rounded to three significant digits}) \\ \cos A &= \frac{b}{c} \Rightarrow \cos 28.0^\circ = \frac{b}{17.4} \Rightarrow \\ b &= 17.4 \cos 28.0^\circ \approx 15.4 \text{ ft} \quad (\text{rounded to three significant digits})\end{aligned}$$

22.  $B = 46.0^\circ$ ,  $c = 29.7$  m  
 $A + B = 90^\circ$   
 $A = 90^\circ - B$   
 $A = 90^\circ - 46.0^\circ$   
 $= 44.0^\circ$




$$\begin{aligned}\cos B &= \frac{a}{c} \Rightarrow \cos 46.0^\circ = \frac{a}{29.7} \Rightarrow \\ a &= 29.7 \cos 46.0^\circ \approx 20.6 \text{ m} \quad (\text{rounded to three significant digits}) \\ \sin B &= \frac{b}{c} \Rightarrow \sin 46.0^\circ = \frac{b}{29.7} \Rightarrow \\ b &= 29.7 \sin 46.0^\circ \approx 21.4 \text{ m} \quad (\text{rounded to three significant digits})\end{aligned}$$

23. Solve the right triangle with  $B = 73.0^\circ$ ,  $b = 128$  in. and  $C = 90^\circ$




$$\begin{aligned}A + B &= 90^\circ \Rightarrow A = 90^\circ - B \Rightarrow \\ A &= 90^\circ - 73.0^\circ = 17.0^\circ \\ \tan B &= \frac{b}{a} \Rightarrow \tan 73.0^\circ = \frac{128}{a} \Rightarrow \\ a &= \frac{128}{\tan 73.0^\circ} \Rightarrow a = 39.1 \text{ in} \quad (\text{rounded to three significant digits}) \\ \sin B &= \frac{b}{c} \Rightarrow \sin 73.0^\circ = \frac{128}{c} \Rightarrow \\ c &= \frac{128}{\sin 73.0^\circ} \Rightarrow c = 134 \text{ in} \quad (\text{rounded to three significant digits})\end{aligned}$$

24.  $A = 61.0^\circ$ ,  $b = 39.2$  cm  
 $A + B = 90^\circ \Rightarrow B = 90^\circ - A \Rightarrow$   
 $B = 90^\circ - 61.0^\circ = 29.0^\circ$   
 $\tan A = \frac{a}{b} \Rightarrow \tan 61.0^\circ = \frac{a}{39.2} \Rightarrow$   
 $a = 39.2 \tan 61.0^\circ \approx 70.7 \text{ cm}$   
 (rounded to three significant digits)  
 $\cos A = \frac{b}{c} \Rightarrow \cos 61.0^\circ = \frac{39.2}{c} \Rightarrow$   
 $c = \frac{39.2}{\cos 61.0^\circ} \approx 80.9 \text{ cm}$   
 (rounded to three significant digits)



25.  $A = 62.5^\circ$ ,  $a = 12.7$  m



$$A + B = 90^\circ \Rightarrow B = 90^\circ - A \Rightarrow$$

$$B = 90^\circ - 62.5^\circ = 27.5^\circ$$


$$\tan A = \frac{a}{b} \Rightarrow \tan 62.5^\circ = \frac{12.7}{b} \Rightarrow$$

$$b = \frac{12.7}{\tan 62.5^\circ} \approx 6.61 \text{ m (rounded to three significant digits)}$$

$$\sin A = \frac{a}{c} \Rightarrow \sin 62.5^\circ = \frac{12.7}{c} \Rightarrow$$

$$c = \frac{12.7}{\sin 62.5^\circ} \approx 14.3 \text{ m (rounded to three significant digits)}$$

26.  $B = 51.7^\circ$ ,  $a = 28.1$  ft



$$A + B = 90^\circ \Rightarrow B = 90^\circ - B \Rightarrow$$

$$A = 90^\circ - 51.7^\circ = 38.3^\circ$$


$$\tan B = \frac{b}{a} \Rightarrow \tan 51.7^\circ = \frac{b}{28.1} \Rightarrow$$

$$b = 28.1 \tan 51.7^\circ \approx 35.6 \text{ ft (rounded to three significant digits)}$$

$$\cos B = \frac{a}{c} \Rightarrow \cos 51.7^\circ = \frac{28.1}{c} \Rightarrow$$

$$c = \frac{28.1}{\cos 51.7^\circ} \approx 45.3 \text{ ft (rounded to three significant digits)}$$

27.  $a = 13$  m,  $c = 22$  m



$$c^2 = a^2 + b^2 \Rightarrow 22^2 = 13^2 + b^2 \Rightarrow$$

$$484 = 169 + b^2 \Rightarrow 315 = b^2 \Rightarrow b \approx 18 \text{ m}$$

(rounded to two significant digits)

We will determine the measurements of both  $A$  and  $B$  by using the sides of the right triangle. In practice, once you find one of the measurements, subtract it from  $90^\circ$  to find the other.

$$\sin A = \frac{a}{c} \Rightarrow \sin A = \frac{13}{22} \Rightarrow$$

$$A \approx \sin^{-1}\left(\frac{13}{22}\right) \approx 36.2215^\circ \approx 36^\circ \text{ (rounded to two significant digits)}$$


(rounded to two significant digits)

$$\cos B = \frac{b}{c} \Rightarrow \cos B = \frac{13}{22} \Rightarrow$$

$$B \approx \cos^{-1}\left(\frac{13}{22}\right) \approx 53.7784^\circ \approx 54^\circ$$

(rounded to two significant digits)

28.  $b = 32$  ft,  $c = 51$  ft



$$c^2 = a^2 + b^2 \Rightarrow 51^2 = a^2 + 32^2 \Rightarrow$$

$$2601 = a^2 + 1024 \Rightarrow 1577 = a^2 \Rightarrow a \approx 40 \text{ ft}$$

(rounded to two significant digits)

We will determine the measurements of both  $A$  and  $B$  by using the sides of the right triangle. In practice, once you find one of the measurements, subtract it from  $90^\circ$  to find the other.

$$\cos A = \frac{b}{c} \Rightarrow \cos A = \frac{32}{51} \Rightarrow$$

$$A \approx \cos^{-1}\left(\frac{32}{51}\right) \approx 51.1377^\circ \approx 51^\circ$$


(rounded to two significant digits)

$$\sin B = \frac{b}{c} \Rightarrow \sin B = \frac{32}{51} \Rightarrow$$

$$B \approx \sin^{-1}\left(\frac{32}{51}\right) \approx 38.8623^\circ \approx 39^\circ \text{ (rounded to two significant digits)}$$

(rounded to two significant digits)

- 29.
- $a = 76.4$
- yd,
- $b = 39.3$
- yd



$$\begin{aligned} c^2 &= a^2 + b^2 \Rightarrow c = \sqrt{a^2 + b^2} \\ &= \sqrt{(76.4)^2 + (39.3)^2} = \sqrt{5836.96 + 1544.49} \\ &= \sqrt{7381.45} \approx 85.9 \text{ yd (rounded to three significant digits)} \end{aligned}$$

We will determine the measurements of both  $A$  and  $B$  by using the sides of the right triangle. In practice, once you find one of the measurements, subtract it from  $90^\circ$  to find the other.


$$\begin{aligned} \tan A &= \frac{a}{b} \Rightarrow \tan A = \frac{76.4}{39.3} \Rightarrow \\ A &\approx \tan^{-1}\left(\frac{76.4}{39.3}\right) \approx 62.7788^\circ \\ &\approx 62^\circ + (.7788 \cdot 60)' \approx 62^\circ 47' \approx 62^\circ 50' \end{aligned}$$

(rounded to three significant digits)

$$\begin{aligned} \tan B &= \frac{b}{a} \Rightarrow \tan B = \frac{39.3}{76.4} \Rightarrow \\ B &\approx \tan^{-1}\left(\frac{39.3}{76.4}\right) \approx 27.2212^\circ \\ &\approx 27^\circ + (.2212 \cdot 60)' \approx 27^\circ 13' \approx 27^\circ 10' \end{aligned}$$

(rounded to three significant digits)

- 30.
- $a = 958$
- m,
- $b = 489$
- m



$$\begin{aligned} c^2 &= a^2 + b^2 \Rightarrow c = \sqrt{a^2 + b^2} = \sqrt{958^2 + 489^2} \\ &= \sqrt{917,764 + 239,121} = \sqrt{1,156,885} \\ &\approx 1075.565887 \approx 1080 \text{ (rounded to three significant digits)} \end{aligned}$$

We will determine the measurements of both  $A$  and  $B$  by using the sides of the right triangle. In practice, once you find one of the measurements, subtract it from  $90^\circ$  to find the other.


$$\begin{aligned} \tan A &= \frac{a}{b} \Rightarrow \tan A = \frac{958}{489} \Rightarrow \\ A &\approx \tan^{-1}\left(\frac{958}{489}\right) \approx 62.9585^\circ \\ &\approx 63^\circ + (.9585 \cdot 60)' \approx 62^\circ 58' \approx 63^\circ 00' \end{aligned}$$

(rounded to three significant digits)

$$\begin{aligned} \tan B &= \frac{b}{a} \Rightarrow \tan B = \frac{489}{958} \Rightarrow \\ B &\approx \tan^{-1}\left(\frac{489}{958}\right) \approx 27.0415^\circ \\ &\approx 27^\circ + (.0415 \cdot 60)' \approx 27^\circ 02' \approx 27^\circ 00' \end{aligned}$$

(rounded to three significant digits)

- 31.
- $a = 18.9$
- cm,
- $c = 46.3$
- cm



$$\begin{aligned} c^2 &= a^2 + b^2 \Rightarrow 46.3^2 = 18.9^2 + b^2 \Rightarrow \\ 2143.69 &= 357.21 + b^2 \Rightarrow 1786.48 = b^2 \Rightarrow \\ b &\approx 42.3 \text{ cm (rounded to three significant digits)} \end{aligned}$$

$$\begin{aligned} \sin A &= \frac{a}{c} \Rightarrow \sin A = \frac{18.9}{46.3} \Rightarrow \\ A &\approx \sin^{-1}\left(\frac{18.9}{46.3}\right) \approx 24.09227^\circ \\ &\approx 24^\circ + (.09227 \cdot 60)' \approx 24^\circ 06' \approx 24^\circ 10' \end{aligned}$$

(rounded to three significant digits)


$$\begin{aligned} \cos B &= \frac{a}{c} \Rightarrow \cos B = \frac{18.9}{46.3} \Rightarrow \\ B &\approx \cos^{-1}\left(\frac{18.9}{46.3}\right) \approx 65.9077^\circ \\ &\approx 65^\circ + (.9077 \cdot 60)' \approx 65^\circ 54' \approx 65^\circ 50' \end{aligned}$$

(rounded to three significant digits)

- 32.
- $b = 219$
- cm,
- $c = 647$
- m

$$\begin{aligned} c^2 &= a^2 + b^2 \\ 647^2 &= a^2 + 219^2 \\ 418,609 &= a^2 + 47,961 \\ 370,648 &= a^2 \\ a &\approx 609 \text{ m} \end{aligned}$$

(rounded to three significant digits)



(continued on next page)

(continued from page 73)

$$\cos A = \frac{b}{c} \Rightarrow \cos A = \frac{219}{647} \Rightarrow$$

$$A \approx \cos^{-1}\left(\frac{219}{647}\right) \approx 70.2154^\circ$$

$$\approx 70^\circ + (.2154 \cdot 60)' \approx 70^\circ 13' \approx 70^\circ 10'$$

(rounded to three significant digits)


$$\sin B = \frac{b}{c} \Rightarrow \sin B = \frac{219}{647} \Rightarrow$$

$$B \approx \sin^{-1}\left(\frac{219}{647}\right) \approx 19.7846^\circ$$

$$\approx 19^\circ + (.7846 \cdot 60)' \approx 19^\circ 47' \approx 19^\circ 50'$$

(rounded to three significant digits)

33.  $A = 53^\circ 24'$ ,  $c = 387.1$  ft  
 $A + B = 90^\circ$   
 $B = 90^\circ - A$   
 $B = 90^\circ - 53^\circ 24'$   
 $= 89^\circ 60' - 53^\circ 24'$   
 $= 36^\circ 36'$




$$\sin A = \frac{a}{c} \Rightarrow \sin 53^\circ 24' = \frac{a}{387.1} \Rightarrow$$

$$a = 387.1 \sin 53^\circ 24' \approx 310.8 \text{ ft (rounded to four significant digits)}$$

$$\cos A = \frac{b}{c} \Rightarrow \cos 53^\circ 24' = \frac{b}{387.1} \Rightarrow$$

$$b = 387.1 \cos 53^\circ 24' \approx 230.8 \text{ ft (rounded to four significant digits)}$$

34.  $A = 13^\circ 47'$ ,  $c = 1285$  m



$$A + B = 90^\circ \Rightarrow B = 90^\circ - A \Rightarrow$$

$$B = 90^\circ - 13^\circ 47' = 89^\circ 60' - 13^\circ 47'$$

$$= 76^\circ 13'$$


$$\sin A = \frac{a}{c} \Rightarrow \sin 13^\circ 47' = \frac{a}{1285} \Rightarrow$$

$$a = 1285 \sin 13^\circ 47' \approx 306.2 \text{ m (rounded to four significant digits)}$$

$$\cos A = \frac{b}{c} \Rightarrow \cos 13^\circ 47' = \frac{b}{1285} \Rightarrow$$

$$b = 1285 \cos 13^\circ 47' \approx 1248 \text{ m (rounded to four significant digits)}$$

35.  $B = 39^\circ 09'$ ,  $c = .6231$  m  
 $A + B = 90^\circ$   
 $B = 90^\circ - A$   
 $B = 90^\circ - 39^\circ 09'$   
 $= 89^\circ 60' - 39^\circ 09'$   
 $= 50^\circ 51'$




$$\sin B = \frac{b}{c} \Rightarrow \sin 39^\circ 09' = \frac{b}{.6231} \Rightarrow$$

$$b = .6231 \sin 39^\circ 09' \approx .3934 \text{ m (rounded to four significant digits)}$$

$$\cos B = \frac{a}{c} \Rightarrow \cos 39^\circ 09' = \frac{a}{.6231} \Rightarrow$$

$$a = .6231 \cos 39^\circ 09' \approx .4832 \text{ m (rounded to four significant digits)}$$

36.  $B = 82^\circ 51'$ ,  $c = 4.825$  cm



$$A + B = 90^\circ \Rightarrow B = 90^\circ - A \Rightarrow$$

$$B = 90^\circ - 82^\circ 51' = 89^\circ 60' - 82^\circ 51'$$

$$= 7^\circ 9'$$

$$\sin B = \frac{b}{c} \Rightarrow \sin 82^\circ 51' = \frac{b}{4.825} \Rightarrow$$

$$b = 4.825 \sin 82^\circ 51' \approx 4.787 \text{ m (rounded to four significant digits)}$$


$$\cos B = \frac{a}{c} \Rightarrow \cos 82^\circ 51' = \frac{a}{4.825} \Rightarrow$$

$$a = 4.825 \cos 82^\circ 51' \approx .6006 \text{ m (rounded to four significant digits)}$$

37. The angle of elevation from X to Y (with Y above X) is the acute angle formed by ray XY and a horizontal ray with endpoint at X.
38. The angle of elevation from X to Y is the acute angle formed by ray XY and a horizontal ray with endpoint at X. Therefore, the angle of elevation cannot be more than  $90^\circ$ .
39. Answers will vary. The angle of elevation and the angle of depression are measured between the line of sight and a horizontal line. So, in the diagram, lines AD and CB are both horizontal. Hence, they are parallel. The line formed by AB is a transversal and angles DAB and ABC are alternate interior angle and thus have the same measure.
40. The angle of depression is measured between the line of sight and a horizontal line. This angle is measured between the line of sight and a vertical line.


41.  $\sin 43^\circ 50' = \frac{d}{13.5}$   
 $d = 13.5 \sin 43^\circ 50' \approx 9.3496000$   
 The ladder goes up the wall 9.35 m. (rounded to three significant digits)

42.  $T = 32^\circ 10'$  and  $S = 57^\circ 50'$



Since  $S + T = 32^\circ 10' + 57^\circ 50' = 89^\circ 60' = 90^\circ$ , triangle  $RST$  is a right triangle. Thus, we have  $\tan 32^\circ 10' = \frac{RS}{53.1}$   
 $RS = 53.1 \tan 32^\circ 10' \approx 33.395727$   
 The distance across the lake is 33.4 m. (rounded to three significant digits)

43. Let  $x$  represent the horizontal distance between the two buildings and  $y$  represent the height of the portion of the building across the street that is higher than the window.




$$\tan 20.0^\circ = \frac{30.0}{x} \Rightarrow x = \frac{30.0}{\tan 20.0^\circ} \approx 82.4$$

$$\tan 50.0^\circ = \frac{y}{x} \Rightarrow y = x \tan 50.0^\circ = \left( \frac{30.0}{\tan 20.0^\circ} \right) \tan 50.0^\circ$$

$$\text{height} = y + 30.0 = \left( \frac{30.0}{\tan 20.0^\circ} \right) \tan 50.0^\circ + 30.0 \approx 128.2295$$

The height of the building across the street is about 128 ft. (rounded to three significant digits)

44. Let  $x$  = the diameter of the sun.



Since the included angle is  $32'$ ,  $\frac{1}{2}(32') = 16'$ .


We will use this angle,  $d$ , and half of the diameter to set up the following equation.

$$\frac{\frac{1}{2}x}{92,919,800} = \tan 16' \Rightarrow x = 2(92,919,800)(\tan 16') \approx 864,943.0189$$

The diameter of the sun is about 864,900 mi. (rounded to four significant digits)

45. The altitude of an isosceles triangle bisects the base as well as the angle opposite the base. The two right triangles formed have interior angles which have the same measure. The lengths of the corresponding sides also have the same measure. Since the altitude bisects the base, each leg (base) of the right triangles is  $\frac{42.36}{2} = 21.18$  in.

Let  $x$  = the length of each of the two equal sides of the isosceles triangle.




$$\cos 38.12^\circ = \frac{21.18}{x} \Rightarrow x \cos 38.12^\circ = 21.18 \Rightarrow x = \frac{21.18}{\cos 38.12^\circ} \approx 26.921918$$

The length of each of the two equal sides of the triangle is 26.92 in. (rounded to four significant digits)



46. The altitude of an isosceles triangle bisects the base as well as the angle opposite the base. The two right triangles formed have interior angles which have the same measure. The lengths of the corresponding sides also have the same measure. Since the altitude bisects the base, each leg (base) of the right triangles are  $\frac{184.2}{2} = 92.10$  cm. Each angle opposite to the base of the right triangles measures  $\frac{1}{2}(68^\circ 44') = 34^\circ 22'$ . Let  $h$  = the altitude.




In triangle ABC,

$$\tan 34^\circ 22' = \frac{92.10}{h} \Rightarrow h \tan 34^\circ 22' = 92.10 \Rightarrow h = \frac{92.10}{\tan 34^\circ 22'} \approx 134.67667$$


The altitude of the triangle is 134.7 cm. (rounded to four significant digits)

47. In order to find the angle of elevation,  $\theta$ , we need to first find the length of the diagonal of the square base. The diagonal forms two isosceles right triangles. Each angle formed by a side of the square and the diagonal measures  $45^\circ$ .



By the Pythagorean theorem,  
 $700^2 + 700^2 = d^2 \Rightarrow 2 \cdot 700^2 = d^2 \Rightarrow d = \sqrt{2 \cdot 700^2} \Rightarrow d = 700\sqrt{2}$

Thus, length of the diagonal is  $700\sqrt{2}$  ft. To find the angle,  $\theta$ , we consider the following isosceles triangle.



The height of the pyramid bisects the base of this triangle and forms two right triangles. We can use one of these triangles to find the angle of elevation,  $\theta$ .

$$\tan \theta = \frac{200}{350\sqrt{2}} \approx .4040610178$$

$$\theta \approx \tan^{-1}(.4040610178) \approx 22.0017$$


Rounding this figure to two significant digits, we have  $\theta \approx 22^\circ$ .

48. Let  $y$  = the height of the spotlight (this measurement starts 6 feet above ground)

$$\tan 30.0^\circ = \frac{y}{1000}$$

$$y = 1000 \cdot \tan 30.0^\circ \approx 577.3502$$

Rounding this figure to three significant digits, we have  $y \approx 577$ .




However, the observer's eye-height is 6 feet from the ground, so the cloud ceiling is  $577 + 6 = 583$  ft.

49. Let  $h$  represent the height of the tower. In triangle  $ABC$  we have


$$\tan 34.6^\circ = \frac{h}{40.6}$$

$$h = 40.6 \tan 34.6^\circ \approx 28.0081$$

The height of the tower is 28.0 m. (rounded to three significant digits)



50. Let  $d$  = the distance from the top  $B$  of the building to the point on the ground  $A$ .



In triangle  $ABC$ ,

$$\sin 32^\circ 30' = \frac{252}{d}$$

$$d = \frac{252}{\sin 32^\circ 30'} \approx 469.0121$$


The distance from the top of the building to the point on the ground is 469 m. (rounded to three significant digits)

51. Let  $x$  = the length of the shadow cast by Diane Carr.

$$\tan 23.4^\circ = \frac{5.75}{x}$$


$$x \tan 23.4^\circ = 5.75$$

$$x = \frac{5.75}{\tan 23.4^\circ} \approx 13.2875$$



The length of the shadow cast by Diane Carr is 13.3 ft. (rounded to three significant digits)

52. Let  $x$  = the horizontal distance that the plan must fly to be directly over the tree.




$$\tan 13^\circ 50' = \frac{10,500}{x}$$

$$x \tan 13^\circ 50' = 10,500$$

$$x = \frac{10,500}{\tan 13^\circ 50'} \approx 42,641.2351$$

The horizontal distance that the plan must fly to be directly over the tree is 42,600 ft. (rounded to three significant digits)

53. Let  $x$  = the height of the taller building;  $h$  = the difference in height between the shorter and taller buildings;  $d$  = the distance between the buildings along the ground.



$$\frac{28.0}{d} = \tan 14^\circ 10' \Rightarrow 28.0 = d \tan 14^\circ 10' \Rightarrow$$

$$d = \frac{28.0}{\tan 14^\circ 10'} \approx 110.9262493 \text{ m}$$

(We hold on to these digits for the intermediate steps.) To find  $h$ , solve


$$\frac{h}{d} = \tan 46^\circ 40'$$

$$h = d \tan 46^\circ 40' \approx (110.9262493) \tan 46^\circ 40'$$

$$\approx 117.5749$$

Thus, the value of  $h$  rounded to three significant digits is 118 m. Since  $x = h + 28.0 = 118 + 28.0 \approx 146 \text{ m}$ , the height of the taller building is 146 m.

54. Let  $\theta$  = the angle of depression.




$$\tan \theta = \frac{39.82}{51.74} \approx .7696173174$$

$$\theta = \tan^{-1}(.7696173174)$$

$$\theta \approx 37.58^\circ \approx 37^\circ 35'$$

55. (a) Let  $x$  = the height of the peak above 14,545 ft.



Since the diagonal of the right triangle formed is in miles, we must first convert this measurement to feet. Since there are 5280 ft in one mile, we have the length of the diagonal is  $27.0134(5280) =$


142,630.752. Find the value of  $x$  by

$$\begin{aligned} \text{solving } \sin 5.82^\circ &= \frac{x}{142,630.752} \\ x &= 142,630.752 \sin 5.82^\circ \\ &\approx 14,463.2674 \end{aligned}$$

Thus, the value of  $x$  rounded to five significant digits is 14,463 ft. Thus, the total height is about  $14,545 + 14,463 = 29,008 \approx 29,000$  ft.

- (b) The curvature of the earth would make the peak appear shorter than it actually is. Initially the surveyors did not think Mt. Everest was the tallest peak in the Himalayas. It did not look like the tallest peak because it was farther away than the other large peaks.

56. Let  $x$  = the distance from the assigned target.



In triangle  $ABC$ , we have

$$\begin{aligned} \tan 0^\circ 0' 30'' &= \frac{x}{234,000} \\ x &= 234,000 \tan 0^\circ 0' 30'' \approx 34.0339 \end{aligned}$$


The distance from the assigned target is 34.0 mi. (rounded to three significant digits)

### Section 2.5: Further Applications of Right Triangles

1. It should be shown as an angle measured clockwise from due north.
2. It should be shown measured from north (or south) in the east (or west) direction.
3. A sketch is important to show the relationships among the given data and the unknowns.


4. The angle of elevation (or depression) from  $X$  to  $Y$  is measured from the horizontal line through  $X$  to the ray  $XY$ .

5.  $(-4, 0)$




The bearing of the airplane measured in a clockwise direction from due north is  $270^\circ$ . The bearing can also be expressed as  $N 90^\circ W$ , or  $S 90^\circ W$ .

6.  $(-3, -3)$




The bearing of the airplane measured in a clockwise direction from due north is  $225^\circ$ . The bearing can also be expressed as  $S 45^\circ W$ .


7.  $(-5, 5)$




The bearing of the airplane measured in a clockwise direction from due north is  $315^\circ$ . The bearing can also be expressed as  $N 45^\circ W$ .

8.  $(0, -2)$ 



The bearing of the airplane measured in a clockwise direction from due north is  $180^\circ$ . The bearing can also be expressed as  $S 0^\circ E$  or  $S 0^\circ W$ .

 9.  $(0, 4)$ 



The bearing of the airplane measured in a clockwise direction from due north is  $0^\circ$ . The bearing can also be expressed as  $N 0^\circ E$  or  $N 0^\circ W$ .

 10.  $(2, 2)$ 



The bearing of the airplane measured in a clockwise direction from due north is  $45^\circ$ . The bearing can also be expressed as  $N 45^\circ E$ .

 11.  $(2, -2)$ 


The bearing of the airplane measured in a clockwise direction from due north is  $135^\circ$ . The bearing can also be expressed as  $S 45^\circ E$ .

 12.  $(5, 0)$ 


The bearing of the airplane measured in a clockwise direction from due north is  $90^\circ$ . The bearing can also be expressed as  $N 90^\circ E$ , or  $S 90^\circ E$ .

 13. All points whose bearing from the origin is  $240^\circ$  lie in quadrant III.


The reference angle,  $\theta'$ , is  $30^\circ$ . For any point,  $(x, y)$  on the ray  $\frac{x}{r} = -\cos \theta'$  and

$\frac{y}{r} = -\sin \theta'$ , where  $r$  is the distance from the point to the origin. Let  $r = 2$ , so

$$\frac{x}{r} = -\cos \theta'$$

$$x = -r \cos \theta' = -2 \cos 30^\circ = -2 \cdot \frac{\sqrt{3}}{2} = -\sqrt{3}$$

$$\frac{y}{r} = -\sin \theta'$$


$$y = -r \sin \theta' = -2 \sin 30^\circ = -2 \cdot \frac{1}{2} = -1$$

(continued on next page)

(continued from page 79)

Thus, a point on the ray is  $(-\sqrt{3}, -1)$ . Since the ray contains the origin, the equation is of the form  $y = mx$ . Substituting the point  $(-\sqrt{3}, -1)$ , we have  $-1 = m(-\sqrt{3}) \Rightarrow m = \frac{-1}{-\sqrt{3}} = \frac{1}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{\sqrt{3}}{3}$ . Thus, the equation of the ray is  $y = \frac{\sqrt{3}}{3}x, x \leq 0$  (since the ray lies in quadrant III).

14. All points whose bearing from the origin is  $150^\circ$  lie in quadrant IV.



The reference angle,  $\theta'$ , is  $60^\circ$ . For any point,  $(x, y)$  on the ray  $\frac{x}{r} = \cos \theta'$  and  $\frac{y}{r} = -\sin \theta'$ , where  $r$  is the distance from the point to the origin. Let  $r = 2$ , so

$$\frac{x}{r} = \cos \theta' \Rightarrow x = r \cos \theta' = 2 \cos 60^\circ = 2 \cdot \frac{1}{2} = 1$$

$$\frac{y}{r} = -\sin \theta'$$


$$y = -r \sin \theta' = -2 \sin 60^\circ = -2 \cdot \frac{\sqrt{3}}{2} = -\sqrt{3}$$

Thus, a point on the ray is  $(1, -\sqrt{3})$ .

Since the ray contains the origin, the equation is of the form  $y = mx$ . Substituting the point  $(1, -\sqrt{3})$ , we have  $-\sqrt{3} = m(1) \Rightarrow m = -\sqrt{3}$ .

Thus, the equation of the ray is  $y = -\sqrt{3}x, x \geq 0$  (since the ray lies in quadrant IV).

15. Let  $x$  = the distance the plane is from its starting point. In the figure, the measure of angle  $ACB$  is  $40^\circ + (180^\circ - 130^\circ) = 40^\circ + 50^\circ = 90^\circ$ . Therefore, triangle  $ACB$  is a right triangle.




Since  $d = rt$ , the distance traveled in 1.5 hr is  $(1.5 \text{ hr})(110 \text{ mph}) = 165 \text{ mi}$ . The distance traveled in 1.3 hr is  $(1.3 \text{ hr})(110 \text{ mph}) = 143 \text{ mi}$ .

Using the Pythagorean theorem, we have  $x^2 = 165^2 + 143^2 \Rightarrow x^2 = 27,225 + 20,449 \Rightarrow x^2 = 47,674 \Rightarrow x \approx 218.3438$

The plane is 220 mi from its starting point. (rounded to two significant digits)

16. Let  $x$  = the distance from the starting point. In the figure, the measure of angle  $ACB$  is  $27^\circ + (180^\circ - 117^\circ) = 27^\circ + 63^\circ = 90^\circ$ .


Therefore, triangle  $ACB$  is a right triangle.



Applying the Pythagorean theorem, we have  $x^2 = 50^2 + 140^2 \Rightarrow x^2 = 2500 + 19,600 \Rightarrow x^2 = 22,100 \Rightarrow x = \sqrt{22,100} \approx 148.6607$

The distance of the end of the trip from the starting point is 150 km. (rounded to two significant digits)

17. Let  $x$  = distance the ships are apart. In the figure, the measure of angle  $CAB$  is  $130^\circ - 40^\circ = 90^\circ$ . Therefore, triangle  $CAB$  is a right triangle.



Since  $d = rt$ , the distance traveled by the first ship in 1.5 hr is

$(1.5 \text{ hr})(18 \text{ knots}) = 27$  nautical mi and the second ship is

$(1.5 \text{ hr})(26 \text{ knots}) = 39$  nautical mi.


Applying the Pythagorean theorem, we have

$$x^2 = 27^2 + 39^2 \Rightarrow x^2 = 729 + 1521 \Rightarrow$$

$$x^2 = 2250 \Rightarrow x = \sqrt{2250} \approx 47.4342$$

The ships are 47 nautical mi apart. (rounded to 2 significant digits)

18. Let  $C$  = the location of the ship, and let  $c$  = the distance between the lighthouses.



$$m\angle BAC = 180^\circ - 129^\circ 43' \\ = 179^\circ 60' - 129^\circ 43' = 50^\circ 17'$$


Since  $50^\circ 17' + 39^\circ 43' = 90^\circ$ , we have a right triangle. Thus,

$$\sin 39^\circ 43' = \frac{3742}{c} \Rightarrow c \sin 39^\circ 43' = 3742 \Rightarrow$$

$$c = \frac{3742}{\sin 39^\circ 43'} \approx 5856.1020$$

The distance between the lighthouses is 5856 m (rounded to four significant digits).

19. Draw triangle  $WDG$  with  $W$  representing Winston-Salem,  $D$  representing Danville, and  $G$  representing Goldsboro. Name any point  $X$  on the line due south from  $D$ .



Since the bearing from  $W$  to  $D$  is  $42^\circ$  (equivalent to  $N 42^\circ E$ ), angle  $WDX$  measures  $42^\circ$ . Since angle  $XDG$  measures  $48^\circ$ , the measure of angle  $D$  is  $42^\circ + 48^\circ = 90^\circ$ .

Thus, triangle  $WDG$  is a right triangle.


Using  $d = rt$  and the Pythagorean theorem, we have

$$WG = \sqrt{(WD)^2 + (DG)^2} \\ = \sqrt{[65(1.1)]^2 + [65(1.8)]^2}$$

$$WG = \sqrt{71.5^2 + 117^2} = \sqrt{5112.25 + 13,689} \\ = \sqrt{18,801.25} \approx 137$$

The distance from Winston-Salem to Goldsboro is approximately 140 mi. (rounded to two significant digits)

20. Let  $x$  = the distance from Atlanta to Augusta.



The line from Atlanta to Macon makes an angle of  $27^\circ + 63^\circ = 90^\circ$ , with the line from Macon to Augusta. Since  $d = rt$ , the distance from Atlanta to Macon is  $60\left(1\frac{1}{4}\right) = 75$  mi.

The distance from Macon to Augusta is  $60\left(1\frac{3}{4}\right) = 105$  mi.


Use the Pythagorean theorem to find  $x$ :

$$x^2 = 75^2 + 105^2 \Rightarrow x^2 = 5635 + 11,025 \Rightarrow$$

$$x^2 = 16,660 \approx 129.0349$$

The distance from Atlanta to Augusta is 130 mi. (rounded to two significant digits)

21. Let  $x$  = distance between the two ships.



The angle between the bearings of the ships is  $180^\circ - (28^\circ 10' + 61^\circ 50') = 90^\circ$ . The triangle formed is a right triangle. The distance traveled at 24.0 mph is

$(4 \text{ hr})(24.0 \text{ mph}) = 96$  mi. The distance traveled at 28.0 mph is

$(4 \text{ hr})(28.0 \text{ mph}) = 112$  mi.


Applying the Pythagorean theorem we have

$$x^2 = 96^2 + 112^2 \Rightarrow x^2 = 9216 + 12,544 \Rightarrow$$

$$x^2 = 21,760 \Rightarrow x = \sqrt{21,760} \approx 147.5127$$

The ships are 148 mi apart. (rounded to three significant digits)

22. Let  $C$  = the location of the transmitter;  
 $a$  = the distance of the transmitter from  $B$ .




The measure of angle  $CBA$  is  
 $90^\circ - 53^\circ 40' = 89^\circ 60' - 53^\circ 40' = 36^\circ 20'$ .  
 The measure of angle  $CAB$  is  
 $90^\circ - 36^\circ 20' = 89^\circ 60' - 36^\circ 20' = 53^\circ 40'$ .  
 Since  $A + B = 90^\circ$ , so  $C = 90^\circ$ . Thus, we have

$$\sin A = \frac{a}{2.50} \Rightarrow \sin 53^\circ 40' = \frac{a}{2.50} \Rightarrow a = 2.50 \sin 53^\circ 40' \approx 2.0140$$

The distance of the transmitter from  $B$  is 2.01 mi. (rounded to 3 significant digits)

23.  $ax = b + cx \Rightarrow ax - cx = b \Rightarrow x(a - c) = b \Rightarrow x = \frac{b}{a - c}$

24. Suppose we have a line that has  $x$ -intercept  $a$  and  $y$ -intercept  $b$ . Assume for the following diagram that  $a$  and  $b$  are both positive. This is not a necessary condition, but it makes the visualization easier.




$\tan \theta = -\tan(180^\circ - \theta) = -\tan \theta'$ . This is because the angle represented by  $180^\circ - \theta$  terminates in quadrant II if  $0^\circ < \theta < 90^\circ$ . If  $90^\circ < \theta < 180^\circ$ , then the angle represented by  $180^\circ - \theta$  terminates in quadrant I. Thus,  $\tan \theta$  and  $\tan(180^\circ - \theta)$  are opposite in sign.

The slope of the line is  $m = -\frac{b}{a}$ , and  $\tan \theta = -\tan(180^\circ - \theta) = -\tan \theta' = -\frac{b}{a}$ . Thus,  $m = -\frac{b}{a} = \tan \theta$ . The point-slope form of the equation of a line is  $y - y_1 = m(x - x_1)$ .

Substituting  $\tan \theta$  for  $m$  into  $y - y_1 = m(x - x_1)$ , we have  $y - y_1 = -\tan \theta(x - x_1)$ .

The line passes through  $(a, 0)$ , so  $y - y_1 = \tan \theta(x - x_1) \Rightarrow y - 0 = \tan \theta(x - a) \Rightarrow y = \tan \theta(x - a)$ .

25. From exercise 24, we have  $y = \tan \theta(x - a) \Rightarrow y = \tan 35^\circ(x - 25)$
26. From exercise 24, we have  $y = \tan \theta(x - a) \Rightarrow y = \tan 15^\circ(x - 5)$
27. Algebraic solution:  
 Let  $x$  = the side adjacent to  $49.2^\circ$  in the smaller triangle.



In the larger right triangle, we have

$$\tan 29.5^\circ = \frac{h}{392 + x} \Rightarrow h = (392 + x) \tan 29.5^\circ$$

In the smaller right triangle, we have

$$\tan 49.2^\circ = \frac{h}{x} \Rightarrow h = x \tan 49.2^\circ$$

Substituting, we have


$$\begin{aligned} x \tan 49.2^\circ &= (392 + x) \tan 29.5^\circ \\ x \tan 49.2^\circ &= 392 \tan 29.5^\circ + x \tan 29.5^\circ \\ x \tan 49.2^\circ - x \tan 29.5^\circ &= 392 \tan 29.5^\circ \\ x(\tan 49.2^\circ - \tan 29.5^\circ) &= 392 \tan 29.5^\circ \\ x &= \frac{392 \tan 29.5^\circ}{\tan 49.2^\circ - \tan 29.5^\circ} \end{aligned}$$



Now substitute this expression for  $x$  in the equation for the smaller triangle to obtain  $h = x \tan 49.2^\circ$

$$\begin{aligned} h &= \frac{392 \tan 29.5^\circ}{\tan 49.2^\circ - \tan 29.5^\circ} \cdot \tan 49.2^\circ \\ &\approx 433.4762 \approx 433 \text{ ft (rounded to three significant digits).} \end{aligned}$$

Graphing calculator solution:

The first line considered is  $y = (\tan 29.5^\circ)x$  and the second is  $y = (\tan 29.5^\circ)(x - 392)$ .






The height of the triangle is 433 ft (rounded to three significant digits).

28. Algebraic solution:

Let  $x$  = the side adjacent to  $52.5^\circ$  in the smaller triangle.



In the larger right triangle, we have

$$\tan 41.2^\circ = \frac{h}{168 + x} \Rightarrow h = (168 + x) \tan 41.2^\circ.$$

In the smaller right triangle, we have

$$\tan 52.5^\circ = \frac{h}{x} \Rightarrow h = x \tan 52.5^\circ.$$

Substituting, we have

$$x \tan 52.5^\circ = (168 + x) \tan 41.2^\circ$$

$$x \tan 52.5^\circ = 168 \tan 41.2^\circ + x \tan 41.2^\circ$$

$$x \tan 52.5^\circ - x \tan 41.2^\circ = 168 \tan 41.2^\circ$$

$$x(\tan 52.5^\circ - \tan 41.2^\circ) = 168 \tan 41.2^\circ$$

$$x = \frac{168 \tan 41.2^\circ}{\tan 52.5^\circ - \tan 41.2^\circ}$$

Now substitute this expression for  $x$  in the equation for the smaller triangle to obtain

$$h = x \tan 52.5^\circ$$




$$h = \frac{168 \tan 41.2^\circ}{\tan 52.5^\circ - \tan 41.2^\circ} \cdot \tan 52.5^\circ$$

$$\approx 448.0432 \approx 448 \text{ m (rounded to three significant digits.)}$$

Graphing calculator solution:


The first line considered is  $y = (\tan 41.2^\circ)x$

and the second is  $y = (\tan 52.5^\circ)(x - 168)$ .



The height of the triangle is 448 m (rounded to three significant digits).

29. Let  $x$  = the distance from the closer point on the ground to the base of height  $h$  of the pyramid.



In the larger right triangle, we have

$$\tan 21^\circ 10' = \frac{h}{135 + x} \Rightarrow h = (135 + x) \tan 21^\circ 10'$$

In the smaller right triangle, we have

$$\tan 35^\circ 30' = \frac{h}{x} \Rightarrow h = x \tan 35^\circ 30'.$$

Substitute for  $h$  in this equation, and solve for  $x$  to obtain the following.

$$(135 + x) \tan 21^\circ 10' = x \tan 35^\circ 30'$$

$$135 \tan 21^\circ 10' + x \tan 21^\circ 10' = x \tan 35^\circ 30'$$

$$135 \tan 21^\circ 10' = x \tan 35^\circ 30' - x \tan 21^\circ 10'$$

$$135 \tan 21^\circ 10' = x(\tan 35^\circ 30' - \tan 21^\circ 10')$$

$$\frac{135 \tan 21^\circ 10'}{\tan 35^\circ 30' - \tan 21^\circ 10'} = x$$

(continued on next page)




(continued from page 83)

Substitute for  $x$  in the equation for the smaller triangle.

$$h = \frac{135 \tan 21^\circ 10'}{\tan 35^\circ 30' - \tan 21^\circ 10'} \tan 35^\circ 30' \\ \approx 114.3427$$

The height of the pyramid is 114 ft. (rounded to three significant digits)

30. Let  $x$  = the distance traveled by the whale as it approaches the tower;  $y$  = the distance from the tower to the whale as it turns.



$$\frac{68.7}{y} = \tan 35^\circ 40' \Rightarrow 68.7 = y \tan 35^\circ 40' \Rightarrow$$

$$y = \frac{68.7}{\tan 35^\circ 40'} \quad \text{and} \quad \frac{68.7}{x+y} = \tan 15^\circ 50'$$


$$68.7 = (x+y) \tan 15^\circ 50'$$

$$x+y = \frac{68.7}{\tan 15^\circ 50'} \Rightarrow x = \frac{68.7}{\tan 15^\circ 50'} - y$$

$$x = \frac{68.7}{\tan 15^\circ 50'} - \frac{68.7}{\tan 35^\circ 40'} \approx 146.5190$$

The whale traveled 147 m as it approached the lighthouse. (rounded to three significant digits)

31. Let  $x$  = the height of the antenna;  $h$  = the height of the house.



In the smaller right triangle, we have

$$\tan 18^\circ 10' = \frac{h}{28} \Rightarrow h = 28 \tan 18^\circ 10'$$

In the larger right triangle, we have

$$\tan 27^\circ 10' = \frac{x+h}{28} \Rightarrow x+h = 28 \tan 27^\circ 10' \Rightarrow$$


$$x = 28 \tan 27^\circ 10' - h$$

$$x = 28 \tan 27^\circ 10' - 28 \tan 18^\circ 10'$$

$$\approx 5.1816$$

The height of the antenna is 5.18 m. (rounded to three significant digits)

32. Let  $x$  = the height of Mt. Whitney above the level of the road;  $y$  = the distance shown in the figure below.



In triangle  $ADC$ ,

$$\tan 22^\circ 40' = \frac{x}{y} \Rightarrow y \tan 22^\circ 40' = x \Rightarrow$$

$$y = \frac{x}{\tan 22^\circ 40'} \quad (1)$$

In triangle  $ABC$

$$\tan 10^\circ 50' = \frac{x}{y+7.00}$$

$$(y+7.00) \tan 10^\circ 50' = x$$

$$y \tan 10^\circ 50' + 7.00 \tan 10^\circ 50' = x$$

$$\frac{x - 7.00 \tan 10^\circ 50'}{\tan 10^\circ 50'} = y \quad (2)$$

Setting equations 1 and 2 equal, we have

$$\frac{x}{\tan 22^\circ 40'} = \frac{x - 7.00 \tan 10^\circ 50'}{\tan 10^\circ 50'}$$

$$x \tan 10^\circ 50' = x \tan 22^\circ 40' - 7.00(\tan 10^\circ 50')(\tan 22^\circ 40')$$

$$7.00(\tan 10^\circ 50')(\tan 22^\circ 40')$$

$$= x \tan 22^\circ 40' - x \tan 10^\circ 50'$$

$$7.00(\tan 10^\circ 50')(\tan 22^\circ 40')$$

$$= x(\tan 22^\circ 40' - \tan 10^\circ 50')$$

$$x = \frac{7.00(\tan 10^\circ 50')(\tan 22^\circ 40')}{\tan 22^\circ 40' - \tan 10^\circ 50'}$$

$$x \approx 2.4725$$

The height of the top of Mt. Whitney above road level is 2.47 km. (rounded to three significant digits)

33. (a) From the figure in the text,

$$d = \frac{b}{2} \cot \frac{\alpha}{2} + \frac{b}{2} \cot \frac{\beta}{2} \\ = \frac{b}{2} \left( \cot \frac{\alpha}{2} + \cot \frac{\beta}{2} \right)$$

- (b) Using the result of part (a), let

$$\alpha = 37'48'', \beta = 42'3'', \text{ and } b = 2.000$$

$$d = \frac{b}{2} \left( \cot \frac{\alpha}{2} + \cot \frac{\beta}{2} \right) \Rightarrow$$


$$d = \frac{2.000}{2} \left( \cot \frac{37'48''}{2} + \cot \frac{42'3''}{2} \right)$$

$$= \cot .315^\circ + \cot .3504166667$$

$$\approx 345.3951$$

The distance between the two point  $P$  and  $Q$  is about 345.4 cm.

34. Let  $h$  = the minimum height above the surface of the earth so a pilot at  $A$  can see an object on the horizon at  $C$ .




Using the Pythagorean theorem, we have

$$\begin{aligned}(4.00 \times 10^3 + h)^2 &= (4.00 \times 10^3)^2 + 125^2 \\(4000 + h)^2 &= 4000^2 + 125^2 \\(4000 + h)^2 &= 16,000,000 + 15,625 \\(4000 + h)^2 &= 16,015,625 \\4000 + h &= \sqrt{16,015,625} \\h &= \sqrt{16,015,625} - 4000 \\&\approx 4001.9526 - 4000 = 1.9526\end{aligned}$$

The minimum height above the surface of the earth would be 1.95 mi. (rounded to 3 significant digits)

35. Let  $x$  = the minimum distance that a plant needing full sun can be placed from the fence.




$$\begin{aligned}\tan 23^\circ 20' &= \frac{4.65}{x} \Rightarrow x \tan 23^\circ 20' = 4.65 \Rightarrow \\x &= \frac{4.65}{\tan 23^\circ 20'} \approx 10.7799\end{aligned}$$

The minimum distance is 10.8 ft. (rounded to three significant digits)

36.  $\tan A = \frac{1.0837}{1.4923} \approx .7261944649$   
 $A \approx \tan^{-1}(.7261944649) \approx 35.987^\circ$   
 $\approx 35^\circ 59.2' \approx 35^\circ 59' 10''$   
 $\tan B = \frac{1.4923}{1.0837} \approx 1.377041617$   
 $B \approx \tan^{-1}(1.377041617) \approx 54.013^\circ$   
 $\approx 54^\circ 00.8' \approx 54^\circ 00' 50''$

37. (a) If  $\theta = 37^\circ$ , then  $\frac{\theta}{2} = \frac{37^\circ}{2} = 18.5^\circ$ .




To find the distance between  $P$  and  $Q$ ,  $d$ , we first note that angle  $QPC$  is a right angle. Hence, triangle  $QPC$  is a right triangle and we can solve

$$\begin{aligned}\tan 18.5^\circ &= \frac{d}{965} \\d &= 965 \tan 18.5^\circ \approx 322.8845\end{aligned}$$

The distance between  $P$  and  $Q$ , is 320 ft. (rounded to two significant digits)


- (b) Since we are dealing with a circle, the distance between  $M$  and  $C$  is  $R$ . If we let  $x$  be the distance from  $N$  to  $M$ , then the distance from  $C$  to  $N$  will be  $R - x$ .



Since triangle  $CNP$  is a right triangle, we can set up the following equation.

$$\begin{aligned}\cos \frac{\theta}{2} &= \frac{R-x}{R} \Rightarrow R \cos \frac{\theta}{2} = R-x \Rightarrow \\x &= R - R \cos \frac{\theta}{2} \Rightarrow x = R \left( 1 - \cos \frac{\theta}{2} \right)\end{aligned}$$

38. Let  $y$  = the common hypotenuse of the two right triangles.



$$\cos 30^\circ 50' = \frac{198.4}{y}$$

$$y = \frac{198.4}{\cos 30^\circ 50'} \approx 231.0571948$$

To find  $x$ , first find the angle opposite  $x$  in the right triangle:

$$52^\circ 20' - 30^\circ 50' = 51^\circ 80' - 30^\circ 50' = 21^\circ 30'$$

$$\sin 21^\circ 30' = \frac{x}{y}$$

$$\sin 21^\circ 30' \approx \frac{x}{231.0571948}$$

$$x \approx 231.0571948 \sin 21^\circ 30'$$

$$\approx 84.6827$$

The length  $x$  is approximate 84.7 m. (rounded)

39. (a)  $\theta \approx \frac{57.3S}{R} = \frac{57.3(336)}{600} = 32.088^\circ$

$$d = R \left( 1 - \cos \frac{\theta}{2} \right)$$

$$= 600(1 - \cos 16.044^\circ) \approx 23.3702 \text{ ft}$$

The distance is 23 ft. (rounded to two significant digits)

(b)  $\theta \approx \frac{57.3S}{R} = \frac{57.3(485)}{600} = 46.3175^\circ$

$$d = R \left( 1 - \cos \frac{\theta}{2} \right)$$

$$= 600(1 - \cos 23.15875^\circ)$$

$$\approx 48.3488$$

The distance is 48 ft. (rounded to two significant digits)

- (c) The faster the speed, the more land needs to be cleared on the inside of the curve.

## Chapter 2: Review Exercises

1.  $\sin A = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{60}{61}$

$$\cos A = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{11}{61}$$

$$\tan A = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{60}{11}$$

$$\cot A = \frac{\text{side adjacent}}{\text{side opposite}} = \frac{11}{60}$$

$$\sec A = \frac{\text{hypotenuse}}{\text{side adjacent}} = \frac{61}{11}$$

$$\csc A = \frac{\text{hypotenuse}}{\text{side opposite}} = \frac{61}{60}$$

2.  $\sin A = \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{40}{58} = \frac{20}{29}$

$$\cos A = \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{42}{58} = \frac{21}{29}$$

$$\tan A = \frac{\text{side opposite}}{\text{side adjacent}} = \frac{40}{42} = \frac{20}{21}$$

$$\cot A = \frac{\text{side adjacent}}{\text{side opposite}} = \frac{42}{40} = \frac{21}{20}$$

$$\sec A = \frac{\text{hypotenuse}}{\text{side adjacent}} = \frac{58}{42} = \frac{29}{21}$$

$$\csc A = \frac{\text{hypotenuse}}{\text{side opposite}} = \frac{58}{40} = \frac{29}{20}$$

3.  $\sin 4\beta = \cos 5\beta$

Sine and cosine are cofunctions, so the sum of the angles is  $90^\circ$ .

$$4\beta + 5\beta = 90^\circ \Rightarrow 9\beta = 90^\circ \Rightarrow \beta = 10^\circ$$

4.  $\sec(2\theta + 10^\circ) = \csc(4\theta + 20^\circ)$

Secant and cosecant are cofunctions, so the sum of the angles is  $90^\circ$ .

$$(2\theta + 10^\circ) + (4\theta + 20^\circ) = 90^\circ$$

$$6\theta + 30^\circ = 90^\circ$$

$$6\theta = 60^\circ \Rightarrow \theta = 10^\circ$$

5.  $\tan(5x + 11^\circ) = \cot(6x + 2^\circ)$

Tangent and cotangent are cofunctions, so the sum of the angles is  $90^\circ$ .

$$(5x + 11^\circ) + (6x + 2^\circ) = 90^\circ$$

$$11x + 13^\circ = 90^\circ$$

$$11x = 77^\circ \Rightarrow x = 7^\circ$$

6.  $\cos\left(\frac{3\theta}{5} + 11^\circ\right) = \sin\left(\frac{7\theta}{10} + 40^\circ\right)$

Sine and cosine are cofunctions, so the sum of the angles is  $90^\circ$ .

$$\left(\frac{3\theta}{5} + 11^\circ\right) + \left(\frac{7\theta}{10} + 40^\circ\right) = 90^\circ$$

$$\left(\frac{6\theta}{10} + 11^\circ\right) + \left(\frac{7\theta}{10} + 40^\circ\right) = 90^\circ$$

$$\frac{13\theta}{10} + 51^\circ = 90^\circ \Rightarrow \frac{13\theta}{10} = 39^\circ$$

$$\theta = 39^\circ \cdot \frac{10}{13} = 30^\circ$$

7.  $\sin 46^\circ < \sin 58^\circ$

In the interval from  $0^\circ$  to  $90^\circ$ , as the angle increases, so does the sine of the angle, so  $\sin 46^\circ < \sin 58^\circ$  is true.

8.  $\cos 47^\circ < \cos 58^\circ$

In the interval from  $0^\circ$  to  $90^\circ$ , as the angle increases, the cosine of the angle decreases, so  $\cos 47^\circ < \cos 58^\circ$  is false.

9.  $\sec 48^\circ \geq \cos 42^\circ$

Using the reciprocal identity,

$$\sec 48^\circ = \frac{1}{\cos 48^\circ}. \text{ Since } 48^\circ \text{ and } 42^\circ \text{ are in}$$

quadrant I,  $\sec 48^\circ$  and  $\cos 42^\circ$  are both positive. Thus

$$0 < \cos 48^\circ < 1 \Rightarrow \frac{1}{\cos 48^\circ} = \sec 48^\circ > 1. \text{ Since}$$

$$0 < \cos 42^\circ < 1, \sec 48^\circ \geq \cos 42^\circ \text{ is true.}$$

10.  $\sin 22^\circ \geq \csc 68^\circ$

Using the reciprocal identity,


$$\csc 68^\circ = \frac{1}{\sin 68^\circ}. \text{ Since } 68^\circ \text{ and } 22^\circ \text{ are in}$$

quadrant I,  $\csc 68^\circ$  and  $\sin 22^\circ$  are both positive. Thus

$$0 < \sin 68^\circ < 1 \Rightarrow \frac{1}{\sin 68^\circ} = \csc 68^\circ > 1. \text{ Since}$$

$$0 < \sin 22^\circ < 1, \sin 22^\circ \geq \csc 68^\circ \text{ is false.}$$

11. The sum of the measures of angles  $A$  and  $B$  is  $90^\circ$ , and, thus, they are complementary angles. Since sine and cosine are cofunctions, we have  $\sin B = \cos(90^\circ - B) = \cos A$ .



12. A  $120^\circ$  angle lies in quadrant II, so the reference angle is  $180^\circ - 120^\circ = 60^\circ$ . Since  $120^\circ$  is in quadrant II, the cosine, tangent, cotangent, and secant are negative.

$$\sin 120^\circ = \sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 120^\circ = -\cos 60^\circ = -\frac{1}{2}$$

$$\tan 120^\circ = -\tan 60^\circ = -\sqrt{3}$$

$$\cot 120^\circ = -\cot 60^\circ = -\frac{1}{\sqrt{3}} = -\frac{\sqrt{3}}{3}$$

$$\sec 120^\circ = -\sec 60^\circ = -2$$

$$\csc 120^\circ = \csc 60^\circ = \frac{2}{\sqrt{3}} = \frac{2\sqrt{3}}{3}$$

13.  $1020^\circ$  is coterminal with  $1020^\circ - 2 \cdot 360^\circ = 300^\circ$ . The reference angle is  $360^\circ - 300^\circ = 60^\circ$ . Because  $1020^\circ$  lies in quadrant IV, the sine, tangent, cotangent, and cosecant are negative.

$$\sin 1020^\circ = -\sin 60^\circ = -\frac{\sqrt{3}}{2}$$

$$\cos 1020^\circ = \cos 60^\circ = \frac{1}{2}$$

$$\tan 1020^\circ = -\tan 60^\circ = -\sqrt{3}$$

$$\cot 1020^\circ = -\cot 60^\circ = -\frac{\sqrt{3}}{3}$$

$$\sec 1020^\circ = \sec 60^\circ = 2$$

$$\csc 1020^\circ = -\csc 60^\circ = -\frac{2\sqrt{3}}{3}$$

14.  $-225^\circ$  is coterminal with  $-225^\circ + 360^\circ = 135^\circ$ . This angle lies in quadrant II. The reference angle is  $180^\circ - 135^\circ = 45^\circ$ . Since  $-225^\circ$  is in quadrant II, the cosine, tangent, cotangent, and secant are negative.

$$\sin(-225^\circ) = \sin 45^\circ = \frac{\sqrt{2}}{2}$$

$$\cos(-225^\circ) = -\cos 45^\circ = -\frac{\sqrt{2}}{2}$$

$$\tan(-225^\circ) = -\tan 45^\circ = -1$$

$$\cot(-225^\circ) = -\cot 45^\circ = -1$$

$$\sec(-225^\circ) = -\sec 45^\circ = -\sqrt{2}$$

$$\csc(-225^\circ) = \csc 45^\circ = \sqrt{2}$$

15.  $-1470^\circ$  is coterminal with  $-1470^\circ + 5 \cdot 360^\circ = 330^\circ$ . This angle lies in quadrant IV. The reference angle is  $360^\circ - 330^\circ = 30^\circ$ . Since  $-1470^\circ$  is in quadrant IV, the sine, tangent, cotangent, and cosecant are negative.

$$\sin(-1470^\circ) = -\sin 30^\circ = -\frac{1}{2}$$

$$\cos(-1470^\circ) = \cos 30^\circ = \frac{\sqrt{3}}{2}$$

$$\tan(-1470^\circ) = -\tan 30^\circ = -\frac{\sqrt{3}}{3}$$

$$\cot(-1470^\circ) = -\cot 30^\circ = -\sqrt{3}$$

$$\sec(-1470^\circ) = \sec 30^\circ = \frac{2\sqrt{3}}{3}$$

$$\csc(-1470^\circ) = -\csc 30^\circ = -2$$

$$16. \sin \theta = -\frac{1}{2}$$

Since  $\sin \theta$  is negative,  $\theta$  must lie in quadrants III or IV. The absolute value of  $\sin \theta$  is  $\frac{1}{2}$ , so the reference angle,  $\theta'$ , is  $30^\circ$ .

The angle in quadrant III will be  $180^\circ + \theta' = 180^\circ + 30^\circ = 210^\circ$ , and the quadrant IV angle is  $360^\circ - \theta' = 360^\circ - 30^\circ = 330^\circ$ .

$$17. \cos \theta = -\frac{1}{2}$$

Since  $\cos \theta$  is negative,  $\theta$  must lie in quadrants II or III. Since the absolute value of  $\cos \theta$  is  $\frac{1}{2}$ , the reference angle,  $\theta'$  must be  $60^\circ$ . The quadrant II angle  $\theta$  equals  $180^\circ - \theta' = 180^\circ - 60^\circ = 120^\circ$ , and the quadrant III angle  $\theta$  equals  $180^\circ + \theta' = 180^\circ + 60^\circ = 240^\circ$ .

$$18. \cot \theta = -1$$

Since  $\cot \theta$  is negative,  $\theta$  must lie in quadrants II or IV. Since the absolute value of  $\cot \theta$  is 1 the reference angle,  $\theta'$  must be  $45^\circ$ . The quadrant II angle  $\theta$  equals  $180^\circ - \theta' = 180^\circ - 45^\circ = 135^\circ$ , and the quadrant IV angle  $\theta$  equals  $360^\circ - \theta' = 360^\circ - 45^\circ = 315^\circ$ .

$$19. \sec \theta = -\frac{2\sqrt{3}}{3}$$


Since  $\sec \theta$  is negative,  $\theta$  must lie in quadrants II or III. Since the absolute value of  $\sec \theta$  is  $\frac{2\sqrt{3}}{3}$ , the reference angle,  $\theta'$  must be  $30^\circ$ . The quadrant II angle  $\theta$  equals  $180^\circ - \theta' = 180^\circ - 30^\circ = 150^\circ$ , and the quadrant III angle  $\theta$  equals  $180^\circ + \theta' = 180^\circ + 30^\circ = 210^\circ$ .

$$20. \cos 60^\circ + 2\sin^2 30^\circ = \frac{1}{2} + 2\left(\frac{1}{2}\right)^2 = \frac{1}{2} + \frac{1}{2} = 1$$

$$21. \tan^2 120^\circ - 2\cot 240^\circ = (-\sqrt{3})^2 - 2\left(\frac{\sqrt{3}}{3}\right) = 3 - \frac{2\sqrt{3}}{3}$$

$$\begin{aligned} 22. \sec^2 300^\circ - 2\cos^2 150^\circ + \tan 45^\circ &= 2^2 - 2\left(-\frac{\sqrt{3}}{2}\right)^2 + 1 \\ &= 4 - 2\left(\frac{3}{4}\right) + 1 = 4 - \frac{3}{2} + 1 = \frac{7}{2} \end{aligned}$$

$$23. \text{(a) } (-3, -3)$$



The distance from the origin is  $r$ :


$$\begin{aligned} r &= \sqrt{x^2 + y^2} \Rightarrow r = \sqrt{(-3)^2 + (-3)^2} \Rightarrow \\ r &= \sqrt{9 + 9} \Rightarrow r = \sqrt{18} \Rightarrow r = 3\sqrt{2} \end{aligned}$$

$$\sin \theta = \frac{y}{r} = \frac{-3}{3\sqrt{2}} = -\frac{1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = -\frac{\sqrt{2}}{2}$$

$$\cos \theta = \frac{x}{r} = \frac{-3}{3\sqrt{2}} = -\frac{1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = -\frac{\sqrt{2}}{2}$$

$$\tan \theta = \frac{y}{x} = \frac{-3}{-3} = 1$$

$$\text{(b) } (1, -\sqrt{3})$$



The distance from the origin is  $r$ :


$$\begin{aligned} r &= \sqrt{x^2 + y^2} = \sqrt{(1)^2 + (-\sqrt{3})^2} \\ &= \sqrt{1 + 3} = \sqrt{4} = 2 \end{aligned}$$

$$\sin \theta = \frac{y}{r} = \frac{-\sqrt{3}}{2} = -\frac{\sqrt{3}}{2}$$

$$\cos \theta = \frac{x}{r} = \frac{1}{2}$$


$$\tan \theta = \frac{y}{x} = \frac{-\sqrt{3}}{1} = -\sqrt{3}$$

For the exercises in this section, be sure your calculator is in degree mode. If your calculator accepts angles in degrees, minutes, and seconds, it is not necessary to change angles to decimal degrees. Keystroke sequences may vary on the type and/or model of calculator being used. Screens shown will be from a TI-83 Plus calculator. To obtain the degree ( $^\circ$ ) and ( $'$ ) symbols, go to the ANGLE menu (2nd APPS).




For Exercises 24–27, the calculation for decimal degrees is indicated for calculators that do not accept degree, minutes, and seconds.

24.  $\sin 72^\circ 30' \approx .95371695$




$$72^\circ 30' = \left(72 + \frac{30}{60}\right)^\circ = 72.5^\circ$$

25.  $\sec 222^\circ 30' \approx -1.3563417$




$$222^\circ 30' = \left(222 + \frac{30}{60}\right)^\circ = 222.5^\circ$$

26.  $\cot 305.6^\circ \approx -.71592968$




27.  $\csc 78^\circ 21' \approx 1.0210339$




$$78^\circ 21' = \left(78 + \frac{21}{60}\right)^\circ = 78.35^\circ$$

28.  $\sec 58.9041^\circ \approx 1.9362132$



29.  $\tan 11.7689^\circ \approx .20834446$



30. If  $\theta = 135^\circ$ ,  $\theta' = 180^\circ - 135^\circ = 45^\circ$ .


If  $\theta = 45^\circ$ ,  $\theta' = 45^\circ$ .

If  $\theta = 300^\circ$ ,  $\theta' = 360^\circ - 300^\circ = 60^\circ$ .

If  $\theta = 140^\circ$ ,  $\theta' = 180^\circ - 140^\circ = 40^\circ$ .


Of these reference angles,  $40^\circ$  is the only one which is not a special angle, so  $D, \tan 140^\circ$ , is the only one which cannot be determined exactly using the methods of this chapter.

31.  $\sin \theta = .8254121$




$$\theta \approx 55.673870^\circ$$

32.  $\cot \theta = 1.1249386$



$$\theta \approx 41.635092^\circ$$

33.  $\cos \theta = .97540415$



$$\theta \approx 12.733938^\circ$$

34.  $\sec \theta = 1.2637891$

```
1/1.2637891
.7912712651
cos-1(Ans)
37.69552761
■
```

$\theta \approx 37.695528^\circ$

35.  $\tan \theta = 1.9633124$

```
tan-1(1.9633124)
63.00828638
```

$\theta \approx 63.008286^\circ$

36.  $\csc \theta = 9.5670466$

```
1/9.5670466
.1045254656
sin-1(Ans)
5.999827299
■
```

$\theta \approx 5.9998273^\circ$

37. Since the value of  $\sin \theta$  is positive in quadrants I and II, the two angles in  $[0^\circ, 360^\circ)$  are approximately  $47.1^\circ$  and  $180^\circ - 47.1^\circ = 132.9^\circ$ .

```
sin-1(.73254290)
47.1000001
■
```

38. Since the value of  $\tan \theta$  is positive in quadrants I and III, the two angles in  $[0^\circ, 360^\circ)$  are approximately  $54.2^\circ$  and  $180^\circ - 54.2^\circ = 234.2^\circ$

```
tan-1(1.3865342)
54.2000001
```

39.  $\sin 50^\circ + \sin 40^\circ = \sin 90^\circ$

```
sin(50)+sin(40)
1.408832053
sin(90)
1
```

Using a calculator gives

$\sin 50^\circ + \sin 40^\circ = 1.408832053$  while  $\sin 90^\circ = 1$ . Thus, the statement is false.

40.  $\cos 210^\circ = \cos 180^\circ \cdot \cos 30^\circ - \sin 180^\circ \cdot \sin 30^\circ$   
 $\cos 210^\circ = -\cos 30^\circ = -\frac{\sqrt{3}}{2}$  and  
 $\cos 180^\circ \cdot \cos 30^\circ - \sin 180^\circ \cdot \sin 30^\circ$   
 $= (-1)\left(\frac{\sqrt{3}}{2}\right) - 0\left(\frac{1}{2}\right) = -\frac{\sqrt{3}}{2}$

Thus, the statement is true.

41.  $\sin 240^\circ = 2 \sin 120^\circ \cos 120^\circ$   
 $\sin 240^\circ = -\sin 60^\circ = -\frac{\sqrt{3}}{2}$  and  
 $2 \sin 120^\circ \cos 120^\circ = 2 \sin 60^\circ (-\cos 60^\circ)$   
 $= 2\left(\frac{\sqrt{3}}{2}\right)\left(-\frac{1}{2}\right) = -\frac{\sqrt{3}}{2}$

Thus, the statement is true.

42.  $\sin 42^\circ + \sin 42^\circ = \sin 84^\circ$

```
sin(42)+sin(42)
1.338261213
sin(84)
.9945218954
```

Using a calculator gives

$\sin 42^\circ + \sin 42^\circ = 1.338261213$  while  $\sin 84^\circ = .9945218954$ . Thus, the statement is false.

43. No,  $\cot 25^\circ = \frac{1}{\tan 25^\circ} \neq \tan^{-1} 25^\circ$ .

44. Since  $\sin 2976^\circ$  is positive and  $\cos 2976^\circ$  is negative,  $2976^\circ$  must lie in quadrant II.

```
sin(2976)
.9945218954
cos(2976)
-.1045284633
```

45. Since  $\cos 1997^\circ$  and  $\sin 1997^\circ$  are both negative,  $1997^\circ$  must lie in quadrant III.


```
cos(1997)
-.956304756
sin(1997)
-.2923717047
```

46. Since  $\sin 4000^\circ$  and  $\cos 4000^\circ$  are both positive,  $4000^\circ$  must lie in quadrant I.

```

sin(4000)
.6427876097
cos(4000)
.7660444431
  
```

47.  $A = 58^\circ 30'$ ,  $c = 748$   
 $A + B = 90^\circ \Rightarrow B = 90^\circ - A \Rightarrow$   
 $B = 90^\circ - 58^\circ 30' = 89^\circ 60' - 58^\circ 30'$   
 $= 31^\circ 30'$   
 $\sin A = \frac{a}{c} \Rightarrow \sin 58^\circ 30' = \frac{a}{748} \Rightarrow$   
 $a = 748 \sin 58^\circ 30' \approx 638$  (rounded to three significant digits)  
 $\cos A = \frac{b}{c} \Rightarrow \cos 58^\circ 30' = \frac{b}{748} \Rightarrow$   
 $b = 748 \cos 58^\circ 30' \approx 391$  (rounded to three significant digits)
48.  $a = 129.70$ ,  $b = 368.10$   
 $c = \sqrt{a^2 + b^2} \Rightarrow c = \sqrt{129.70^2 + 368.10^2}$   
 $\approx 390.28$  (rounded to five significant digits)  
 $\tan A = \frac{a}{b} = \frac{129.70}{368.10} \Rightarrow$   
 $A = \tan^{-1}\left(\frac{129.70}{368.10}\right) \approx 19.41^\circ$   
 $\approx 19^\circ + (.41 \cdot 60)' \approx 19^\circ 25'$   
 $\tan B = \frac{b}{a} = \frac{368.10}{129.70} \Rightarrow$   
 $B = \tan^{-1}\left(\frac{368.10}{129.70}\right) \approx 70.59^\circ$   
 $\approx 70^\circ + (.59 \cdot 60)' \approx 70^\circ 35'$
49.  $A = 39.72^\circ$ ,  $b = 38.97$  m




- $A + B = 90^\circ \Rightarrow B = 90^\circ - A \Rightarrow$   
 $B = 90^\circ - 39.72^\circ = 50.28^\circ$   
 $\tan A = \frac{a}{b} \Rightarrow \tan 39.72^\circ = \frac{a}{38.97} \Rightarrow$   
 $a = 38.97 \tan 39.72^\circ \approx 32.38$  m (rounded to four significant digits)

$$c \cos A = \frac{b}{c} \Rightarrow \cos 39.72^\circ = \frac{38.97}{c} \Rightarrow$$

$$c \cos 39.72^\circ = 38.97 \Rightarrow$$


$$c = \frac{38.97}{\cos 39.72^\circ} \approx 50.66$$
 m  
 (rounded to five significant digits)

50.  $B = 47^\circ 53'$ ,  $b = 298.6$  m



$A + B = 90^\circ \Rightarrow A = 90^\circ - B \Rightarrow$   
 $A = 90^\circ - 47^\circ 53' = 89^\circ 60' - 47^\circ 53'$   
 $= 42^\circ 7'$   
 $\tan B = \frac{b}{a} \Rightarrow \tan 47^\circ 53' = \frac{298.6}{a} \Rightarrow$   
 $a \tan 47^\circ 53' = 298.6 \Rightarrow$   
 $a = \frac{298.6}{\tan 47^\circ 53'} \approx 270.0$  m  
 (rounded to four significant digits)  
 $\sin B = \frac{b}{c} \Rightarrow \sin 47^\circ 53' = \frac{298.6}{c} \Rightarrow$   
 $c \sin 47^\circ 53' = 298.6 \Rightarrow$   
 $c = \frac{298.6}{\sin 47^\circ 53'} \approx 402.5$  m  
 (rounded to four significant digits)

51. Let  $x$  = the height of the tree.




$$\tan 70^\circ = \frac{x}{50} \Rightarrow x = 50 \tan 70^\circ \approx 137$$
 ft

52.  $r = \sqrt{12^2 + 5^2} = \sqrt{144 + 25} = \sqrt{169} = 13$   
 $\tan \theta = \frac{5}{12} \Rightarrow \theta = \tan^{-1}\left(\frac{5}{12}\right) \approx 23^\circ$




53. Let  $x$  = height of the tower.



$$\begin{aligned} \tan 38^\circ 20' &= \frac{x}{93.2} \\ x &= 93.2 \tan 38^\circ 20' \\ x &\approx 73.6930 \end{aligned}$$

The height of the tower is 73.7 ft. (rounded to three significant digits)


54. Let  $h$  = height of the tower.



$$\begin{aligned} \tan 29.5^\circ &= \frac{h}{36.0} \\ h &= 36.0 \tan 29.5^\circ \\ h &\approx 20.3678 \end{aligned}$$

The height of the tower is 20.4 m. (rounded to three significant digits)

55. Let  $x$  = length of the diagonal




$$\begin{aligned} \cos 35.65^\circ &= \frac{15.24}{x} \\ x &= \frac{15.24}{\cos 35.65^\circ} \approx 18.7548 \end{aligned}$$

The length of the diagonal is 18.75 cm (rounded to three significant digits).

56. Let  $x$  = the length of the equal sides of an isosceles triangle.

Divide the isosceles triangle into two congruent right triangles




$$d = \frac{1}{2}(49.28) = 24.64 \text{ and}$$

$$\theta = \frac{1}{2}(58.746^\circ) = 29.373^\circ$$

$$\begin{aligned} \sin \theta &= \frac{d}{s} \Rightarrow \sin 29.373^\circ = \frac{24.64}{s} \Rightarrow \\ s &= \frac{24.64}{\sin 29.373^\circ} \approx 50.2352 \end{aligned}$$

Each side is 50.24 m long (rounded to 4 significant digits).

57. Draw triangle  $ABC$  and extend the north-south lines to a point  $X$  south of  $A$  and  $S$  to a point  $Y$ , north of  $C$ .



Angle  $ACB = 344^\circ - 254^\circ = 90^\circ$ , so  $ABC$  is a right triangle.

Angle  $BAX = 32^\circ$  since it is an alternate interior angle to  $32^\circ$ .

Angle  $YCA = 360^\circ - 344^\circ = 16^\circ$

Angle  $XAC = 16^\circ$  since it is an alternate interior angle to angle  $YCA$ .

Angle  $BAC = 32^\circ + 16^\circ = 48^\circ$ .


In triangle  $ABC$ ,

$$\cos A = \frac{AC}{AB} \Rightarrow \cos 48^\circ = \frac{780}{AB} \Rightarrow$$

$$AB \cos 48^\circ = 780 \Rightarrow AB = \frac{780}{\cos 48^\circ} \approx 1165.6917$$

The distance from  $A$  to  $B$  is 1200 m. (rounded to two significant digits)

58. Draw triangle  $ABC$  and extend north-south lines from points  $A$  and  $B$ . Angle  $ABX$  is  $55^\circ$  (alternate interior angles of parallel lines cut by a transversal have the same measure) so Angle  $ABC$  is  $55^\circ + 35^\circ = 90^\circ$ .




Since angle  $ABC$  is a right angle, use the Pythagorean theorem to find the distance from  $A$  to  $C$ .

$$(AC)^2 = 80^2 + 74^2 \Rightarrow (AC)^2 = 6400 + 5476 \Rightarrow$$

$$(AC)^2 = 11,876 \Rightarrow AC = \sqrt{11,876} \approx 108.9771$$

It is 110 km from  $A$  to  $C$ . (rounded to two significant digits)

59. Suppose  $A$  is the car heading south at 55 mph,  $B$  is the car heading west, and point  $C$  is the intersection from which they start. After two hours, using  $d = rt$ ,  $AC = 55(2) = 110$ . Angle  $ACB$  is a right angle, so triangle  $ACB$  is a right triangle. The bearing of  $A$  from  $B$  is  $324^\circ$ , so angle  $CAB = 360^\circ - 324^\circ = 36^\circ$ .




$$\cos \angle CAB = \frac{AC}{AB} \Rightarrow \cos 36^\circ = \frac{110}{AB} \Rightarrow$$

$$AB = \frac{110}{\cos 36^\circ} \approx 135.9675$$

The distance from  $A$  to  $B$  is about 140 mi (rounded to two significant digits).

60. Let  $x$  = the leg opposite angle  $A$



$$\tan A = \frac{x}{k} \Rightarrow x = k \tan A \text{ and}$$

$$\tan B = \frac{h+x}{k} \Rightarrow x = k \tan B - h. \text{ So,}$$

$$k \tan A = k \tan B - h$$

$$h = k \tan B - k \tan A = k(\tan B - \tan A)$$

- 61.–62. Answers will vary.

$$63. h = R \left( \frac{1}{\cos\left(\frac{180T}{P}\right)} - 1 \right)$$

- (a) Let  $R = 3955$  mi,  $T = 25$  min,  $P = 140$  min.

$$h = R \left( \frac{1}{\cos\left(\frac{180T}{P}\right)} - 1 \right)$$

$$h = 3955 \left( \frac{1}{\cos\left(\frac{180 \cdot 25}{140}\right)} - 1 \right) \approx 715.9424$$

The height of the satellite is approximately 716 mi.

- (b) Let  $R = 3955$  mi,  $T = 30$  min,  $P = 140$  min.

$$h = R \left( \frac{1}{\cos\left(\frac{180T}{P}\right)} - 1 \right)$$

$$h = 3955 \left( \frac{1}{\cos\left(\frac{180 \cdot 30}{140}\right)} - 1 \right) \approx 1103.6349$$

The height of the satellite is approximately 1104 mi.

64. (a) From the figure, we see that

$$\sin \theta = \frac{x_Q - x_P}{d} \Rightarrow x_Q = x_P + d \sin \theta.$$

Similarly, we have

$$\cos \theta = \frac{y_Q - y_P}{d} \Rightarrow y_Q = y_P + d \cos \theta.$$

- (b) Let  $(x_P, y_P) = (123.62, 337.95)$ ,

$$\theta = 17^\circ 19' 22'', \text{ and } d = 193.86 \text{ ft.}$$

$$x_Q = x_P + d \sin \theta \Rightarrow$$

$$x_Q = 123.62 + 193.86 \sin 17^\circ 19' 22'' \approx 181.3427$$

$$y_Q = y_P + d \cos \theta \Rightarrow$$


$$y_Q = 337.95 + 193.86 \cos 17^\circ 19' 22'' \approx 523.0170$$

Thus, the coordinates of  $Q$  are  $(181.34, 523.02)$ , rounded to five significant digits.

## Chapter 2: Chapter Test

$$\begin{aligned}
 1. \quad \sin A &= \frac{\text{side opposite}}{\text{hypotenuse}} = \frac{12}{13} \\
 \cos A &= \frac{\text{side adjacent}}{\text{hypotenuse}} = \frac{5}{13} \\
 \tan A &= \frac{\text{side opposite}}{\text{side adjacent}} = \frac{12}{5} \\
 \cot A &= \frac{\text{side adjacent}}{\text{side opposite}} = \frac{5}{12} \\
 \sec A &= \frac{\text{hypotenuse}}{\text{side adjacent}} = \frac{13}{5} \\
 \csc A &= \frac{\text{hypotenuse}}{\text{side opposite}} = \frac{13}{12}
 \end{aligned}$$

2. Apply the relationships between the lengths of the sides of a  $30^\circ - 60^\circ$  right triangle first to the triangle on the right to find the values of  $y$  and  $w$ . In the  $30^\circ - 60^\circ$  right triangle, the side opposite the  $60^\circ$  angle is  $\sqrt{3}$  times as long as the side opposite to the  $30^\circ$  angle. The length of the hypotenuse is 2 times as long as the shorter leg (opposite the  $30^\circ$  angle).



Thus, we have  $y = 4\sqrt{3}$  and  $w = 2(4) = 8$ .

Apply the relationships between the lengths of the sides of a  $45^\circ - 45^\circ$  right triangle next to the triangle on the left to find the values of  $x$  and  $z$ . In the  $45^\circ - 45^\circ$  right triangle, the sides opposite the  $45^\circ$  angles measure the same.

The hypotenuse is  $\sqrt{2}$  times the measure of a leg. Thus, we have  $x = 4$  and  $z = 4\sqrt{2}$ .

$$3. \quad \sin(B + 15^\circ) = \cos(2B + 30^\circ)$$

Since sine and cosine are cofunctions, the sum of the angles is  $90^\circ$ . So,

$$(B + 15^\circ) + (2B + 30^\circ) = 90^\circ$$

$$3B + 45^\circ = 90^\circ$$

$$3B = 45^\circ \Rightarrow B = 15^\circ$$

$$4. \quad \text{(a)} \quad \sin 24^\circ < \sin 48^\circ$$

In the interval from  $0^\circ$  to  $90^\circ$ , as the angle increases, so does the sine of the angle, so  $\sin 24^\circ < \sin 48^\circ$  is true.

$$\text{(b)} \quad \cos 24^\circ < \cos 48^\circ$$

In the interval from  $0^\circ$  to  $90^\circ$ , as the angle increases, so the cosine of the angle decreases, so  $\cos 24^\circ < \cos 48^\circ$  is false.

$$\begin{aligned}
 \text{(c)} \quad \cos(60^\circ + 30^\circ) &= \cos 60^\circ \cos 30^\circ - \sin 60^\circ \sin 30^\circ \\
 \cos(60^\circ + 30^\circ) &= \cos 90^\circ = 0 \\
 \cos 60^\circ \cos 30^\circ - \sin 60^\circ \sin 30^\circ &= \frac{1}{2} \left( \frac{\sqrt{3}}{2} \right) - \frac{\sqrt{3}}{2} \left( \frac{1}{2} \right) = 0
 \end{aligned}$$

Thus, the statement is true.

5. A  $240^\circ$  angle lies in quadrant III, so the reference angle is  $240^\circ - 180^\circ = 60^\circ$ . Since  $240^\circ$  is in quadrant III, the sine, cosine, secant, and cosecant are negative.

$$\sin 240^\circ = -\sin 60^\circ = -\frac{\sqrt{3}}{2}$$

$$\cos 240^\circ = -\cos 60^\circ = -\frac{1}{2}$$

$$\tan 240^\circ = \tan 60^\circ = \sqrt{3}$$

$$\cot 240^\circ = \cot 60^\circ = \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3}$$

$$\sec 240^\circ = -\sec 60^\circ = -2$$

$$\csc 240^\circ = -\csc 60^\circ = -\frac{2}{\sqrt{3}} = -\frac{2\sqrt{3}}{3}$$

6.  $-135^\circ$  is coterminal with  $-135^\circ + 360^\circ = 225^\circ$ . This angle lies in quadrant III. The reference angle is  $225^\circ - 180^\circ = 45^\circ$ . Since  $-135^\circ$  is in quadrant III, the sine, cosine, secant, and cosecant are negative.

$$\sin(-135^\circ) = -\sin 45^\circ = -\frac{\sqrt{2}}{2}$$

$$\cos(-135^\circ) = -\cos 45^\circ = -\frac{\sqrt{2}}{2}$$

$$\tan(-135^\circ) = \tan 45^\circ = 1$$

$$\cot(-135^\circ) = \cot 45^\circ = 1$$

$$\sec(-135^\circ) = -\sec 45^\circ = -\sqrt{2}$$

$$\csc(-135^\circ) = -\csc 45^\circ = -\sqrt{2}$$

7.  $990^\circ$  is coterminal with  $990^\circ - 2 \cdot 360^\circ = 270^\circ$ , which is the reference angle.

$$\sin 990^\circ = \sin 270^\circ = -1$$

$$\cos 990^\circ = \cos 270^\circ = 0$$

$$\tan 990^\circ = \tan 270^\circ \text{ undefined}$$

$$\cot 990^\circ = \cot 270^\circ = 0$$

$$\sec 990^\circ = \sec 270^\circ \text{ undefined}$$

$$\csc 990^\circ = \csc 270^\circ = -1$$

8.  $\cos \theta = -\frac{\sqrt{2}}{2}$

Since  $\cos \theta$  is negative,  $\theta$  must lie in quadrant II or quadrant III. The absolute value of  $\cos \theta$  is  $\frac{\sqrt{2}}{2}$ , so  $\theta' = 45^\circ$ . The quadrant II angle  $\theta$  equals  $180^\circ - \theta' = 180^\circ - 45^\circ = 135^\circ$ , and the quadrant III angle  $\theta$  equals  $180^\circ + \theta' = 180^\circ + 45^\circ = 225^\circ$ .


9.  $\csc \theta = -\frac{2\sqrt{3}}{3}$

Since  $\csc \theta$  is negative,  $\theta$  must lie in quadrant III or quadrant IV. The absolute value of  $\csc \theta$  is  $\frac{2\sqrt{3}}{3}$ , so  $\theta' = 60^\circ$ . The quadrant III angle  $\theta$  equals  $180^\circ + \theta' = 180^\circ + 60^\circ = 240^\circ$ , and the quadrant IV angle  $\theta$  equals  $360^\circ - \theta' = 360^\circ - 60^\circ = 300^\circ$ .


10.  $\tan \theta = 1 \Rightarrow \theta = 45^\circ$  or  $\theta = 225^\circ$

11.  $\tan \theta = 1.6778490$

Since  $\cot \theta = \frac{1}{\tan \theta} = (\tan \theta)^{-1}$ , we can use division or the inverse key (multiplicative inverse).




12. (a)  $\sin 78^\circ 21' \approx .97939940$




$78^\circ 21' = \left(78 + \frac{21}{60}\right)^\circ = 78.35^\circ$


(b)  $\tan 117.689^\circ \approx -1.9056082$



(c)  $\sec 58.9041^\circ \approx 1.9362432$




13.  $\sin \theta = .27843196$



$\theta \approx 16.16664145^\circ$

14.  $A = 58^\circ 30', a = 748$



$A + B = 90^\circ \Rightarrow B = 90^\circ - A \Rightarrow$

$B = 90^\circ - 58^\circ 30' = 31^\circ 30'$


$\tan A = \frac{a}{b} \Rightarrow \tan 58^\circ 30' = \frac{748}{b} \Rightarrow$

$b = \frac{748}{\tan 58^\circ 30'} \approx 458$  (rounded to three significant digits)

$\sin A = \frac{a}{c} \Rightarrow \sin 58^\circ 30' = \frac{748}{c} \Rightarrow$

$c = \frac{748}{\sin 58^\circ 30'} \approx 877$  (rounded to three significant digits)


15. Let  $\theta$  = the measure of the angle that the guy wire makes with the ground.



$\sin \theta = \frac{71.3}{77.4}$

$\theta = \sin^{-1}\left(\frac{71.3}{77.4}\right) \approx 67.1^\circ \approx 67^\circ 10'$

16. Let  $x$  = the height of the flagpole.




$$\tan 32^\circ 10' = \frac{x}{24.7}$$

$$x = 24.7 \tan 32^\circ 10' \approx 15.5344$$

The flagpole is approximately 15.5 ft high.  
(rounded to three significant digits)


17. Let  $h$  = the height of the top of mountain above the cabin. Then  $2000 + h$  = the height of the mountain.



$$\tan 26^\circ = \frac{h}{14,000} \Rightarrow h \approx 6800 \text{ (rounded to two}$$

significant digits). Thus, the height of the mountain is about  $6800 + 2000 = 8800$  ft.


18. Let  $x$  = distance the ships are apart.  
In the figure, the measure of angle  $CAB$  is  $122^\circ - 32^\circ = 90^\circ$ . Therefore, triangle  $CAB$  is a right triangle.



Since  $d = rt$ , the distance traveled by the first ship in 2.5 hr is  
 $(2.5 \text{ hr})(16 \text{ knots}) = 40$  nautical mi and the second ship is  
 $(2.5 \text{ hr})(24 \text{ knots}) = 60$  nautical mi.  
 Applying the Pythagorean theorem, we have  
 $x^2 = 40^2 + 60^2 \Rightarrow x^2 = 1600 + 3600 \Rightarrow$   
 $x^2 = 5200 \Rightarrow x = \sqrt{5200} \approx 72.111$

The ships are 72 nautical mi apart. (rounded to 2 significant digits)

19. Draw triangle  $ACB$  and extend north-south lines from points  $A$  and  $C$ . Angle  $ACD$  is  $62^\circ$  (alternate interior angles of parallel lines cut by a transversal have the same measure), so Angle  $ACB$  is  $62^\circ + 28^\circ = 90^\circ$ .




Since angle  $ACB$  is a right angle, use the Pythagorean theorem to find the distance from  $A$  to  $B$ .

$$(AB)^2 = 75^2 + 53^2 \Rightarrow (AB)^2 = 5625 + 2809 \Rightarrow$$

$$(AB)^2 = 8434 \Rightarrow AB = \sqrt{8434} \approx 91.8368$$

It is 92 km from the pier to the boat, rounded to two significant digits.

20. Let  $x$  = the side adjacent to  $52.5^\circ$  in the smaller triangle.



In the larger triangle, we have

$$\tan 41.2^\circ = \frac{h}{168 + x} \Rightarrow h = (168 + x) \tan 41.2^\circ.$$

In the smaller triangle, we have

$$\tan 52.5^\circ = \frac{h}{x} \Rightarrow h = x \tan 52.5^\circ.$$

Substitute for  $h$  in this equation to solve for  $x$ .

$$(168 + x) \tan 41.2^\circ = x \tan 52.5^\circ$$

$$168 \tan 41.2^\circ + x \tan 41.2^\circ = x \tan 52.5^\circ$$

$$168 \tan 41.2^\circ = x \tan 52.5^\circ - x \tan 41.2^\circ$$

$$168 \tan 41.2^\circ = x (\tan 52.5^\circ - \tan 41.2^\circ)$$

$$\frac{168 \tan 41.2^\circ}{\tan 52.5^\circ - \tan 41.2^\circ} = x$$

Substituting for  $x$  in the equation for the smaller triangle gives

$$h = x \tan 52.5^\circ$$

$$h = \frac{168 \tan 41.2^\circ \tan 52.5^\circ}{\tan 52.5^\circ - \tan 41.2^\circ} \approx 448.0432$$

The height of the triangle is approximately 448 m. (rounded to three significant digits)

**Chapter 2: Quantitative Reasoning**

$$D = \frac{v^2 \sin \theta \cos \theta + v \cos \theta \sqrt{(v \sin \theta)^2 + 64h}}{32}$$

All answers are rounded to four significant digits.

1. Since  $v = 44$  ft per sec and  $h = 7$  ft, we have  $D = \frac{44^2 \sin \theta \cos \theta + 44 \cos \theta \sqrt{(44 \sin \theta)^2 + 64 \cdot 7}}{32}$

If  $\theta = 40^\circ$ ,  $D = \frac{1936 \sin 40 \cos 40 + 44 \cos 40 \sqrt{(44 \sin 40)^2 + 448}}{32} \approx 67.00$  ft.

If  $\theta = 42^\circ$ ,  $D = \frac{1936 \sin 42 \cos 42 + 44 \cos 42 \sqrt{(44 \sin 42)^2 + 448}}{32} \approx 67.14$  ft

If  $\theta = 45^\circ$ ,  $D = \frac{1936 \sin 45 \cos 45 + 44 \cos 45 \sqrt{(44 \sin 45)^2 + 448}}{32} \approx 66.84$  ft

As  $\theta$  increases,  $D$  increases and then decreases.

2. Since  $h = 7$  ft and  $\theta = 42^\circ$ , we have  $D = \frac{v^2 \sin 42 \cos 42 + v \cos 42 \sqrt{(v \sin 42)^2 + 64h}}{32}$

If  $v = 43$ ,  $D = \frac{43^2 \sin 42 \cos 42 + 43 \cos 42 \sqrt{(43 \sin 42)^2 + 448}}{32} \approx 64.40$  ft

If  $v = 44$ ,  $D = \frac{44^2 \sin 42 \cos 42 + 44 \cos 42 \sqrt{(44 \sin 42)^2 + 448}}{32} \approx 67.14$  ft

If  $v = 45$ ,  $D = \frac{45^2 \sin 42 \cos 42 + 45 \cos 42 \sqrt{(45 \sin 42)^2 + 448}}{32} \approx 69.93$  ft

As  $v$  increases,  $D$  increases.

3. The velocity affects the distance more. The shot-putter should concentrate on achieving as large a value of  $v$  as possible.