
# TEST BANK



# MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Find the slope of the line passing through the given pair of points.

1) (1, 2) and (7, 7)

A)  $\frac{9}{8}$ 

B)  $\frac{5}{6}$  C)  $\frac{5}{6}$  D)  $\frac{6}{5}$ 

1) \_\_\_\_\_

2) (2, 4) and (-8, 3)

D) 10

2) \_\_\_\_\_

3) \_\_\_\_\_

4) \_\_\_\_\_

3) (-6, -5) and (-6, 7)

B) Not defined

C) 0

D) 1

4) (-5, 9) and (6, 9)

A) Not defined

C) 18

B) 0

D) 18 11

5) (18, 9) and (16, -2) A)  $\frac{7}{34}$  B)  $\frac{2}{11}$ 

C)  $\frac{11}{2}$ 

D)  $\frac{11}{2}$ 

5) \_\_\_\_\_

6) \_\_\_\_\_

Find the slope of the line.

A) 0

B)  $\frac{3}{4}$ 

C)  $\frac{4}{3}$ 

D) 1

7) y = 4x - 6

A) 0

B) 1

C) -4

D) 4

8) 5x + 2y = -27A)  $\frac{27}{2}$ 

C)  $\frac{5}{2}$ 

8) \_\_\_\_\_

9) 4x - 3y = 27A)  $\frac{3}{4}$ 

C) - 9

D)  $\frac{4}{3}$ 

9) \_\_\_\_\_

10) \_\_\_\_\_

11) \_\_\_\_\_

10) The x-axis

A) 0

C) 1

B) -1

D) Not defined

11) x = -10

A) 1

C) 0

B) -10

D) Not defined

12) A line parallel to 
$$4y - 5x = 8$$

D)  $\frac{5}{4}$ 

12) \_\_\_\_\_

13) \_\_\_\_\_

14) \_\_\_\_\_

15) \_\_\_\_\_

16) \_\_\_\_\_

17) \_\_\_\_\_

18) \_\_\_\_\_

19) \_\_\_\_\_

20) \_\_\_\_\_

21) \_\_\_\_\_

13) A line parallel to 
$$-2x = 7y + 11$$
  
A)  $\frac{11}{2}$  B)  $\frac{2}{7}$ 

14) A line perpendicular to 
$$2x + 7y = -36$$
  
A)  $\frac{2}{7}$  B) 2 C)  $\frac{7}{2}$ 

15) A line perpendicular to 
$$5x = 2y + 3$$
  
A)  $\frac{2}{5}$  B)  $\frac{3}{2}$  C)  $\frac{5}{2}$ 

Find an equation in slope-intercept form (where possible) for the line.

16) Through 
$$(0, 1)$$
,  $m = \frac{1}{2}$ 

 $y = -\frac{1}{2}x + 1$   $y = -\frac{1}{2}x - 1$ 

B) 
$$\frac{1}{2}$$

$$y = \frac{1}{2}x + 1$$

17) Through 
$$(11, 5)$$
,  $m = 2$ 

A) y = -2x + 17

C) y = 2x - 17

B) 
$$y = 2x + 5$$

D) 
$$y = -2x + 1$$

18) Through 
$$(5, 0)$$
,  $m = -1$ 

A) y = 5x

C) 
$$y = x - 5$$

B) 
$$y = -5x$$

D) 
$$y = -x + 5$$

### 19) Through (3, 2), m = 3

A) y = -3x + 11

C) 
$$y = -3x - 7$$

B) 
$$y = 3x - 7$$

D) 
$$y = 3x + 11$$

#### 20) Through (-9, 8), m = 0

A) y = 8

D) 
$$x = -9$$

## 21) Through (-8, -5), with undefined slope

A)  $\frac{5}{5}$ -  $\frac{5}{8}$ -  $\frac{5}{8}$ -  $\frac{5}{8}$  -  $\frac{5}{8}$  = 0

D) 
$$x = -8$$

22) Through (2, 3), m = -

A) 
$$\frac{2}{5} \times \frac{4}{5}$$
  
C)  $\frac{2}{5} \times \frac{19}{5}$ 

b) 
$$\frac{2}{5}$$
  $\frac{19}{5}$   
D)  $\frac{2}{5}$   $\frac{4}{5}$ 

23)

Through (0, -2), m = 
$$\frac{5}{2}$$

A)  $\frac{5}{2}$ 
 $y = x + 2$ 

C)  $\frac{5}{2}$ 
 $y = x - 2$ 

B) 
$$y = -\frac{5}{2}x + 2$$
D) 
$$\frac{5}{2}$$

$$y = -\frac{5}{2}x - 2$$

24) 
$$\begin{array}{ccc} & \frac{5}{8} \\ & \text{Through (5, 3), m = -} \\ & A) & \frac{5}{8} & \frac{49}{8} \\ & y = -x + \\ & C) & \frac{5}{8} & \frac{25}{8} \\ & y = x + \end{array}$$

B) 
$$\frac{5}{8} \times \frac{49}{8}$$
 $y = x - 25$ 
 $y = x + 25$ 

26) Through (3, -2) and (-1, -13)  
A) 
$$\frac{11}{4}$$
  $\frac{10}{11}$   
 $y = x - C$   
C)  $\frac{11}{4}$   $\frac{41}{4}$   
 $y = x - C$ 

B) 
$$\frac{11}{4}$$
  $\frac{25}{4}$   $y = -\frac{x}{11}$   $\frac{34}{11}$   $y = -\frac{x}{11}$ 

27) Through (-7, -2) and (0, 3)  
A) 
$$\frac{5}{3}x + 3$$
  
C)  $\frac{5}{7}y = -x + 3$ 

B) 
$$\frac{5}{3}x + 3$$
D)  $\frac{5}{7}$ 
 $y = x + 3$ 

28) Through (-9, 0) and (8, 2)  
A) 
$$\frac{2}{17}$$
  $\frac{18}{17}$   
 $y = x + 14$ 

B) 
$$\frac{3}{2}$$
  
 $y = -\frac{3}{2}$   
 $y = -\frac{18}{17}$   
 $y = -\frac{18}{17}$ 

29) Through (0, -8) and (-6, 3)  
A) 
$$\frac{11}{6}$$
  
 $y = x - 8$   
C)  $\frac{8}{9}$   $\frac{7}{3}$   
 $y = x - 8$ 

B) 
$$\frac{11}{6}$$
  
 $y = -\frac{x}{8} \cdot \frac{7}{3}$   
 $y = -\frac{x}{3} \cdot \frac{7}{3}$ 

B) 
$$\frac{9}{17}$$
  $\frac{233}{17}$ 

$$y = x + 1$$
D)  $\frac{3}{5} + \frac{29}{5}$ 

A) 
$$y = 0.4x - 4.3$$
  
C)  $y = -0.4x + 1.3$ 

B) 
$$y = 2.5x - 19$$

D) 
$$y = -2.5x + 16$$

A) 
$$\frac{7}{2}$$
  
- x - 2y = 0  
C) x = 4

B) 
$$y = -2$$

D) 
$$\frac{2}{7}$$

A) 
$$\frac{1}{6}$$
 -  $x - 6y = 0$ 

B) 
$$x = -1$$

C) 
$$-6x - 1y = 0$$

D) 
$$y = -3$$

A) 
$$y = \frac{1}{4}$$
C) 
$$\frac{1}{4}$$

B) 
$$y = -4x - 8$$

D) 
$$y = 4x - 8$$

35) Through (-9, -1), perpendicular to 
$$6x + 5y = -59$$

A) 
$$\frac{5}{6} \times \frac{13}{2}$$
  
 $y = \frac{6}{5} \times + \frac{13}{2}$   
C)  $\frac{6}{5} \times -39$ 

B) 
$$\frac{5}{6}$$
  $\frac{13}{2}$ 

D) 
$$\frac{5}{6}$$
  $y = x$ 

36) Through (7, -7), parallel to 
$$5x - 2y = 51$$

A) 
$$\frac{7}{2}$$
  $\frac{5}{2}$   
 $y = x - \frac{5}{2}$   
C)  $\frac{2}{5}$   $\frac{7}{5}$ 

B) 
$$\frac{5}{2}$$
  $\frac{49}{2}$ 

$$y = -\frac{5}{2}x + \frac{49}{2}$$

37) Through (6, 2), parallel to 
$$-7x + 5y = -17$$

A) 
$$\frac{7}{5}$$
  $\frac{32}{5}$   
C)  $\frac{6}{5}$   $\frac{17}{5}$ 

B) 
$$\frac{7}{5}$$
  $\frac{32}{5}$ 
D)  $\frac{5}{7}$   $\frac{2}{7}$ 

38) Through (-4, -9), perpendicular to 
$$-7x + 9y = 109$$

38) Through (-4, -9), perpendicular to 
$$-7x + 9y = 109$$
  
A)  $\frac{7}{9} + \frac{7}{9}$   
B)  $\frac{9}{7}$   
 $y = -x$ 

C) 
$$\frac{9}{7}$$
  $\frac{99}{7}$ 

D) 
$$\frac{4}{9}$$
  $\frac{109}{9}$ 

39) Through (-3, 3), perpendicular to -2x + 5y = 21

A) 
$$y = -\frac{5}{2}x - \frac{9}{2}$$
C) 5 9

B) 
$$\frac{5}{2}$$

C) 
$$\frac{5}{2}$$
  $\frac{9}{2}$ 

B) 
$$\frac{5}{2}$$
  
 $y = -\frac{x}{x}$   
D)  $\frac{2}{5}$   
 $y = -\frac{x}{y} - \frac{9}{x}$ 

40) Through (-8, 5), perpendicular to x = 1

A) 
$$x = 1$$

B) 
$$y = 5$$

C) 
$$v = 1$$

C) 
$$y = 1$$
 D)  $y = -5$ 

41) The line with y-intercept -7 and perpendicular to x + 2y = 6

$$y = -\frac{1}{2}x + 1$$

B) 
$$\frac{1}{2}$$
  $y = \frac{1}{2}$   $-\frac{1}{2}$ 

C) 
$$y = 2x - 7$$

D) 
$$y = -2x - 7$$

42) The line with x-intercept 5 and perpendicular to 6x - y = -9

39) \_\_\_\_\_

40) \_\_\_\_\_

$$A) \frac{1}{6} x + \frac{5}{6}$$

B) 
$$\frac{1}{6}$$


C) 
$$y = -6x + 30$$

D) 
$$\frac{1}{6}$$
  $\frac{5}{6}$ 


Find the slope of the line.

43)





44)




A) 6

B) -1 C) -6 D) 1

45) \_\_\_\_\_

46) \_\_\_

45)




A) 1

B) -1

C) -5 D) 5


46)



A) 2

C) - 2 D)  $\frac{1}{2}$ 

47)




- C) 3

- B) 0
- D) undefined

49) \_\_\_\_\_

48)




- A) 2
- C) 0


- B) undefined
- D) -2


# Graph the equation.


49) 
$$y = 3x - 5$$




A)









D)









A)





B)




51) 2y + 9x = -18











C)


















D)













54) 
$$2x + y = -4$$











C)
















D)





56) 
$$x = -3$$


A)





B)



















C)















D)



#### Solve the problem.

59) In a certain city, the cost of a taxi ride is computed as follows: There is a fixed charge of \$2.45 as soon as you get in the taxi, to which a charge of \$2.20 per mile is added. Find a linear equation that can be used to determine the cost, C, of an x-mile taxi ride.

A) C = 2.45x + 2.20

B) C = 4.65x

C) C = 2.20x + 2.45

D) C = 3.15x

60) After two years on the job, an engineer's salary was \$45,000. After seven years on the job, her salary was \$57,500. Let y represent her salary after x years on the job. Assuming that the change in her salary over time can be approximated by a straight line, give an equation for this line in the form y = mx + b.

A) y = 2500x + 40,000

B) y = 12,500x + 20,000

C) y = 12,500x + 45,000

D) y = 2500x + 45,000

61) Suppose that the population of a certain town, in thousands, was 105 in 1990 and 141 in 2002. Assume that the population growth can be approximated by a straight line. Find the equation of a line which will estimate the population of the town, in thousands, in any given year since 1990.

61) \_\_\_\_\_

62) \_\_\_

63) \_\_\_\_\_

59)

60) \_\_\_

A) y = 4.25x + 90 where x is the number of years since 1990

B) y = 2.5x + 105 where x is the number of years since 1990

C) y = -3x + 177 where x is the number of years since 1990

D) y = 3x + 105 where x is the number of years since 1990

62) Assume that the sales of a certain appliance dealer can be approximated by a straight line. Suppose that sales were \$ 8500 in 1982 and \$ 58,500 in 1987. Let x=0 represent 1982. Find the equation giving yearly sales S.

A) S = 10,000x + 58,500

B) S = 10,000x + 8500

C) S = 50,000x + 58,500

D) S = 50,000x + 8500

63) The cost of owning a home includes both fixed costs and variable utility costs. Assume that it costs \$1859 per month for mortgage and insurance payments and it costs an average of \$3.03 per unit for natural gas, electricity, and water usage. Determine a linear equation that computes the annual cost of owning this home if x utility units are used.

A) y = -3.03x + 22,308

B) y = -3.03x + 1859

C) 
$$y = 3.03x + 22,308$$

D) 
$$y = 3.03x + 1859$$

64) In a lab experiment 4 grams of acid were produced in 18 minutes and 7 grams in 38 minutes. Let y be the grams produced in x minutes. Write a linear equation for grams produced.

A) 
$$\frac{3}{20}$$
  $\frac{13}{10}$   
C)  $\frac{3}{20}$   $\frac{13}{10}$ 

B) 
$$\frac{20}{3}$$
  $\frac{13}{10}$ 

D)  $\frac{3}{20}$   $\frac{13}{10}$ 
 $y = x$ 

65) A biologist recorded 9 snakes on 16 acres in one area and 19 snakes 65) \_ on 19 acres in another area. Let y be the number of snakes in x acres. Write a linear equation for the number of snakes.

A) 
$$\frac{10}{3}$$
  $\frac{133}{3}$   
C)  $\frac{10}{3}$   $\frac{133}{3}$ 

B) 
$$\frac{10}{3}$$
  $\frac{133}{3}$ 
 $y = -\frac{1}{3}x + \frac{1}{3}$ 
 $y = \frac{1}{3}$ 
 $y = \frac{1}{3}$ 

66) The following data show the list price, x, in thousands of dollars, and 66) \_\_\_\_ the dealer invoice price, y, also in thousands of dollars, for a variety of sport utility vehicles. Find a linear equation that approximates the data, using the points (16.5, 16.1) and (20.0, 18.3).

 ne ponius (10.0, 10.1) una (20.0, 10						
List Price	Dealer Invoice Price					
16.5	16.1					
17.6	17.0					
20.7	18.2					
23.1	19.3					
20.0	18.3					
24.6	21.0					

A) 
$$y = 0.629x + 5.73$$

B) 
$$y = 1.59x - 9.11$$

C) 
$$y = 1.59x - 10.2$$

D) 
$$y = 0.629x + 6.38$$

67) The information in the chart gives the salary of a person for the stated years. Model and (3, 26,400).

Year, x | Salary, y

ation in the chart gives the salary of a person for the stated	67)	
el the data with a linear function using the points $(1, 24,700)$		
00).		
The state of the s		

Year, x	Salary, y
1990, 0	\$23,500
1991, 1	\$24,700
1992, 2	\$25,200
1993, 3	\$26,400
1994, 4	\$27,200

A) 
$$y = 850x + 23,850$$

B) 
$$y = -1097x + 23,850$$

68) \_\_\_

C) 
$$y = 28.1x + 23,850$$

D) 
$$y = 850x$$

68) The change in a certain engineer's salary over time can be approximated by the linear equation y = 1500x + 47,500 where y represents salary in dollars and x represents number of years on the job. According to this equation, after how many years on the job was the engineer's salary \$61,000?

	A) 11 years	B) 10 years	C) 9 years	D) 8 years		
69)	The relationship bet the dealer invoice pr trucks can be approx Use this equation to	69)	_			
	with a list price of A) 24.727 thousan	d dollars	B) 17.478 thou			
	C) 21.986 thousan	d dollars	D) 14.658 thou	sand dollars		
70)	Suppose the sales of a particular brand of appliance satisfy the relationship $S = 60x + 3300$ , where S represents the number of sales in year x, with $X = 0$ corresponding to 1982. Find the number of sales in					_
	19 96.					
	A) 8220 sales C) 8280 sales		B) 4140 sales D) 4080 sales			
71)	The mathematical m	anufacturing x ite ost to produce 6	ems during a mont 00 items?	th. Based on this,	71)	_
	A) \$ 600,000	B) \$ 66.67	C) \$ 0.11	D) \$ 540,000		
72)	Suppose the function elapsed, in minutes, time. Find the actual minutes.  A) 21.3 min  C) 61.08 min	for t minutes of a	a person's estimate	of the elapsed	72)	_
73)	A car rental compan car and \$0.17 per n How many miles did A) 319 mi	73)	_			
	A) 319 IIII	D) 209 IIII	C) 143 mi	D) 136 IIII		
74)	If an object is dropped second), of the object by 32 and adding 10 velocity, V, in terms predict the velocity (A) 184.1 feet per second 182.1 feet per sec	t after t seconds to the result. Wr of the number of of the object at tir econd	can be obtained b ite an equation exp seconds, t. Use t	y multiplying toressing the his function to er second	74)	_
75)	The information in t stated years. Mode 24,400) and (3, 26,80) the year 2007.	he chart below gi ll the data with a	ives the salary of a linear function usi	person for the ng the points (1,	75) Year, x 1990, 0 1991, 1 1992, 2 1993, 3 1994, 4	Salary, y \$23,500 \$24,400 \$25,200 \$26,800 \$27,200

78) \_\_\_\_\_

79) \_\_\_\_

80) \_\_\_\_\_

81) \_\_\_\_\_

82) \_\_\_\_\_

83) \_\_\_\_\_

84) \_\_\_\_\_

85)

86) \_\_\_\_\_

87) \_\_\_\_\_

- 76) In order to receive a B in a course, it is necessary to get an average of 80% correct on two one-hour exams of 100 points each, on one midterm exam of 200 points, and on one final exam of 500 points. If a student scores 92, and 84 on the one-hour exams, and 145 on the midterm exam, what is the minimum score on the final exam that the person can
  - A) 399
- B) 309
- C) 579
- D) 444

Evaluate the function as indicated.

get and still earn a B?

- 77) Find f(1) when f(x) = -13x 16.
- B) -29
- C) 3
- D) -14.6

- 78) Find f(5) when f(x) = 3x 1.
  - A) 16
- B) 2
- C) 10
- D) 14

- 79) Find f(0) when f(x) = -6x 10.
  - A) 0
- B) -6
- C) -16
- D) -10

- 80) Find f(6.0) when f(x) = -5.5x + 1.
  - A) -32
- B) -34
- C) -32.9
- D) 34

- 81) Find f(7.3) when f(x) = 3x + 3.9.
  - A) 22.29
- B) 25.8
- C) -18
- D) 18

- 82)  $\left(\frac{1}{2}\right)$  when g(x) = 8 9x. A)  $\frac{7}{2}$  B)  $\frac{25}{2}$

- 83) Find f(3.2) when f(x) = 8.
  - A) 8
- C) 25.6
- D) 3.2

- 84) Find f(-r) when f(x) = 2 4x.
  - A) 2 4r
- B) 2 + 4r
- C) 2 + rx
- D) r 4x
- 85) Find  $g(^{n^2})$  when g(x) = -7 + 2x. and  $g(^{11-})$  when g(x) = -7 + 2x. A)  $_{-7 + 2}n^2$  B)  $_{-7 + 2}x^2$  C)  $_{-7 + n^2}$  D)  $_{-7 - 2}n^2$

- 86) Find g(a + 1) when g(x) = 4x + 1.
- B) 4a + 1 C) 4a 1
- D) 4a + 5
- Write a cost function for the problem. Assume that the relationship is linear.
  - 87) A moving firm charges a flat fee of \$ 40 plus \$ 35 per hour. Let C(x) be the cost in dollars of using the moving firm for x hours.
    - A) C(x) = 40x 35
- B) C(x) = 35x + 40
- C) C(x) = 40x + 35
- D) C(x) = 35x 40

88) A cab company charges a base rate of \$ 1.00 plus 
$$10$$
 cents per minute. Let  $C(x)$  be the cost in dollars of using the cab for  $x$  minutes.

A) 
$$C(x) = 1.00x - 0.10$$

B) 
$$C(x) = 0.10x - 1.00$$

C) 
$$C(x) = 1.00x + 0.10$$

D) 
$$C(x) = 0.10x + 1.00$$

89) An electrician charges a fee of \$ 55 plus \$ 40 per hour. Let 
$$C(x)$$
 be the cost in dollars of using the electrician for x hours.

A) 
$$C(x) = 55x + 40$$

B) 
$$C(x) = 55x - 40$$

C) 
$$C(x) = 40x + 55$$

D) 
$$C(x) = 40x - 55$$

A) 
$$C(x) = 29x - 6$$

B) 
$$C(x) = 29x + 6$$

C) 
$$C(x) = 6x + 29$$

D) 
$$C(x) = 6x - 29$$

A) 
$$C(x) = 582x + 3320$$

B) 
$$C(x) = 1164x + 410$$

C) 
$$C(x) = 582x + 410$$

D) 
$$C(x) = 1164x + 3320$$

93) \_\_\_\_\_

A) 
$$C(x) = 150x + 5000$$

B) 
$$C(x) = 150x + 500$$


C) 
$$C(x) = 17x + 5000$$


D) 
$$C(x) = 17x + 500$$


#### Solve the problem.

$$\frac{2}{p} = S(q) = \frac{2}{3}$$
 and  $p = D(q) = 12 - \frac{2}{3}$  p is the price in dollars and q is the quantity of


where p is the price in dollars and q is the quantity of pencil sharpeners (in hundreds). Graph these functions on the same axes (graph the supply function as a dashed line and the demand function as a solid line). Also, find the equilibrium quantity and the equilibrium price.








Equilibrium quantity: 720 Equilibrium price: \$ 4.80



Equilibrium quantity: 900 Equilibrium price: \$ 6





Equilibrium quantity: 480

94) Let the supply and demand functions for raspberry-flavored licorice be given by


$$p = S(q) = \frac{\frac{5}{4}}{q}$$
 and  $p = D(q) = \frac{\frac{3}{4}}{q}$ ,

where p is the price in dollars and q is the number of batches. Graph these functions on the same axes (graph the supply function as a dashed line and the demand function as a solid line). Also, find the equilibrium quantity and the equilibrium price.






Equilibrium quantity: 35.00 Equilibrium price: \$ 28




Equilibrium quantity: 43.75

Equilibrium

price: \$ 35



Equilibrium quantity: 35 Equilibrium price: \$ 43.75



Equilibrium quantity: 28 Equilibrium price: \$ 35.00

95) Given the supply and demand functions below, find the price when the demand is 145.

95) \_\_\_\_\_

$$S(p) = 9p + 12$$

$$D(p) = 280 - 9p$$

96) Suppose that the demand and price for a certain model of graphing calculator are related by p = D(q) = 93 - 2.25q, where p is the price (in dollars) and q is the demand (in hundreds). Find the price if the demand is 300 calculators.

96) \_\_\_\_\_

- A) \$ 25.50
- B) \$ 99.75
- C) \$86.25
- D) \$ 160.50
- 97) Given the supply and demand functions below, find the demand when p = \$12.


97) \_\_\_\_\_


$$S(p) = 5p$$

$$D(p) = 120 - 4p$$

- B) 48
- C) 132
- D) 72
- 98) Suppose that the demand and price for a certain model of graphing


calc ulator





108) Midtown Delivery Service delivers packages which cost \$ 2.30 per

pack age to



- 122) Find the temperature at which the Celsius and Fahrenheit scales coincide.

- A) -40°
- B) 38°
- C) 0°
- D) -26°


123) For the following table of data,


123) \_\_\_\_\_

122) \_\_


- a. Draw a scatterplot.
- b. Calculate the correlation coefficient.
- c. Calculate the least squares line and graph it on the scatterplot.
- d. Predict the y-value when x is 22.
- x
   1
   2
   3
   4
   5
   6
   7
   8
   9

   y
   5
   4.5
   4
   4
   3
   3.5
   2.5
   2
   1







A) a.



- b. 0.965
- c. Y = 0.45x 5.53




- d. 4.37
- B) a.




b. 0.965

c. 
$$Y = -0.45x$$


+ 5.53




C) a.




c. Y = -0.45x - 5.53



D) a.



c. Y = -0.45x + 5.53



d. -4.37


- 124) For the following table of data,
  - a. Draw a scatterplot.
  - b. Calculate the correlation coefficient.
  - c. Calculate the least squares line and graph it on the scatterplot.


d. the y-valu

Predie


ct when

X	-4	-3	-2	-1	0	1	2	3	4
v	-3.5	-2	-1	-1.5	1	0.5	1.5	2	4






A) a.




b. 0.966

c. 
$$Y = 0.82x + 0.11$$




d. -17.11

C) a.




B) a.




b. -0.966


c. 
$$Y = 0.82x + 0.11$$




d. -17.33







- b. 0.966
- c. Y = 0.82x 0.11





d. -17.33

125) For the following table of data,


- 125) \_\_\_\_\_
- a. Draw a scatterplot.
- b. Calculate the correlation coefficient.
- c. Calculate the least squares line and graph it on the scatterplot.
- d. Predict the y-value when x is 14.

x	1	2	3	4	5	6	7	8	9
у	10	12	11	14	15	17	22	19	24






b. 0.950 c. Y = 1.7x + 7.5




31


.3




- b. 0.950
- c. Y = 1.7x




b. 0.950



D) a.



- b. 0.903
- c. Y = 2x + 6.5



Find the coefficient of correlation.

d.

32.3

126) Consider the data points with the following coordinates:

- A) 0
- B) 0.4383
- C) -0. 4383
- D) 0.3901

126)

127) \_\_\_\_

128) \_\_

127) The test scores of 6 randomly picked students and the number of hours they prepared are as follows:

-۲	arca a	iic a	<i>j</i> 101.	LOVVS	•		
]	Hours	5	10	4	6	10	9
	Score	64	86	69	86	59	87

- A) -0.6781
- B) -0.2242
- C) 0.2242
- D) 0.6781
- 128) The test scores of 6 randomly picked students and the number of hours they prepared are as follows:

Hours	4	10	5	5	3	3
Score	54	99	56	99	70	72

- A) 0.2015
- B) -0.2241
- C) -0.6781
- D) 0.6039

129) Consider the data points with the following coordinates:	129)
x     57     53     59     61     53     56     60       y     156     164     163     177     159     175     151	/
y 156 164 163 177 159 175 151	
A) 0.2145 B) -0.0783 C) -0.0537 D) 0.1085	
130) Consider the data points with the following coordinates:	130)
x     62     53     64     52     52     54     58       y     158     176     151     164     164     174     162	
y 158 176 151 164 164 174 162	
A) -0.7749 B) 0 C) -0.0810 D) 0.7537	
101) Consider the data relations that he following conditions	101)
131) Consider the data points with the following coordinates:	131)
x         121         101         128         160         154         126         134           y         171         152         168         157         164         169         160	
A) -0.0781 B) 0.0537 C) 0.5370 D) 0.2245	
11) 0.0701	
132) The following are costs of advertising (in thousands of dollars) and the	132)
number of products sold (in thousands):	/ ====
Cost 9 2 3 4 2 5 9 10	
Cost         9         2         3         4         2         5         9         10           Number         85         52         55         68         67         86         83         73	
A) 0.2353 B) -0.0707 C) 0.7077 D) 0.2456	
133) The following are costs of advertising (in thousands of dollars) and the	133)
number of products sold (in thousands):	
Cost 6 3 7 6 10 4 7 7 Number 54 75 91 57 96 52 92 100	
A) 0.6112 B) 0.6756 C) -0.3707 D) 0.2635	
A) 0.0112 b) 0.0730 C) -0.3707 b) 0.2033	
134) The following are the temperatures on randomly chosen days and the	134)
amount a certain kind of plant grew (in millimeters):	101)
Temp 62 76 50 51 71 46 51 44 79	
Temp 62 76 50 51 71 46 51 44 79 Growth 36 39 50 13 33 33 17 6 16	
A) 0 B) -0.2105 C) 0.1955 D) 0.2563	
135) The following are the temperatures on randomly chosen days and the	135)
amount a certain kind of plant grew (in millimeters):	
Temp 77 88 85 61 64 72 73 63 74	
Growth 39 17 12 22 15 29 14 25 43	
A) -0.0953 B) 0.0396 C) 0 D) -0.3105	
Find the equation of the least assessed line	
Find the equation of the least squares line.	2.0
136) Ten students in a graduate program were randomly selected. Their	3.9
grade point averages (GPAs) when they entered the program were between 3.5 and 4.0. The following data were obtained regarding their	
GPAs on entering the program versus their current GPAs.	4.0
Of 115 of effecting the program versus their current of 115.	4.0
Entering GPA (x) Current GPA (y)	3.9
3.5 3.6	0.7
3.8 3.7	
3.6 3.9	3.7
3.6	3.5
3.5 3.9	
2.0	

3.9

3.8

3

3

4

.0

.8

A) 
$$y = 2.51 + 0.329x$$

C) 
$$y = 5.81 + 0.497x$$

B) 
$$y = 4.91 + 0.0212x$$

D) 
$$y = 3.67 + 0.0313x$$

137) \_\_\_\_\_

138) \_\_\_\_\_

139)

58

7

137) The paired data below consist of the test scores of 6 randomly selected students and the number of hours they studied for the test.

A) 
$$y = 33.7 + 2.14x$$

B) 
$$y = 33.7 - 2.14x$$

C) 
$$y = -67.3 + 1.07x$$

D) 
$$y = 67.3 + 1.07x$$

138) The paired data below consist of the costs of advertising (in thousands of dollars) and the number of products sold (in thousands).

A) 
$$y = 55.8 + 2.79x$$

B) 
$$y = 55.8 - 2.79x$$

C) 
$$y = 26.4 + 1.42x$$

D) 
$$y = -26.4 - 1.42x$$

139) The paired data below consist of the temperatures on randomly chosen days and the amount a certain kind of plant grew (in millimeters).

A) 
$$y = -14.6 - 0.211x$$

B) 
$$y = 7.30 + 0.122x$$

C) 
$$y = 7.30 - 0.112x$$

D) 
$$y = 14.6 + 0.211x$$

140) A study was conducted to compare the average time spent in the lab each week versus course grade for computer students. The results are recorded in the table below.

Number of hours spent in lab (x) Grade (percent)(y)

51

16

A) 
$$y = 0.930 + 44.3x$$

C) 
$$y = 88.6 - 1.86x$$

B) 
$$y = 1.86 + 88.6x$$

D) 
$$y = 44.3 + 0.930x$$

141) \_\_\_\_\_

141) Two separate tests are designed to measure a student's ability to solve problems. Several students are randomly selected to take both tests and the results are shown below.

A) 
$$y = 0.930 - 19.4x$$

B) 
$$y = -19.4 - 0.930x$$

C) 
$$y = 19.4 + 0.930x$$

D) 
$$y = -0.930 + 19.4x$$

142) Managers rate employees according to job performance and attitude. The results for several randomly selected employees are given below.

142) \_\_\_\_\_

Attitude (x)	59									
Performance (y)	72	67	78	82	75	87	92	83	87	78

- A) y = 11.7 + 1.02x
- B) y = 2.81 + 1.35x
- C) y = 92.3 0.669x

- D) y = -47.3 + 2.02x
- 143) Two different tests are designed to measure employee productivity and dexterity. Several employees of a company are randomly selected and asked to complete the tests. The results are below.

143)	

Dexterity (x)	23	25	28	21	21	25	26	30	34	36
Productivity (y)	49	53	59	42	47	53	55	63	67	75

- A) y = 5.05 + 1.91x
- B) y = 2.36 + 2.03x

C) y = 75.3 - 0.329x

- D) y = 10.7 + 1.53x
- 144) In the table below, x represents the number of years since 2000 and y represents annual sales (in thousands of dollars) for a clothing company.

A) y = 3.31x + 23.8

B) y = 5.18x + 20.6

C) y = 2.61x + 25.9

- D) y = 4.37x + 21.7
- 145) In the table below, x represents the number of years since 2000 and y represents the population (in thousands) of the town Boomville.



A) y = 18x + 8

B) y = 25x - 5

C) y = 12x + 20

D) y = 28x - 10

## Solve the problem.


- 146) Find an equation for the least squares line representing weight, in pounds, as a function of height, in inches, of men. Then, predict the weight of a man who is 68 inches tall to the nearest tenth of a pound. The following data are the (height, weight) pairs for 8 men: (66, 150), (68, 160), (69, 166), (70, 175), (71, 181), (72, 191), (73, 198), (74, 206).
- 146) \_\_\_\_\_

A) 161.2 pounds

B) 151.4 pounds

C) 160.0 pounds

- D) 165.1 pounds
- 147) Find an equation for the least squares line representing weight, in pounds, as a function of height, in inches, of men. Then, predict the height of a man who is 145 pounds to the nearest tenth of an inch. The following data are the (height, weight) pairs for 8 men: (66, 150), (68,
- 160), 175),
- (69, (71,
- 166), 181),
- (70, (72,



151) The paired data below consist of the test scores of 6 randomly selected students and the number of hours they studied for the test. Use the

equa of the tion least

squares line to predict the score on the test of a student who studies 15 hours.	151)				
Hours (x Score (y)	) 5 10 4 6 10 0 64 86 69 86 59	9 87			
	A) 83.3	B) 78.3	C) 88.3	D) 89.8	
152)	The paired data belo of dollars) and the no equation of the least if the cost of advertis	umber of products squares line to pre	sold (in thousands	s). Use the	152)
	Cost (x) Number (y)	9 2 3 4 2 9 85 52 55 68 67	5     9     10       86     83     73		
	A) 97.65 products C) 94.65 products		B) 104.35 produc D) 41,905.8 produ		
153)	The paired data belo days and the amoun the equation of the lethe temperature is	t a certain kind of geast squares line to	plant grew (in mill	imeters). Use	153)
	$\frac{\text{Temp }(x)}{\text{Growth }(y)}$	62 76 50 51 7 36 39 50 13	71 46 51 44 79 33 33 17 6 16	-	
	A) 31.48 mm C) 29.80 mm		B) 32.76 mm D) 32.12 mm		
154)	In the table below, x represents annual sa company. Use the l the year 2006. Round	les (in thousands c east squares regre	of dollars) for a consistency for a consistency of the second sec	clothing	154)
	Year (x) 3 Sales (y) 30 A) \$147,000	2 3 4 5 0 40 60 90 130 B) \$140,000	- C) \$145,000	D) \$142,000	
155)	A study was conducted each week versus correcorded in the table to predict the grade of	urse grade for com below. Use the e	puter students. The quation of the leas	e results are t squares line	of

Grade 7 155) \_\_\_\_
(percent) 89
(y) 15

81

16

46

10

Provide an appropriate response.

156) Find k so that the line through (3, k) and (1, -2) is parallel to 2x - 4y = 7. 156) \_\_\_\_\_\_ Find k so that the line is perpendicular to 4x + 4y = -4.

A) 3; -4

B) - 1; - 4

C) 3; 0

D) - 1; 0

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

157) John has been a teacher at West Side High School for the past 12 years. His salary during that time can be modeled by the linear equation y = 800x + 33,000 where x is the number of years since he began teaching at West Side and y is his salary in dollars. Explain what the slope, 800, represents in this context.

157) \_\_\_\_\_

158) If a company decides to make a new product, there are fixed costs and variable costs associated with this new product. Explain the differences of the two types of costs and why they occur. Use an example to illustrate your point.

158) \_\_\_\_\_

159) Give a definition or an example of the word or phrase: Undefined slope

159) \_\_\_\_\_

160) Why is the slope of a horizontal line equal to zero? Give an example.

160) \_\_\_\_\_

161) Explain what is wrong with the statement "The line has no slope."

161) \_\_\_\_\_

162) Why is the slope of a vertical line undefined?

162) \_\_\_\_\_

163) Can an equation of a vertical line be written in slope-intercept form? Explain.

163) \_\_\_\_\_

164) The total number of reported cases of AIDS in the United States has risen from 372 in 1981 to 100,000 in 1989 and 200,000 in 1992. Does a linear equation fit this data? Explain.

164) \_\_\_\_\_

165) Show that the points  $P_1(2,4)$ ,  $P_2(5,2)$ , and  $P_3(7,5)$  are the vertices of a right triangle.

165) \_\_\_\_\_

- 1) B
- 2) A
- 3) B
- 4) B
- 5) D
- 6) C
- 7) D
- 8) B
- 9) D
- 10) A
- 11) D
- 12) B
- 13) B
- 14) C
- 15) A
- 16) D
- 17) C 18) D
- 19) B
- 20) A 21) D
- 22) C
- 23) C
- 24) A
- 25) C
- 26) C
- 27) D
- 28) A
- 29) B
- 30) D
- 31) B
- 32) C
- 33) D
- 34) A
- 35) A
- 36) B
- 37) A
- 38) B
- 39) A
- 40) B 41) C
- 42) D
- 43) D
- 44) B
- 45) A
- 46) A
- 47) D
- 48) C 49) D
- 50) D
- 51) A

- 52) C
- 53) B
- 54) B
- 55) B
- 56) D
- 57) B
- 58) A
- 59) C
- 60) A
- 61) D
- 62) B
- 63) C
- 64) C
- 65) A
- 66) A
- 67) A
- 68) C
- 69) B
- 70) B
- 71) A
- 72) A
- 73) C
- 74) B
- 75) B
- 76) A
- 77) B
- 78) D
- 79) D
- 80) A
- 81) B
- 82) A
- 83) A
- 84) B
- 85) A
- 86) D
- 87) B
- 88) D
- 89) C
- 90) C
- 91) C
- 92) B
- 93) C
- 94) C
- 95) A 96) C
- 97) D
- 98) B
- 99) B
- 100) D
- 101) A 102) D
- 103) A

- 104) C
- 105) C
- 106) B
- 100) D
- 107) A
- 108) D
- 109) C
- 110) D
- 111) A
- 112) D
- 113) D
- 114) B
- 115) B
- 116) C
- 117) A
- 118) C
- 119) A
- 120) A
- 121) A
- 122) A
- 123) D
- 124) A
- 125) A
- 126) B
- 127) C
- 128) D
- 129) D
- 130) A
- 131) B
- 132) C 133) A
- 134) C
- 135) A
- 136) D
- 137) D
- 107) D
- 138) A
- 139) D
- 140) C
- 141) C
- 142) A
- 143) A
- 144) A
- 145) B
- 146) A
- 147) A
- 148) C
- 149) C
- 150) B
- 151) A
- 152) A 153) A
- 154) C
- 155) B

- 156) D
- 157) The slope of 800 indicates that during his 12 years at the school, John's salary has increased by approximately \$800 per year.
- 158) Fixed costs occur only once. These costs may be startup costs related to the production of the new product. Variable costs depend on how much product is made. These costs may consist of labor, material, and maintenance.

For example, a company decided to make oak filing cabinets. Fixed costs would include the costs of purchasing and renovating plant space and the cost of manufacturing equipment. Variable costs would include the cost labor and the cost of materials.

- 159) An equation such as ax + c = 0 has an undefined slope. (Answers may vary.)
- 160) Answers may vary. One possibility: The slope of a horizontal line is equal to zero because the y-values do not change as the x-values change. For example, the points (3, 4) and (7, 4)  $\underline{4-4} = \underline{0}$

are two points on a horizontal line. The slope of this line is zero because  $m = \begin{pmatrix} 7 - 3 \\ = \end{pmatrix} = \begin{pmatrix} 4 \\ = \end{pmatrix}$ 

- 161) Answers may vary. One possibility: It is not specific enough. The slope of a horizontal line is 0, while the slope of a vertical line is undefined.
- 162) Answers may vary. One possibility: Let (a, b) and (a, c),  $b \neq c$ , be any two different points on a vertical line. The slope of the line =  $\frac{y_1 y_2}{x_1 x_2} = \frac{b c}{a a} = \frac{b c}{0}$ . Division by zero is undefined.
- 163) No. In the slope-intercept form of the equation of a line, x is multiplied by slope; however, the slope of a vertical line is undefined. (Explanations will vary.)
- 164) No, the data cannot be modeled by a linear equation because the reported cases are not increasing at a constant rate. Assume a linear equation, and examine the slope of the two line segments. The slope of the segment from (0, 372) to (8, 100,000) is 12,453.5 while the slope of the segment from (8, 100,000) to (11, 200,000) is  $33,333.\overline{3}$ . (Explanations will vary.)
- 165) Answers will vary. One possibility: The slope of the line through  $P_1$  and  $P_2$  is -2/3. The slope of the line through  $P_2$  and  $P_3$  is 3/2. Therefore, since the product of these slopes is  $^{-1}$ , the lines are perpendicular and constitute a right angle in the triangle, making the triangle formed by these points a right triangle.