


TEST BANK



SULLIVAN



Precalculus⁸

Ch. 2 Functions and Their Graphs


2.1 Functions

1 Determine Whether a Relation Represents a Function

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Determine whether the relation represents a function. If it is a function, state the domain and range.

1)



A) function

domain: {3, 7, 11, 15}

range: {9, 21, 33, 45}


B) function

domain: {9, 21, 33, 45}

range: {3, 7, 11, 15}

C) not a function

2)



A) function

domain: {Bob, Ann, Dave}

range: {carrots, peas, squash}


B) function

domain: {carrots, peas, squash}

range: {Bob, Ann, Dave}

C) not a function

3)



A) function

domain: {Bob, Ann, Dave}

range: {Ms. Lee, Mr. Bar}

B) function

domain: {Ms. Lee, Mr. Bar}

range: {Bob, Ann, Dave}

C) not a function

4) $\{(-3, -6), (1, 4), (4, 1), (7, -1)\}$

A) function

domain: $\{-3, 1, 4, 7\}$

range: $\{-6, 4, 1, -1\}$

B) function

domain: $\{-6, 4, 1, -1\}$

range: $\{-3, 1, 4, 7\}$

C) not a function

5) $\{(11, -4), (-5, -3), (-5, 0), (4, 3), (20, 5)\}$

A) function

domain: $\{11, 4, -5, 20\}$

range: $\{-4, -3, 0, 3, 5\}$

B) function

domain: $\{-4, -3, 0, 3, 5\}$

range: $\{11, 4, -5, 20\}$

C) not a function

6) $\{(-4, 11), (-3, 4), (0, -5), (3, 4), (5, 20)\}$

A) function

domain: $\{-4, -3, 0, 3, 5\}$

range: $\{11, 4, -5, 20\}$

B) function

domain: $\{11, 4, -5, 20\}$

range: $\{-4, -3, 0, 3, 5\}$

C) not a function

7) $\{(8.66, 15.86), (8.666, -15.9), (\frac{9}{7}, 0), (1.29, -8)\}$

A) function

domain: $\{8.66, 8.666, \frac{9}{7}, 1.29\}$

range: $\{15.86, -15.9, 0, -8\}$

B) function

Determine whether the equation defines y as a function of x .

8) $y = x^4$

A) function

B) not a function

9) $y = \frac{1}{x}$

A) function

B) not a function

10) $y = |x|$

A) function

B) not a function

11) $y^2 = 4 - x^2$

A) function

B) not a function

12) $y = \pm\sqrt{1 - 3x}$

A) function

B) not a function

13) $x = y^2$

A) function

B) not a function

14) $y^2 + x = 2$

A) function

B) not a function

15) $y = 2x^2 - 5x + 6$

A) function

B) not a function

$$16) y = \frac{4x + 1}{x - 2}$$

A) function

B) not a function

$$17) x^2 + 2y^2 = 1$$

A) function

B) not a function

$$18) x - 3y = 9$$

A) function

B) not a function

$$19) -5x + x^2 - 63 = y$$

A) function

B) not a function

2 Find the Value of a Function

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Find the value for the function.

1) Find $f(-2)$ when $f(x) = x^2 - 5x - 1$.

A) 13

B) 15

C) -5

D) -7

2) Find $f(1)$ when $f(x) = \frac{x^2 - 5}{x + 2}$.

A) $-\frac{4}{3}$

B) 2

C) $\frac{1}{3}$

D) 4

3) Find $f(-9)$ when $f(x) = |x| - 6$.

A) 3

B) -15

C) 15

D) -3

4) Find $f(3)$ when $f(x) = \sqrt{x^2 + 5x}$.

A) $2\sqrt{6}$

B) $\sqrt{34}$

C) $\sqrt{30}$

D) $\sqrt{14}$

5) Find $f(-x)$ when $f(x) = 3x^2 - 2x + 4$.

A) $3x^2 + 2x + 4$

B) $-3x^2 + 2x - 4$

C) $3x^2 + 2x - 4$

D) $-3x^2 + 2x + 4$

6) Find $f(-x)$ when $f(x) = \frac{x}{x^2 + 4}$.

A) $\frac{-x}{x^2 + 4}$

B) $\frac{-x}{x^2 - 4}$

C) $\frac{x}{-x^2 + 4}$

D) $\frac{-x}{-x^2 + 4}$

7) Find $-f(x)$ when $f(x) = -3x^2 + 4x + 1$.

A) $3x^2 - 4x - 1$

B) $-3x^2 - 4x + 1$

C) $-3x^2 - 4x - 1$

D) $3x^2 - 4x + 1$

8) Find $-f(x)$ when $f(x) = |x| - 9$.

A) $-|x| + 9$

B) $| -x | - 9$

C) $-|x| - 9$

D) $| -x | + 9$

- 9) Find $f(x - 1)$ when $f(x) = 4x^2 + 3x + 5$.
- A) $4x^2 - 5x + 6$ B) $-5x^2 + 4x + 6$ C) $4x^2 - 5x + 12$ D) $4x^2 + 23x + 12$
- 10) Find $f(x + 1)$ when $f(x) = \frac{x^2 - 9}{x - 3}$.
- A) $\frac{x^2 + 2x - 8}{x - 2}$ B) $\frac{x^2 + 2x + 10}{x - 2}$ C) $\frac{x^2 + 2x - 8}{x + 4}$ D) $\frac{x^2 - 8}{x - 2}$
- 11) Find $f(2x)$ when $f(x) = 3x^2 - 5x + 2$.
- A) $12x^2 - 10x + 2$ B) $6x^2 - 10x + 2$ C) $6x^2 - 10x + 4$ D) $12x^2 - 10x + 4$
- 12) Find $f(2x)$ when $f(x) = \sqrt{7x^2 + 4x}$.
- A) $\sqrt{28x^2 + 8x}$ B) $2\sqrt{7x^2 + 4x}$ C) $\sqrt{14x^2 + 8x}$ D) $\sqrt{14x^2 + 16x}$
- 13) Find $f(x + h)$ when $f(x) = 3x^2 + 2x + 1$.
- A) $3x^2 + 6xh + 3h^2 + 2x + 2h + 1$ B) $3x^2 + 3h^2 + 2x + 2h + 1$
 C) $3x^2 + 3h^2 + 8x + 8h + 1$ D) $3x^2 + 3xh + 3h^2 + 2x + 2h + 1$
- 14) Find $f(x + h)$ when $f(x) = \frac{3x + 7}{8x - 5}$.
- A) $\frac{3x + 3h + 7}{8x + 8h - 5}$ B) $\frac{3x + 3h + 7}{8x - 5}$ C) $\frac{3x + 10h}{8x + 3h}$ D) $\frac{3x + 7h}{8x - 5h}$

Solve the problem.

- 15) If $f(x) = 5x^3 + 5x^2 - x + C$ and $f(-2) = 1$, what is the value of C ?
- A) $C = 19$ B) $C = 3$ C) $C = -61$ D) $C = -21$
- 16) If $f(x) = \frac{x - B}{x - A}$, $f(2) = 0$, and $f(-7)$ is undefined, what are the values of A and B ?
- A) $A = -7, B = 2$ B) $A = 2, B = -7$ C) $A = 7, B = -2$ D) $A = -2, B = 7$
- 17) If $f(x) = \frac{x - 4A}{-12x + 3}$ and $f(-12) = 12$, what is the value of A ?
- A) $A = -444$ B) $A = 444$ C) $A = -150$ D) $A = 150$
- 18) If a rock falls from a height of 50 meters on Earth, the height H (in meters) after x seconds is approximately $H(x) = 50 - 4.9x^2$.
 What is the height of the rock when $x = 1.9$ seconds? Round to the nearest hundredth, if necessary.
- A) 32.31 m B) 67.69 m C) 40.69 m D) 32.67 m
- 19) If a rock falls from a height of 40 meters on Earth, the height H (in meters) after x seconds is approximately $H(x) = 40 - 4.9x^2$. When does the rock strike the ground? Round to the nearest hundredth, if necessary.
- A) 2.86 sec B) 8.16 sec C) 1.29 sec D) 1.67 sec

- 20) It has been determined that the number of fish $f(t)$ that can be caught in t minutes in a certain pond using a certain bait is $f(t) = 0.23t + 1$, for $t > 10$. Find the approximate number of fish that can be caught if you fish for 13 minutes.
- A) About 3 fish B) About 9 fish C) About 15 fish D) About 17 fish
- 21) The function $P(d) = 1 + \frac{d}{33}$ gives the pressure, in atmospheres (atm), at a depth d feet in the sea. Find the pressure at 37 feet.
- A) $\frac{70}{33}$ atm B) $\frac{37}{33}$ atm C) $\frac{4}{33}$ atm D) $\frac{38}{33}$ atm
- 22) The function F described by $F(C) = \frac{9}{5}C + 32$ gives the Fahrenheit temperature corresponding to the Celsius temperature C . Find the Fahrenheit temperature equivalent to -30°C .
- A) -22°F B) -76°F C) -130°F D) -184°F

3 Find the Domain of a Function

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Find the domain of the function.

1) $f(x) = -4x + 4$

- A) all real numbers B) $\{x \mid x \geq -4\}$ C) $\{x \mid x \neq 0\}$ D) $\{x \mid x > 0\}$

2) $f(x) = x^2 + 7$

- A) all real numbers B) $\{x \mid x \geq -7\}$ C) $\{x \mid x > -7\}$ D) $\{x \mid x \neq -7\}$

3) $f(x) = \frac{x^2}{x^2 + 1}$

- A) all real numbers B) $\{x \mid x \neq -1\}$ C) $\{x \mid x > -1\}$ D) $\{x \mid x \neq 0\}$

4) $g(x) = \frac{3x}{x^2 - 1}$

- A) $\{x \mid x \neq -1, 1\}$ B) $\{x \mid x \neq 0\}$ C) $\{x \mid x > 1\}$ D) all real numbers

5) $h(x) = \frac{x - 3}{x^3 - 4x}$

- A) $\{x \mid x \neq -2, 0, 2\}$ B) $\{x \mid x \neq 0\}$ C) $\{x \mid x \neq 3\}$ D) all real numbers

6) $f(x) = \sqrt{21 - x}$

- A) $\{x \mid x \leq 21\}$ B) $\{x \mid x \neq 21\}$ C) $\{x \mid x \leq \sqrt{21}\}$ D) $\{x \mid x \neq \sqrt{21}\}$

7) $\frac{x}{\sqrt{x - 8}}$

- A) $\{x \mid x > 8\}$ B) $\{x \mid x \geq 8\}$ C) $\{x \mid x \neq 8\}$ D) all real numbers

4 Form the Sum, Difference, Product, and Quotient of Two Functions

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

For the given functions f and g , find the requested function and state its domain.

1) $f(x) = 7 - 7x$; $g(x) = -3x + 7$

Find $f + g$.

A) $(f + g)(x) = -10x + 14$; all real numbers

B) $(f + g)(x) = -3x + 7$; $\{x \mid x \neq \frac{7}{3}\}$

C) $(f + g)(x) = 4x$; all real numbers

D) $(f + g)(x) = -4x + 14$; $\{x \mid x \neq -\frac{7}{2}\}$

2) $f(x) = 4x - 9$; $g(x) = 6x - 6$

Find $f - g$.

A) $(f - g)(x) = -2x - 3$; all real numbers

B) $(f - g)(x) = -2x - 15$; $\{x \mid x \neq -\frac{15}{2}\}$

C) $(f - g)(x) = 10x - 15$; $\{x \mid x \neq 1\}$

D) $(f - g)(x) = 2x + 3$; all real numbers

3) $f(x) = 9x + 2$; $g(x) = 7x - 3$

Find $f \cdot g$.

A) $(f \cdot g)(x) = 63x^2 - 13x - 6$; all real numbers

B) $(f \cdot g)(x) = 63x^2 + 11x - 6$; $\{x \mid x \neq -6\}$

C) $(f \cdot g)(x) = 16x^2 - 13x - 1$; all real numbers

D) $(f \cdot g)(x) = 63x^2 - 6$; $\{x \mid x \neq -6\}$

4) $f(x) = 5x + 1$; $g(x) = 2x - 1$

Find $\frac{f}{g}$.

A) $(\frac{f}{g})(x) = \frac{5x + 1}{2x - 1}$; $\{x \mid x \neq \frac{1}{2}\}$

B) $(\frac{f}{g})(x) = \frac{5x + 1}{2x - 1}$; $\{x \mid x \neq -\frac{1}{5}\}$

C) $(\frac{f}{g})(x) = \frac{2x - 1}{5x + 1}$; $\{x \mid x \neq \frac{1}{2}\}$

D) $(\frac{f}{g})(x) = \frac{2x - 1}{5x + 1}$; $\{x \mid x \neq -\frac{1}{5}\}$

5) $f(x) = 16 - x^2$; $g(x) = 4 - x$

Find $f + g$.

A) $(f + g)(x) = -x^2 - x + 20$; all real numbers

B) $(f + g)(x) = 4 + x$; $\{x \mid x \neq -4\}$

C) $(f + g)(x) = -x^2 + x + 12$; all real numbers

D) $(f + g)(x) = x^3 - 4x^2 - 16x + 64$; all real numbers

6) $f(x) = x - 8$; $g(x) = 4x^2$

Find $f + g$.

A) $(f + g)(x) = 4x^2 + x - 8$; all real numbers

B) $(f + g)(x) = 4x^2 - x + 8$; all real numbers

C) $(f + g)(x) = 4x^2 + x - 8$; $\{x \mid x \neq 8\}$

D) $(f + g)(x) = -4x^2 + x - 8$; all real numbers

7) $f(x) = 2x^3 - 1$; $g(x) = 6x^2 - 1$

Find $f \cdot g$.

A) $(f \cdot g)(x) = 12x^5 - 2x^3 - 6x^2 + 1$; all real numbers

B) $(f \cdot g)(x) = 12x^6 - 2x^3 - 6x^2 + 1$; all real numbers

C) $(f \cdot g)(x) = 12x^5 - 2x^3 - 6x^2 + 1$; $\{x \mid x \neq 0\}$

D) $(f \cdot g)(x) = 2x^3 + 6x^2 + 1$; all real numbers

8) $f(x) = \sqrt{x}$; $g(x) = 5x - 6$

Find $\frac{f}{g}$.

A) $(\frac{f}{g})(x) = \frac{\sqrt{x}}{5x-6}$; $\{x \mid x \geq 0, x \neq \frac{6}{5}\}$

B) $(\frac{f}{g})(x) = \frac{\sqrt{x}}{5x-6}$; $\{x \mid x \neq \frac{6}{5}\}$

C) $(\frac{f}{g})(x) = \frac{\sqrt{x}}{5x-6}$; $\{x \mid x \neq 0\}$

D) $(\frac{f}{g})(x) = \frac{5x-6}{\sqrt{x}}$; $\{x \mid x \geq 0\}$

9) $f(x) = \sqrt{3-x}$; $g(x) = \sqrt{x-1}$

Find $f \cdot g$.

A) $(f \cdot g)(x) = \sqrt{(3-x)(x-1)}$; $\{x \mid 1 \leq x \leq 3\}$

B) $(f \cdot g)(x) = \sqrt{(3-x)(x-1)}$; $\{x \mid x \geq 0\}$

C) $(f \cdot g)(x) = \sqrt{(3-x)(x-1)}$; $\{x \mid x \neq 1, x \neq 3\}$

D) $(f \cdot g)(x) = \sqrt{-x^2-3}$; $\{x \mid x \neq 3\}$

10) $f(x) = \frac{7x+3}{2x-3}$; $g(x) = \frac{3x}{2x-3}$

Find $f + g$.

A) $(f + g)(x) = \frac{10x+3}{2x-3}$; $\{x \mid x \neq \frac{3}{2}\}$

B) $(f + g)(x) = \frac{4x-3}{2x-3}$; $\{x \mid x \neq \frac{3}{2}\}$

C) $(f + g)(x) = \frac{10x+3}{2x-3}$; $\{x \mid x \neq \frac{3}{2}, x \neq -\frac{3}{10}\}$

D) $(f + g)(x) = \frac{10x+3}{2x-3}$; $\{x \mid x \neq 0\}$

11) $f(x) = \sqrt{x+11}$; $g(x) = \frac{4}{x}$

Find $f \cdot g$.

A) $(f \cdot g)(x) = \frac{4\sqrt{x+11}}{x}$; $\{x \mid x \geq -11, x \neq 0\}$

B) $(f \cdot g)(x) = \frac{\sqrt{4x+44}}{x}$; $\{x \mid x \geq -11, x \neq 0\}$

C) $(f \cdot g)(x) = \sqrt{\frac{4x+44}{x}}$; $\{x \mid x \geq -11, x \neq 0\}$

D) $(f \cdot g)(x) = \sqrt{\frac{15}{x}}$; $\{x \mid x \neq 0\}$

Solve the problem.

12) Given $f(x) = \frac{1}{x}$ and $(\frac{f}{g})(x) = \frac{x+4}{x^2+6x}$, find the function g .

A) $g(x) = \frac{x+6}{x+4}$

B) $g(x) = \frac{x+4}{x+6}$

C) $g(x) = \frac{x-6}{x-4}$

D) $g(x) = \frac{x-4}{x-6}$

13) Find $(f + g)(-4)$ when $f(x) = x - 1$ and $g(x) = x - 3$.

A) -12

B) -10

C) -6

D) -4

14) Find $(f - g)(3)$ when $f(x) = 5x^2 + 2$ and $g(x) = x + 1$.

A) 43

B) -50

C) 45

D) 49

15) Find $(fg)(-2)$ when $f(x) = x + 1$ and $g(x) = 2x^2 + 16x - 3$.

A) 27

B) -15

C) 81

D) 31

16) Find $\left(\frac{f}{g}\right)(-2)$ when $f(x) = 3x - 5$ and $g(x) = 2x^2 + 14x + 2$.

A) $\frac{11}{18}$

B) $-\frac{1}{9}$

C) 2

D) 0

Find and simplify the difference quotient of f , $\frac{f(x+h) - f(x)}{h}$, $h \neq 0$, for the function.

17) $f(x) = 7x - 1$

A) 7

B) $7 + \frac{-2}{h}$

C) $7 + \frac{14(x-1)}{h}$

D) 0

18) $f(x) = x^2 + 6x - 1$

A) $2x + h + 6$

B) $\frac{2x^2 + 2x + 2xh + h^2 + h - 2}{h}$

C) $2x + h - 1$

D) 1

19) $f(x) = \frac{1}{8x}$

A) $\frac{-1}{8x(x+h)}$

B) $\frac{-1}{x(x+h)}$

C) $\frac{1}{8x}$

D) 0

Solve the problem.

20) Suppose that $P(x)$ represents the percentage of income spent on housing in year x and $I(x)$ represents income in year x . Determine a function H that represents total housing expenditures in year x .


A) $H(x) = (P \cdot I)(x)$

B) $H(x) = (P + I)(x)$

C) $H(x) = (I - P)(x)$

D) $H(x) = \left(\frac{I}{P}\right)(x)$

- 21) The following graph shows the private, public and total national school enrollment for students for select years from 1970 through 2000.



- i) How is the graph for total school enrollment, T , determined from the graph of the private enrollment, r , and the public enrollment, u ?
- ii) During which 10-year period did the total number of students enrolled increase the least?
- iii) During which 10-year period did the total number of students enrolled increase the most?
- A) i) T is the sum of r and u .
 ii) 1970 - 1980
 iii) 1990-2000
- B) i) T is the sum of r and u .
 ii) 1990-2000
 iii) 1970-1980
- C) i) T is the sum of r and u .
 ii) 1970 - 1980
 iii) 1980-1990
- D) i) T is the difference of r and u .
 ii) 1970 - 1980
 iii) 1990-2000
- 22) A firm is considering a new product. The accounting department estimates that the total cost, $C(x)$, of producing x units will be

$$C(x) = 95x + 3420.$$
 The sales department estimates that the revenue, $R(x)$, from selling x units will be

$$R(x) = 105x,$$
 but that no more than 734 units can be sold at that price. Find and interpret $(R - C)(734)$.
- A) \$3920 profit, income exceeds cost
 It is worth it to develop product.
- B) -\$3920 loss, cost exceeds income
 It is not worth it to develop product.
- C) \$150,220 profit, income exceeds cost
 It is worth it to develop product.
- D) \$1076 profit, income exceeds cost
 It is worth it to develop product.
- 23) The function $f(t) = -0.13t^2 + 0.52t + 30.5$ models the U.S. population in millions, ages 65 and older, where t represents years after 1990. The function $g(t) = 0.51t^2 + 11.7t + 106.5$ models the total yearly cost of Medicare in billions of dollars, where t represents years after 1990. What does the function $\frac{g}{f}$ represent? Find $\frac{g}{f}(5)$.
- A) Cost per person in thousands of dollars. \$5.95 thousand
- B) Cost per person in thousands of dollars. \$0.20 thousand
- C) Cost per person in thousands of dollars. \$0.17 thousand
- D) Cost per person in thousands of dollars. \$12.18 thousand


2.2 The Graph of a Function

1 Identify the Graph of a Function

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Determine whether the graph is that of a function. If it is, use the graph to find its domain and range, the intercepts, if any, and any symmetry with respect to the x -axis, the y -axis, or the origin.

1)



A) function

domain: $\{x \mid x \leq -2 \text{ or } x \geq 2\}$

range: all real numbers

intercepts: $(-2, 0)$, $(2, 0)$

symmetry: x -axis, y -axis, origin

B) function

domain: all real numbers

range: $\{y \mid y \leq -2 \text{ or } y \geq 2\}$

intercepts: $(-2, 0)$, $(2, 0)$

symmetry: y -axis

C) function

domain: $\{x \mid -2 \leq x \leq 2\}$


range: all real numbers

intercepts: $(-2, 0)$, $(2, 0)$

symmetry: x -axis, y -axis

D) not a function

2)



A) function

domain: $\{x \mid x > 0\}$

range: all real numbers

intercept: $(1, 0)$

symmetry: none

C) function

domain: all real numbers

range: $\{y \mid y > 0\}$

intercept: $(1, 0)$

symmetry: none

B) function

domain: $\{x \mid x > 0\}$


range: all real numbers

intercept: $(0, 1)$

symmetry: origin

D) not a function

3)



A) function

domain: $\{x \mid -\pi \leq x \leq \pi\}$

range: $\{y \mid -1 \leq y \leq 1\}$

intercepts: $(-\pi, 0), (-\frac{3\pi}{4}, 0), (-\frac{\pi}{2}, 0), (-\frac{\pi}{4}, 0), (0, 0), (\frac{\pi}{4}, 0), (\frac{\pi}{2}, 0), (\frac{3\pi}{4}, 0), (\pi, 0)$

symmetry: origin

B) function

domain: $\{x \mid -1 \leq x \leq 1\}$

range: $\{y \mid -\pi \leq y \leq \pi\}$

intercepts: $(-\pi, 0), (-\frac{3\pi}{4}, 0), (-\frac{\pi}{2}, 0), (-\frac{\pi}{4}, 0), (0, 0), (\frac{\pi}{4}, 0), (\frac{\pi}{2}, 0), (\frac{3\pi}{4}, 0), (\pi, 0)$

symmetry: none

C) function

domain: all real numbers


range: $\{y \mid -1 \leq y \leq 1\}$

intercepts: $(-\pi, 0), (-\frac{3\pi}{4}, 0), (-\frac{\pi}{2}, 0), (-\frac{\pi}{4}, 0), (0, 0), (\frac{\pi}{4}, 0), (\frac{\pi}{2}, 0), (\frac{3\pi}{4}, 0), (\pi, 0)$

symmetry: origin

D) not a function

4)




A) function
 domain: all real numbers
 range: $\{y \mid y \leq 9\}$
 intercepts: $(-2, 0)$, $(0, 8)$, $(4, 0)$
 symmetry: none

C) function
 domain: all real numbers
 range: $\{y \mid y \leq 9\}$
 intercepts: $(0, -2)$, $(8, 0)$, $(0, 4)$
 symmetry: none

B) function
 domain: $\{x \mid x \leq 9\}$
 range: all real numbers
 intercepts: $(-2, 0)$, $(0, 8)$, $(4, 0)$
 symmetry: y-axis

D) not a function

5)




A) function
 domain: $\{x \mid -9 \leq x \leq 9\}$
 range: $\{y \mid -9 \leq y \leq 9\}$
 intercepts: $(-9, 0)$, $(0, -9)$, $(0, 9)$, $(9, 0)$
 symmetry: x-axis, y-axis, origin

C) function
 domain: $\{x \mid -9 \leq x \leq 9\}$
 range: $\{y \mid -9 \leq y \leq 9\}$
 intercepts: $(-9, 0)$, $(0, -9)$, $(0, 0)$, $(0, 9)$, $(9, 0)$
 symmetry: origin

B) function
 domain: $\{x \mid -9 \leq x \leq 9\}$
 range: $\{y \mid -9 \leq y \leq 9\}$
 intercepts: $(-9, 0)$, $(0, -9)$, $(0, 9)$, $(9, 0)$
 symmetry: x-axis, y-axis

D) not a function

6)



A) function

domain: $\{x \mid x \geq -2\}$

range: $\{y \mid y \leq 0\}$

intercepts: $(-2, 0), (0, -2), (2, 0)$

symmetry: none

C) function

domain: all real numbers

range: all real numbers

intercepts: $(-2, 0), (0, -2), (2, 0)$

symmetry: none

B) function

domain: $\{x \mid x \leq 0\}$


range: $\{y \mid y \geq -2\}$

intercepts: $(-2, 0), (0, -2), (2, 0)$

symmetry: y-axis

D) not a function

7)



A) function

domain: all real numbers

range: $\{y \mid y = 7 \text{ or } y = 8\}$

intercept: $(0, 8)$

symmetry: none

C) function

domain: all real numbers

range: all real numbers

intercept: $(0, 8)$

symmetry: none

B) function

domain: $\{x \mid x = 7 \text{ or } x = 8\}$

range: all real numbers

intercept: $(8, 0)$

symmetry: x-axis


D) not a function

2 Obtain Information from or about the Graph of a Function

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

The graph of a function f is given. Use the graph to answer the question.

- 1) Use the graph of f given below to find $f(50)$.




A) 0

B) 50

C) 100

D) 60


- 2) Is $f(-25)$ positive or negative?



A) positive

B) negative


3) Is $f(40)$ positive or negative?



A) positive

B) negative

4) For what numbers x is $f(x) = 0$?




A) -60, 70, 100

B) $(-100, -60)$, $(70, 100)$

C) $(-60, 70)$

D) -60

5) For what numbers x is $f(x) > 0$?




A) $[-20, -12), (14, 20)$

B) $(-12, 14)$

C) $(-12, \infty)$

D) $(-\infty, -12)$

6) For what numbers x is $f(x) < 0$?




A) $(-60, 70)$

B) $[-100, -60), (70, 100)$

C) $(-60, \infty)$


D) $(-\infty, -60)$

7) What is the domain of f ?




- A) $\{x \mid -20 \leq x \leq 20\}$ B) $\{x \mid -16 \leq x \leq 22\}$ C) all real numbers D) $\{x \mid x \geq 0\}$

8) What are the x-intercepts?




- A) -6, 7, 10 B) -10, -6, 7, 10 C) -6, 7 D) -6

9) What is the y-intercept?




- A) -30 B) 35 C) 50 D) -40

10) How often does the line $y = -5$ intersect the graph?




- A) once B) twice C) three times D) does not intersect

11) How often does the line $y = 4$ intersect the graph?



- A) once B) twice C) three times D) does not intersect

12) For which of the following values of x does $f(x) = 40$?



- A) 80 B) 140 C) 100 D) 40

Answer the question about the given function.

13) Given the function $f(x) = 7x^2 + 14x - 9$, is the point $(-1, -16)$ on the graph of f ?

- A) Yes B) No

14) Given the function $f(x) = -6x^2 + 12x + 7$, is the point $(2, -5)$ on the graph of f ?

- A) Yes B) No

15) Given the function $f(x) = 5x^2 + 10x + 2$, if $x = -1$, what is $f(x)$? What point is on the graph of f ?

- A) -3; $(-1, -3)$ B) -3; $(-3, -1)$ C) 17; $(-1, 17)$ D) 17; $(17, -1)$





- 16) Given the function $f(x) = -6x^2 - 12x + 1$, what is the domain of f ?
- A) all real numbers B) $\{x \mid x \geq -1\}$ C) $\{x \mid x \leq -1\}$ D) $\{x \mid x \geq 1\}$
- 17) Given the function $f(x) = x^2 + 4x - 21$, list the x -intercepts, if any, of the graph of f .
- A) $(-7, 0), (3, 0)$ B) $(7, 0), (3, 0)$ C) $(-7, 0), (1, 0)$ D) $(7, 0), (-3, 0)$
- 18) Given the function $f(x) = -5x^2 - 10x + 8$, list the y -intercept, if there is one, of the graph of f .
- A) 8 B) -2 C) 13 D) -7
- 19) Given the function $f(x) = \frac{x^2 - 6}{x + 1}$, is the point $(2, -\frac{2}{3})$ on the graph of f ?
- A) Yes B) No
- 20) Given the function $f(x) = \frac{x^2 - 2}{x - 3}$, is the point $(-2, -\frac{6}{5})$ on the graph of f ?
- A) Yes B) No
- 21) Given the function $f(x) = \frac{x^2 - 4}{x + 2}$, if $x = 1$, what is $f(x)$? What point is on the graph of f ?
- A) -1; $(1, -1)$ B) -1; $(-1, 1)$ C) $\frac{5}{3}; (1, \frac{5}{3})$ D) $\frac{5}{3}; (\frac{5}{3}, 1)$
- 22) Given the function $f(x) = \frac{x^2 + 4}{x + 9}$, what is the domain of f ?
- A) $\{x \mid x \neq -9\}$ B) $\{x \mid x \neq 9\}$ C) $\{x \mid x \neq 4\}$ D) $\{x \mid x \neq -\frac{4}{9}\}$
- 23) Given the function $f(x) = \frac{x^2 + 6}{x - 7}$, list the x -intercepts, if any, of the graph of f .
- A) $(6, 0), (-6, 0)$ B) $(7, 0)$ C) $(-\sqrt{6}, 0)$ D) none
- 24) Given the function $f(x) = \frac{x^2 + 5}{x + 7}$, list the y -intercept, if there is one, of the graph of f .
- A) $(0, \frac{5}{7})$ B) $(\frac{5}{7}, 0)$ C) $(0, -7)$ D) $(0, -5)$

Solve the problem.

- 25) If an object weighs m pounds at sea level, then its weight W (in pounds) at a height of h miles above sea level is given approximately by $W(h) = m \left(\frac{4000}{4000 + h} \right)^2$. How much will a man who weighs 165 pounds at sea level weigh on the top of a mountain which is 14,494 feet above sea level? Round to the nearest hundredth of a pound, if necessary.
- A) 164.77 pounds B) 7.72 pounds C) 165.23 pounds D) 165 pounds


Match the function with the graph that best describes the situation.

26) The amount of rainfall as a function of time, if the rain fell more and more softly.




27) The height of an animal as a function of time.


A)




B)



C)




D)



SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.


Solve the problem.

28) Michael decides to walk to the mall to do some errands. He leaves home, walks 2 blocks in 8 minutes at a constant speed, and realizes that he forgot his wallet at home. So Michael runs back in 7 minutes. At home, it takes him 3 minutes to find his wallet and close the door. Michael walks 4 blocks in 12 minutes and then decides to jog to the mall. It takes him 7 minutes to get to the mall which is 2 blocks away. Draw a graph of Michael's distance from home (in blocks) as a function of time.



MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

- 29) A steel can in the shape of a right circular cylinder must be designed to hold 700 cubic centimeters of juice (see figure). It can be shown that the total surface area of the can (including the ends) is given by $S(r) = 2\pi r^2 + \frac{1400}{r}$, where r is the radius of the can in centimeters. Using the TABLE feature of a graphing utility, find the radius that minimizes the surface area (and thus the cost) of the can. Round to the nearest tenth of a centimeter.



- A) 4.8 cm B) 6 cm C) 4 cm D) 0 cm
- 30) The concentration C (arbitrary units) of a certain drug in a patient's bloodstream can be modeled using $C(t) = \frac{t}{(0.62t + 1.612)^2}$, where t is the number of hours since a 500 milligram oral dose was administered. Using the TABLE feature of a graphing utility, find the time at which the concentration of the drug is greatest. Round to the nearest tenth of an hour.
- A) 2.6 hours B) 4.1 hours C) 3.4 hours D) 4.9 hours


2.3 Properties of Functions

1 Determine Even and Odd Functions from a Graph

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.


The graph of a function is given. Decide whether it is even, odd, or neither.

1)



- A) even B) odd C) neither

2)




A) even

B) odd

C) neither

3)




A) even

B) odd

C) neither

4)




A) even

B) odd

C) neither

5)




A) even

B) odd

C) neither

6)




A) even

B) odd

C) neither

7)




A) even

B) odd

C) neither

8)



A) even

B) odd

C) neither

2 Identify Even and Odd Functions from the Equation

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Determine algebraically whether the function is even, odd, or neither.

1) $f(x) = -2x^3$

A) even

B) odd

C) neither

2) $f(x) = 4x^4 - x^2$

A) even

B) odd

C) neither

3) $f(x) = -3x^2 - 2$

A) even

B) odd

C) neither

4) $f(x) = 4x^3 - 7$

A) even

B) odd

C) neither

5) $f(x) = \sqrt[3]{x}$

A) even

B) odd

C) neither

6) $f(x) = \sqrt{x}$

A) even

B) odd

C) neither

7) $\sqrt[3]{7x^2 + 8}$

A) even

B) odd

C) neither

8) $f(x) = \frac{1}{x^2}$

A) even

B) odd

C) neither

9) $f(x) = \frac{x}{x^2 + 5}$

A) even

B) odd

C) neither

10) $f(x) = \frac{-x^3}{2x^2 - 9}$

A) even

B) odd

C) neither

11) $f(x) = \frac{2x}{|x|}$

A) even

B) odd


C) neither

3 Use a Graph to Determine Where a Function Is Increasing, Decreasing, or Constant

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

The graph of a function is given. Determine whether the function is increasing, decreasing, or constant on the given interval.

1) $(-5, -\frac{5}{2})$




A) decreasing

B) increasing

C) constant

2) $(-\frac{3}{2}, 0)$




A) increasing

B) decreasing

C) constant

3) (0, 1)




A) decreasing

B) increasing

C) constant

4) (1, 5)




A) increasing

B) decreasing

C) constant

5) (0, 3)




A) constant

B) increasing

C) decreasing

6) $(-1, 0)$




A) increasing

B) decreasing

C) constant

7) $(3, \infty)$




A) decreasing

B) increasing

C) constant

8) $(0, \infty)$




A) increasing

B) decreasing

C) constant

9) (1, 2)




A) increasing

B) decreasing

C) constant

10) (4, 3)




A) decreasing

B) constant

C) increasing

11) (-1, 0)




A) constant

B) decreasing

C) increasing

12) $(-2.5, 2.2)$




A) increasing

B) decreasing

C) constant

Use the graph to find the intervals on which it is increasing, decreasing, or constant.

13)




A) Increasing on $(-\infty, 0)$; decreasing on $(0, \infty)$

B) Decreasing on $(-\infty, 0)$; increasing on $(0, \infty)$

C) Decreasing on $(-\infty, \infty)$

D) Increasing on $(-\infty, \infty)$


14)



- A) Increasing on $(-\infty, \infty)$
 C) Decreasing on $(-\infty, \infty)$


- B) Decreasing on $(-\infty, 0)$; increasing on $(0, \infty)$
 D) Increasing on $(-\infty, 0)$; decreasing on $(0, \infty)$

15)



- A) Decreasing on $\left(-\pi, -\frac{\pi}{2}\right)$ and $\left(\frac{\pi}{2}, \pi\right)$; increasing on $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
 B) Increasing on $\left(-\pi, -\frac{\pi}{2}\right)$ and $\left(\frac{\pi}{2}, \pi\right)$; decreasing on $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
 C) Decreasing on $(-\pi, 0)$; increasing on $(0, \pi)$
 D) Increasing on $(-\infty, \infty)$

16)




- A) Decreasing on $(-3, -2)$ and $(2, 4)$; increasing on $(-1, 1)$; constant on $(-2, -1)$ and $(1, 2)$
- B) Decreasing on $(-3, -2)$ and $(2, 4)$; increasing on $(-1, 1)$
- C) Decreasing on $(-3, -1)$ and $(1, 4)$; increasing on $(-2, 1)$
- D) Increasing on $(-3, -2)$ and $(2, 4)$; decreasing on $(-1, 1)$; constant on $(-2, -1)$ and $(1, 2)$

4 Use a Graph to Locate Local Maxima and Local Minima

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.


The graph of a function f is given. Use the graph to answer the question.

- 1) Find the numbers, if any, at which f has a local maximum. What are the local maxima?




- A) f has a local maximum at $x = 0$; the local maximum is 3
- B) f has a local maximum at $x = -1$ and 1; the local maximum is 0
- C) f has a local maximum at $x = 1$; the local maximum is 3
- D) f has no local maximum

2) Find the numbers, if any, at which f has a local minimum. What are the local minima?




- A) f has a local minimum at $x = -3$ and 3 ; the local minimum is 0
- B) f has a local minimum at $x = 0$; the local minimum is 1
- C) f has a local minimum at $x = -3$; the local minimum is 0
- D) f has no local minimum

3) Find the numbers, if any, at which f has a local maximum. What are the local maxima?




- A) f has a local maximum at $x = 0$; the local maximum is 1
- B) f has a local maximum at $x = -\pi$ and π ; the local maximum is -1
- C) f has a local maximum at $-\pi$; the local maximum is 1
- D) f has no local maximum

4) Find the numbers, if any, at which f has a local minimum. What are the local minima?



- A) f has a local minimum at $x = 0$; the local minimum is -2
- B) f has a local minimum at $x = -\pi$ and π ; the local minimum is 2
- C) f has a local minimum at $x = -\pi$; the local minimum is -2
- D) f has no local minimum

5)



Find the numbers, if any, at which f has a local maximum. What are the local maxima?

- A) f has a local maximum at $x = -8$ and 2.2 ; the local maximum at -8 is 5 ; the local maximum at 2.2 is 3.9
- B) f has a local maximum at $x = 5$ and 3.9 ; the local maximum at 5 is -8 ; the local maximum at 3.9 is 2.2
- C) f has a local minimum at $x = -8$ and 2.2 ; the local minimum at -8 is 5 ; the local minimum at 2.2 is 3.9
- D) f has a local minimum at $x = 5$ and 3.9 ; the local minimum at 5 is -8 ; the local minimum at 3.9 is 2.2

5 Use Graphing Utility to Approx Local Maxima and Local Minima and Determine Where a Func Is Incrs or Decrs

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Use a graphing utility to graph the function over the indicated interval and approximate any local maxima and local minima. Determine where the function is increasing and where it is decreasing. If necessary, round answers to two decimal places.

1) $f(x) = x^3 - 3x + 3$; $(-2, 2)$

A) local maximum at $(-1, 5)$
local minimum at $(1, 1)$
increasing on $(-2, -1)$ and $(1, 2)$
decreasing on $(-1, 1)$

C) local maximum at $(-1, 5)$
local minimum at $(1, 1)$
increasing on $(-1, 1)$
decreasing on $(-2, -1)$ and $(1, 2)$

B) local maximum at $(1, 1)$
local minimum at $(-1, 5)$
increasing on $(-2, -1)$ and $(1, 2)$
decreasing on $(-1, 1)$

D) local maximum at $(1, 1)$
local minimum at $(-1, 5)$
increasing on $(-2, -1)$
decreasing on $(-1, 1)$

2) $f(x) = x^3 - 4x^2 + 6$; $(-1, 4)$

A) local maximum at $(0, 6)$
local minimum at $(2.67, -3.48)$
increasing on $(-1, 0)$ and $(2.67, 4)$
decreasing on $(0, 2.67)$

C) local maximum at $(0, 6)$
local minimum at $(2.67, -3.48)$
increasing on $(0, 2.67)$
decreasing on $(-1, 0)$ and $(2.67, 4)$

B) local maximum at $(2.67, -3.48)$
local minimum at $(0, 6)$
increasing on $(-1, 0)$ and $(2.67, 4)$
decreasing on $(0, 2.67)$

D) local maximum at $(2.67, -3.48)$
local minimum at $(0, 6)$
increasing on $(0, 2.67)$
decreasing on $(-1, 0)$ and $(2.67, 4)$

3) $f(x) = x^5 - x^2$; $(-2, 2)$

A) local maximum at $(0, 0)$
local minimum at $(0.74, -0.33)$
increasing on $(-2, 0)$ and $(0.74, 2)$
decreasing on $(0, 0.74)$

C) local maximum at $(0, 0)$
local minimum at $(0.74, -0.33)$
increasing on $(0, 0.74)$
decreasing on $(-2, 0)$ and $(0.74, 2)$

B) local maximum at $(0.74, -0.33)$
local minimum at $(0, 0)$
increasing on $(-2, 0)$ and $(0.74, 2)$
decreasing on $(0, 0.74)$

D) local maximum at $(0.74, -0.33)$
local minimum at $(0, 0)$
increasing on $(0, 0.74)$
decreasing on $(-2, 0)$ and $(0.74, 2)$

4) $f(x) = -0.3x^3 + 0.2x^2 + 4x - 5$; $(-4, 5)$

A) local maximum at $(2.34, 1.61)$
local minimum at $(-1.9, -9.82)$
increasing on $(-1.9, 2.34)$
decreasing on $(-4, -1.9)$ and $(2.34, 5)$

C) local maximum at $(2.34, 1.61)$
local minimum at $(-1.9, -9.82)$
increasing on $(-4, -1.9)$ and $(2.34, 5)$
decreasing on $(-1.9, 2.34)$

B) local maximum at $(-1.9, -9.82)$
local minimum at $(2.34, 1.61)$
increasing on $(-1.9, 2.34)$
decreasing on $(-4, -1.9)$ and $(2.34, 5)$

D) local maximum at $(-1.9, -9.82)$
local minimum at $(2.34, 1.61)$
increasing on $(-4, -1.9)$ and $(2.34, 5)$
decreasing on $(-1.9, 2.34)$

5) $f(x) = 0.15x^4 + 0.3x^3 - 0.8x^2 + 5$; $(-4, 2)$

- A) local maximum at $(0, 5)$
 local minima at $(-2.55, 1.17)$ and $(1.05, 4.65)$
 increasing on $(-2.55, 0)$ and $(1.05, 2)$
 decreasing on $(-4, -2.55)$ and $(0, 1.05)$

- C) local maximum at $(0, 5)$
 local minima at $(-2.55, 1.17)$ and $(1.05, 4.65)$
 increasing on $(-4, -2.55)$ and $(0, 1.05)$
 decreasing on $(-2.55, 0)$ and $(1.05, 2)$

- B) local maximum at $(-2.55, 1.17)$ and $(1.05, 4.65)$
 local minima at $(0, 5)$
 increasing on $(-2.55, 0)$ and $(1.05, 2)$
 decreasing on $(-4, -2.55)$ and $(0, 1.05)$

- D) local maximum at $(-2.55, 1.17)$ and $(1.05, 4.65)$
 local minima at $(0, 5)$
 increasing on $(-4, -2.55)$ and $(0, 1.05)$
 decreasing on $(-2.55, 0)$ and $(1.05, 2)$

Use a graphing utility to graph the function over the indicated interval and approximate any local maxima and local minima. If necessary, round answers to two decimal places.

6) $f(x) = x^2 + 2x - 3$; $(-5, 5)$

- A) local minimum at $(-1, -4)$
 C) local minimum at $(1, 4)$

- B) local maximum at $(-1, 4)$
 D) local maximum at $(1, -4)$

7) $f(x) = 2 + 8x - x^2$; $(-5, 5)$

- A) local maximum at $(4, 18)$
 C) local minimum at $(-4, 18)$

- B) local minimum at $(4, 50)$
 D) local maximum at $(-4, 50)$

8) $f(x) = x^3 - 3x^2 + 1$; $(-5, 5)$

- A) local maximum at $(0, 1)$
 local minimum at $(2, -3)$
 C) local minimum at $(2, -3)$

- B) local minimum at $(0, 1)$
 local maximum at $(2, -3)$
 D) none

9) $f(x) = x^3 - 12x + 2$; $(-5, 5)$

- A) local maximum at $(-2, 18)$
 local minimum at $(2, -14)$
 C) local minimum at $(0, 0)$

- B) local maximum at $(-2, 18)$
 local minimum at $(0, 0)$
 local minimum at $(2, -14)$
 D) none


10) $f(x) = x^4 - 5x^3 + 3x^2 + 9x - 3$; $(-5, 5)$

- A) local minimum at $(-0.57, -6.12)$
 local maximum at $(1.32, 5.64)$
 local minimum at $(3, -3)$
 C) local minimum at $(-3, -3)$
 local maximum at $(-1.32, 5.64)$
 local minimum at $(0.57, -6.12)$

- B) local minimum at $(-1, -6)$
 local maximum at $(1, 6)$
 local minimum at $(3, -3)$
 D) local minimum at $(-0.61, -5.64)$
 local maximum at $(1.41, 6.12)$
 local minimum at $(3, -3)$

Solve the problem.

- 11) The height s of a ball (in feet) thrown with an initial velocity of 70 feet per second from an initial height of 3 feet is given as a function of time t (in seconds) by $s(t) = -16t^2 + 70t + 3$. What is the maximum height? Round to the nearest hundredth, if necessary.



- A) 79.56 ft B) 90.5 ft C) 76.75 ft D) -54.5 ft

Solve.

- 12) John owns a hotdog stand. He has found that his profit is represented by the equation $P(x) = -x^2 + 64x + 76$, with P being profits and x the number of hotdogs sold. How many hotdogs must he sell to earn the most profit?
- A) 32 hotdogs B) 44 hotdogs C) 33 hotdogs D) 22 hotdogs
- 13) Bob owns a watch repair shop. He has found that the cost of operating his shop is given by $c(x) = 3x^2 - 216x + 86$, where c is cost and x is the number of watches repaired. How many watches must he repair to have the lowest cost?
- A) 36 watches B) 30 watches C) 86 watches D) 43 watches
- 14) John owns a hotdog stand. His profit is represented by the equation $P(x) = -x^2 + 14x + 54$, with P being profits and x the number of hotdogs sold. What is the most he can earn?
- A) \$103 B) \$117 C) \$75 D) \$49
- 15) A rock falls from a tower that is 117.6 m high. As it is falling, its height is given by the formula $h(t) = 117.6 - 4.9t^2$. How many seconds will it take for the rock to hit the ground ($h=0$)? Round to the nearest tenth.
- A) 4.9 sec B) 10.8 sec C) 2800 sec D) 24 sec
- 16) A projectile is thrown upward so that its distance above the ground after t seconds is $h(t) = -16t^2 + 532t$. After how many seconds does it reach its maximum height? Round to the nearest second.
- A) 17 sec B) 9 sec C) 28.5 sec D) 38 sec
- 17) A rock falls from a tower that is 176 ft high. As it is falling, its height is given by the formula $h(t) = 176 - 16t^2$. How many seconds will it take for the rock to hit the ground ($h=0$)? Round to the nearest tenth.
- A) 3.3 sec B) 13.3 sec C) 1936 sec D) 10.9 sec

6 Find the Average Rate of Change of a Function

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Find the average rate of change for the function between the given values.

1) $f(x) = -2x + 4$; from 1 to 2

A) -2

B) -4

C) 2

D) 4

2) $f(x) = x^2 + 2x$; from 1 to 6

A) 9

B) $\frac{48}{5}$

C) $\frac{15}{2}$

D) 8

3) $f(x) = 8x^3 + 6x^2 - 2$; from 5 to 8

A) 1110

B) $\frac{4478}{3}$

C) $\frac{1665}{4}$

D) $\frac{2239}{4}$

4) $f(x) = \sqrt{2x}$; from 2 to 8

A) $\frac{1}{3}$

B) 2

C) 7

D) $-\frac{3}{10}$

5) $f(x) = \frac{3}{x-2}$; from 4 to 7

A) $-\frac{3}{10}$

B) 2

C) 7

D) $\frac{1}{3}$

6) $f(x) = 4x^2$; from 0 to $\frac{7}{4}$

A) 7

B) 2

C) $\frac{1}{3}$

D) $-\frac{3}{10}$

7) $f(x) = -3x^2 - x$; from 5 to 6

A) -34

B) -2

C) $\frac{1}{2}$

D) $-\frac{1}{6}$

8) $f(x) = x^3 + x^2 - 8x - 7$; from 0 to 2

A) -2

B) -28

C) $\frac{1}{2}$

D) $-\frac{1}{6}$

9) $f(x) = \sqrt{2x-1}$; from 1 to 5

A) $\frac{1}{2}$

B) -2

C) -28

D) $-\frac{1}{6}$

10) $f(x) = \frac{3}{x+2}$; from 1 to 4

A) $-\frac{1}{6}$

B) -2

C) -28

D) $\frac{1}{2}$

Find an equation of the secant line containing (1, f(1)) and (2, f(2)).

11) $f(x) = x^3 - x$

A) $y = 6x - 6$

B) $y = 6x + 6$

C) $y = -6x - 6$

D) $y = -6x + 6$

12) $f(x) = \frac{6}{x+5}$

A) $y = -\frac{1}{7}x + \frac{8}{7}$

B) $y = \frac{1}{7}x + \frac{6}{7}$

C) $y = \frac{6}{7}x + \frac{1}{7}$

D) $y = \frac{1}{7}x + \frac{4}{3}$

13) $f(x) = \sqrt{x+15}$

A) $y = (\sqrt{17} - 4)x - \sqrt{17} + 8$

B) $y = (-\sqrt{17} + 4)x + \sqrt{17} - 8$

C) $y = (\sqrt{17} - 4)x + \sqrt{17} - 8$

D) $y = (-\sqrt{17} - 4)x - \sqrt{17} + 8$

For the function, find the average rate of change of f from 1 to x:

$\frac{f(x) - f(1)}{x - 1}, x \neq 1$

14) $f(x) = -7x$

A) -7

B) -8

C) $\frac{-7}{x-1}$

D) 0

15) $f(x) = x^3 + x$

A) $x^2 + x + 2$

B) $x^2 + 2$

C) 1

D) $\frac{x^3 + x + 2}{x - 1}$

16) $f(x) = \frac{5}{x+4}$

A) $-\frac{1}{x+4}$

B) $\frac{1}{x+4}$

C) $\frac{5}{(x-1)(x+4)}$

D) $\frac{5}{x(x+4)}$

17) $f(x) = \sqrt{x+24}$

A) $\frac{\sqrt{x+24} - 5}{x-1}$

B) $\frac{\sqrt{x+24} + 5}{x+1}$

C) $\frac{\sqrt{x+24} - 5}{x+1}$

D) $\frac{\sqrt{x+24} + 5}{x-1}$

Solve the problem.

18) From April through December 2000, the stock price of QRS Company had a roller coaster ride. The chart below indicates the price of the stock at the beginning of each month during that period. Find the monthly average rate of change in price between June and September.

Month	Price
April (x = 1)	115
May	109
June	88
July	99
August	94
September	113
October	92
November	85
December	64

A) \$8.33 per month

B) -\$8.33 per month

C) \$12.50 per month

D) -\$12.50 per month

- 19) Along with incomes, people's charitable contributions have steadily increased over the past few years. The table below shows the average deduction for charitable contributions reported on individual income tax returns for the period 1993 to 1998. Find the average rate of change between 1995 and 1997.

Year	Charitable Contributions
1993	\$1780
1994	\$2390
1995	\$2480
1996	\$2840
1997	\$3020
1998	\$3130

- A) \$270 per year B) \$540 per year C) \$325 per year D) \$315 per year
- 20) A deep sea diving bell is being lowered at a constant rate. After 11 minutes, the bell is at a depth of 600 ft. After 50 minutes the bell is at a depth of 2000 ft. What is the average rate of lowering per minute? Round to the nearest hundredth is needed.
- A) 35.9 ft per minute B) 0.03 ft per minute C) 28.0 ft per minute D) 40.0 ft per minute

Find and simplify the difference quotient $\frac{f(x+h) - f(x)}{h}$, $h \neq 0$ for the given function.

21) $f(x) = 6x - 5$

A) 6

B) $6 + \frac{-10}{h}$

C) $6 + \frac{12(x-5)}{h}$

D) 0

22) $f(x) = 8x^2$

A) $8(2x+h)$

B) $\frac{16}{h} + x + 8h$

C) $\frac{8(2x^2 + 2xh + h^2)}{h}$

D) 8

23) $f(x) = 4$

A) 0

B) 1

C) $1 + \frac{8}{h}$

D) 4

24) $f(x) = \frac{1}{2x}$

A) $\frac{-1}{2x(x+h)}$

B) $\frac{-1}{x(x+h)}$

C) $\frac{1}{2x}$

D) 0

25) $f(x) = x^2 + 6x + 7$

A) $2x + h + 6$

B) $\frac{2x^2 + 2x + 2xh + h^2 + h + 14}{h}$

C) $2x + h + 7$

D) 1


2.4 Library of Functions; Piecewise-defined Functions

1 Graph the Functions Listed in the Library of Functions

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Match the graph to the function listed whose graph most resembles the one given.

1)



A) square function

B) cube function

C) absolute value function

D) reciprocal function

2)



A) constant function

B) linear function

C) absolute value function

D) reciprocal function

3)




A) square root function

B) square function

C) cube root function

D) cube function

4)




A) absolute value function

B) square function

C) linear function

D) reciprocal function

5)



A) linear function

C) absolute value function

B) constant function

D) reciprocal function

6)




A) cube function

C) square function

B) cube root function

D) square root function

7)



A) reciprocal function

C) absolute value function

B) square root function

D) square function

8)



A) cube root function


C) square root function

B) cube function


D) square function

Graph the function.


9) $f(x) = x$




A)




B)




C)




D)




10) $f(x) = x^2$




A)




B)




C)




D)




11) $f(x) = x^3$




A)




B)




C)




D)




12) $f(x) = \sqrt{x}$




A)




B)




C)




D)




13) $f(x) = \frac{1}{x}$




A)




B)




C)




D)




14) $f(x) = |x|$




A)




B)




C)




D)




15) $f(x) = \sqrt[3]{x}$




A)




B)




C)




D)




16) $f(x) = 4$




A)




B)



C)



D)




2 Graph Piecewise-defined Functions

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.


Graph the function.

1)


$$f(x) = \begin{cases} x - 4 & \text{if } x < 1 \\ -1 & \text{if } x \geq 1 \end{cases}$$




A)




B)



C)




D)




2)


$$f(x) = \begin{cases} -x + 3 & \text{if } x < 2 \\ 2x - 3 & \text{if } x \geq 2 \end{cases}$$




A)




B)



C)





D)



Solve the problem.

3) Assume it costs 25 cents to mail a letter weighing one ounce or less, and then 20 cents for each additional ounce or fraction of an ounce. Let $L(x)$ be the cost of mailing a letter weighing x ounces. Graph $y = L(x)$.




2.5 Graphing Techniques: Transformations

1 Graph Functions Using Vertical and Horizontal Shifts

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Match the correct function to the graph.

1)




A) $y = \sqrt{x-1}$

B) $y = \sqrt{x}$

C) $y = \sqrt{x+1}$

D) $y = x - 1$

2)



A) $y = |2 - x|$

B) $y = |x + 2|$

C) $y = |1 - x|$

D) $y = x - 2$

Write an equation that results in the indicated translation.

3) The squaring function, shifted 5 units upward

A) $y = x^2 + 5$

B) $y = x^2 - 5$

C) $y = \frac{x^2}{5}$

D) $y = 5x^2$

4) The absolute value function, shifted 5 units to the right

A) $y = |x - 5|$

B) $y = |x + 5|$

C) $y = |x| - 5$

D) $y = |x| + 5$

5) The absolute value function, shifted 7 units upward

A) $y = |x| + 7$

B) $y = |x - 7|$

C) $y = |x + 7|$

D) $y = |x| - 7$

6) The square root function, shifted 9 units to the right

A) $y = \sqrt{x - 9}$

B) $y = \sqrt{x + 9}$

C) $y = \sqrt{x} + 9$

D) $y = \sqrt{x} - 9$

7) The square root function, shifted 7 units to the left

A) $y = \sqrt{x + 7}$

B) $y = \sqrt{x - 7}$

C) $y = \sqrt{x} + 7$

D) $y = \sqrt{x} - 7$

8) The square root function, shifted 7 units upward

A) $y = \sqrt{x} + 7$

B) $y = \sqrt{x - 7}$

C) $y = \sqrt{x + 7}$

D) $y = \sqrt{x} - 7$

9) The square root function, shifted 8 units downward

A) $y = \sqrt{x} - 8$

B) $y = \sqrt{x - 8}$

C) $y = \sqrt{x + 8}$

D) $y = \sqrt{x} + 8$

Suppose the point (2, 4) is on the graph of $y = f(x)$. Find a point on the graph of the given function.

10) $y = f(x + 1)$

A) (1, 4)

B) (3, 4)

C) (2, 3)

D) (2, 5)

11) $f(x) + 4$

A) (2, 8)


B) (2, -4)

C) (6, 4)


D) (-2, 4)

Graph the function by starting with the graph of the basic function and then using the techniques of shifting, compressing, stretching, and/or reflecting.


12) $f(x) = x^2 + 5$




A)




B)




C)




D)




13) $f(x) = (x + 4)^2$




A)




B)




C)




D)




14) $f(x) = (x - 3)^2 + 5$




A)




B)




C)




D)




15) $f(x) = x^3 - 2$




A)




B)




C)




D)




16) $f(x) = (x - 2)^3$




A)




B)




C)




D)




17) $f(x) = (x + 6)^3 + 1$




A)




B)




C)




D)




18) $f(x) = \sqrt{x} - 4$




A)




B)




C)




D)




19) $f(x) = \sqrt{x-2}$




A)




B)




C)




D)




20) $f(x) = \sqrt{x+4} - 3$




A)




B)




C)




D)




21) $f(x) = \sqrt{x-3} + 5$




A)




B)




C)




D)




22) $f(x) = |x| + 5$




A)




B)




C)




D)




23) $f(x) = |x + 1|$




A)




B)




C)




D)




24) $f(x) = |x + 7| + 7$




A)




B)




C)




D)




25) $f(x) = \frac{1}{x} + 2$




A)




B)




C)




D)




26) $f(x) = \frac{1}{x+3}$




A)




B)




C)




D)




$$27) f(x) = \frac{1}{x-1} - 1$$




A)




B)



C)




D)




Using transformations, sketch the graph of the requested function.


28) The graph of a function f is illustrated. Use the graph of f as the first step toward graphing the function $F(x)$, where $F(x) = f(x + 2) - 1$.




A)




B)



C)



D)



Solve the problem.

29) Suppose that the x -intercepts of the graph of $y = f(x)$ are 9 and 5. What are the x -intercepts of $y = f(x + 8)$?

A) 1 and -3

B) 17 and 13

C) 72 and 40

D) 9 and 13

30) Suppose that the x -intercepts of the graph of $y = f(x)$ are 9 and 7. What are the x -intercepts of $y = f(x - 3)$?

A) 12 and 10

B) 6 and 4

C) 27 and 21

D) 9 and 4

31) Suppose that the function $y = f(x)$ is increasing on the interval $(8, 9)$. Over what interval is the graph of $y = f(x + 7)$ increasing?

A) $(1, 2)$

B) $(15, 16)$

C) $(56, 63)$

D) $(8, 9)$

32) Suppose that the function $y = f(x)$ is increasing on the interval $(2, 6)$. Over what interval is the graph of $y = f(x - 4)$ increasing?

A) $(6, 10)$


B) $(-2, 2)$

C) $(8, 24)$


D) $(2, 6)$

Complete the square and then use the shifting technique to graph the function.


33) $f(x) = x^2 + 10x$




A)




B)




C)




D)




34) $f(x) = x^2 + 2x - 2$




A)




B)



C)



D)



Solve the problem.

- 35) The following numerical representation for f computes the average number of hours of television watched per day based on year of birth x .

x	1975	1980	1983	1988	1990	1992	1995
$f(x)$	2	2.5	3	3.5	4	3.5	4

Give a numerical representation for a function g that computes the average number of hours of television watched per day for the year x , where $x = 0$ corresponds to the birth year 1975. Write an equation that shows the relationship between $f(x)$ and $g(x)$.

A)

x	0	5	8	13	15	17	20
$g(x)$	2	2.5	3	3.5	4	3.5	4

 $f(x) = g(x - 1975)$

B)

x	75	80	83	88	90	92	95
$g(x)$	2	2.5	3	3.5	4	3.5	4

 $f(x) = g(x - 1900)$

C)

x	0	5	8	13	15	17	20
$g(x)$	2	2.5	3	3.5	4	3.5	4


 $f(x) = g(x + 1975)$

D)

x	0	5	8	13	15	17	20
$g(x)$	2	2.5	3	3.5	4	3.5	4

 $f(x) = g(x) - 1975$



- 36) Suppose a cold front is passing through the United States at noon with a shape described by the function $y = \frac{1}{20}x^2$, where each unit represents 100 miles. St. Louis, Missouri is located at $(0, 0)$, and the positive y -axis points north.




Suppose the front moves south 340 miles and west 120 miles and maintains its shape. Give the equation for the new front and plot the new position of the front.

A) $y = \frac{1}{20}(x + 1.2)^2 - 3.4$


B) $y = \frac{1}{20}(x - 1.2)^2 - 3.4$



$$C) y = \frac{1}{20}(x - 1.2)^2 + 3.4$$



$$D) y = -\frac{1}{20}(x + 1.2)^2 - 3.4$$



2 Graph Functions Using Compressions and Stretches

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Write the equation that results in the desired transformation.

1) The squaring function, vertically stretched by a factor of 10

A) $y = 10x^2$

B) $y = -10x^2$

C) $y = (x - 10)^2$

D) $y = 10(x - 10)x^2$

2) The cubing function, vertically compressed by a factor of .3

A) $y = .3x^3$

B) $y = .3\sqrt[3]{x}$

C) $y = (x - .3)^3$

D) $y = (x + .3)^3$

Suppose the point (2, 4) is on the graph of $y = f(x)$. Find a point on the graph of the given function.

3) $y = 2f(x)$

A) (2, 8)


B) (4, 4)

C) (1, 4)


D) (5, 1)

Graph the function by starting with the graph of the basic function and then using the techniques of shifting, compressing, stretching, and/or reflecting.


4) $f(x) = 7x^2$




A)




B)




C)




D)




5) $f(x) = \frac{1}{5}x^2$




A)




B)




C)




D)




6) $f(x) = 2x^3$




A)




B)




C)




D)




7) $f(x) = \frac{1}{3}x^3$




A)




B)




C)




D)




8) $f(x) = 4\sqrt{x}$




A)




B)




C)




D)




9) $f(x) = \frac{1}{2}\sqrt{x}$




A)




B)




C)




D)




10) $f(x) = 2|x|$




A)




B)




C)




D)




11) $f(x) = \frac{1}{2}|x|$




A)




B)




C)




D)




12) $f(x) = \frac{2}{x}$




A)




B)




C)




D)




13) $f(x) = \frac{1}{3x}$




A)




B)




C)




D)




14) $f(x) = 3(x + 1)^2 + 3$




A)




B)



C)





D)




Use the accompanying graph of $y = f(x)$ to sketch the graph of the indicated equation.


15) $y = 2f(x)$




A)




B)





C)




D)




16) $y = -\frac{1}{2}f(x)$




A)




B)





C)




D)




17) $y = -\frac{1}{3}f(x+2) + 5$




A)




B)



C)



D)



Solve the problem.

18) Suppose that the x-intercepts of the graph of $y = f(x)$ are 9 and 5. What are the x-intercepts of $y = 6f(x)$?

A) 9 and 5

B) 3 and -1

C) 15 and 11


D) 45 and 30

3 Graph Functions Using Reflections about the x-Axis and the y-Axis

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Match the correct function to the graph.

1)



A) $y = -2x^2 + 1$

B) $y = -2x^2$

C) $y = -2x^2 - 1$

D) $y = 1 - x^2$

Find the function.

2) Find the function that is finally graphed after the following transformations are applied to the graph of $y = |x|$. The graph is shifted right 3 units, stretched by a factor of 3, shifted vertically down 2 units, and finally reflected across the x-axis.

A) $y = -(3|x - 3| - 2)$

B) $y = -3|x - 3| - 2$

C) $y = -(3|x + 3| - 2)$

D) $y = 3|-x - 3| - 2$

3) Find the function that is finally graphed after the following transformations are applied to the graph of $y = \sqrt{x}$. The graph is shifted up 8 units, reflected about the x-axis, and finally shifted right 5 units.

A) $y = -\sqrt{x - 5} - 8$

B) $y = -\sqrt{x - 5} + 8$

C) $y = \sqrt{-x + 5} - 8$

D) $y = -\sqrt{x + 5} + 8$

4) Find the function that is finally graphed after the following transformations are applied to the graph of $y = \sqrt{x}$. The graph is shifted up 8 units, reflected about the x-axis, and finally shifted right 5 units.

A) $y = -\sqrt{x - 5} - 8$

B) $y = -\sqrt{x - 5} + 8$

C) $y = \sqrt{-x + 5} - 8$

D) $y = -\sqrt{x + 5} + 8$

Suppose the point (2, 4) is on the graph of $y = f(x)$. Find a point on the graph of the given function.

5) The reflection of the graph of $y = f(x)$ across the x-axis

A) (2, -4)

B) (-2, -4)

C) (2, 4)

D) (-2, 4)

6) The reflection of the graph of $y = f(x)$ across the y-axis

A) (-2, 4)


B) (2, 4)

C) (-2, -4)


D) (2, -4)

Graph the function by starting with the graph of the basic function and then using the techniques of shifting, compressing, stretching, and/or reflecting.


7) $f(x) = -x^2$




A)




B)




C)




D)




8) $f(x) = (-x)^2$




A)




B)




C)




D)




9) $f(x) = -x^3$




A)




B)




C)




D)




10) $f(x) = (-x)^3$




A)




B)




C)




D)




11) $f(x) = -\sqrt{x}$




A)




B)




C)




D)




12) $f(x) = \sqrt{-x}$




A)




B)




C)




D)




13) $f(x) = -|x|$




A)




B)




C)




D)




14) $f(x) = |-x|$




A)




B)




C)




D)




15) $f(x) = -\frac{1}{x}$




A)




B)




C)




D)




16) $f(x) = -(x + 6)^2 - 2$




A)




B)




C)




D)




17) $f(x) = -2(x + 1)^2 + 2$




A)




B)



C)



D)



Solve the problem.

18) Suppose that the x-intercepts of the graph of $y = f(x)$ are 9 and 5. What are the x-intercepts of $y = f(-x)$?

A) -9 and -5

B) 9 and 5

C) 9 and -5


D) -9 and 5

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

- 8) A right triangle has one vertex on the graph of $y = x^2$ at (x, y) , another at the origin, and the third on the (positive) y -axis at $(0, y)$. Express the area A of the triangle as a function of x .

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

- 9) The figure shown here shows a rectangle inscribed in an isosceles right triangle whose hypotenuse is 4 units long. Express the area A of the rectangle in terms of x .



- A) $A(x) = 2x(2 - x)$ B) $A(x) = x(2 - x)$ C) $A(x) = 2x(x - 2)$ D) $A(x) = 2x^2$

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

- 10) A wire 20 feet long is to be cut into two pieces. One piece will be shaped as a square and the other piece will be shaped as an equilateral triangle. Express the total area A enclosed by the pieces of wire as a function of the length x of a side of the equilateral triangle. What is the domain of A ?

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

- 11) A farmer's silo is the shape of a cylinder with a hemisphere as the roof. If the height of the silo is 105 feet and the radius of the hemisphere is r feet, express the volume of the silo as a function of r .

- A) $V(r) = \pi(105 - r)r^2 + \frac{2}{3} \pi r^3$ B) $V(r) = 105\pi r^2 + \frac{8}{3} \pi r^3$
 C) $V(r) = \pi(105 - r)r^3 + \frac{4}{3} \pi r^2$ D) $V(r) = \pi(105 - r) + \frac{4}{3} \pi r^2$

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

- 12) The volume V of a square-based pyramid with base sides s and height h is $V = \frac{1}{3}s^2h$. If the height is half of the length of a base side, express the volume V as a function of s .

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.


- 13) A farmer's silo is the shape of a cylinder with a hemisphere as the roof. If the radius of the hemisphere is 10 feet and the height of the silo is h feet, express the volume of the silo as a function of h .

A) $V(h) = 100 \pi(h - 10) + \frac{2000}{3} \pi$ B) $V(h) = 100 \pi h + \frac{4000}{3} \pi h^2$
 C) $V(h) = 100 \pi(h^2 - 10) + \frac{5000}{3} \pi$ D) $V(h) = 4100 \pi(h - 10) + \frac{500}{7} \pi$

- 14) From a 40-inch by 40-inch piece of metal, squares are cut out of the four corners so that the sides can then be folded up to make a box. Let x represent the length of the sides of the squares, in inches, that are cut out. Express the volume of the box as a function of x .

A) $V(x) = 4x^3 - 160x^2 + 1600x$ B) $V(x) = 2x^3 - 120x^2$
 C) $V(x) = 4x^3 - 160x^2$ D) $V(x) = 2x^3 - 120x^2 + 40x$

- 15) A box with an open top is to be constructed from a rectangular piece of cardboard with dimensions 11 inches by 21 inches by cutting out equal squares of side x at each corner and then folding up the sides as in the figure. Express the volume V of the box as a function of x .



A) $V(x) = x(11 - 2x)(21 - 2x)$ B) $V(x) = (11 - 2x)(21 - 2x)$
 C) $V(x) = x(11 - x)(21 - x)$ D) $V(x) = (11 - x)(21 - x)$

- 16) A rectangular box with volume 321 cubic feet is built with a square base and top. The cost is \$1.50 per square foot for the top and the bottom and \$2.00 per square foot for the sides. Let x represent the length of a side of the base. Express the cost the box as a function of x .

A) $C(x) = 3x^2 + \frac{2568}{x}$ B) $C(x) = 3x^2 + \frac{1284}{x}$ C) $C(x) = 2x^2 + \frac{2568}{x}$ D) $C(x) = 4x + \frac{2568}{x^2}$

- 17) The price p and the quantity x sold of a certain product obey the demand equation:

$$p = -\frac{1}{6}x + 200, \{x | 0 \leq x \leq 600\}$$

What is the revenue to the nearest dollar when 200 units are sold?

A) \$33,333 B) \$46,667 C) \$40,000 D) \$100,000

- 18) Let $P = (x, y)$ be a point on the graph of $y = \sqrt{x}$. Express the distance d from P to the point $(1, 0)$ as a function of x .

A) $d(x) = x^2 - x + 1$ B) $d(x) = \sqrt{x^2 + 2x + 2}$
 C) $d(x) = x^2 + 2x + 2$ D) $d(x) = \sqrt{x^2 - x + 1}$

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

- 19) The price p and x , the quantity of a certain product sold, obey the demand equation

$$p = -\frac{1}{10}x + 100, \{x \mid 0 \leq x \leq 1000\}$$

- Express the revenue R as a function of x .
 - What is the revenue if 450 units are sold?
 - Graph the revenue function using a graphing utility.
 - What quantity x maximizes revenue? What is the maximum revenue?
 - What price should the company charge to maximize revenue?
- 20) Two boats leave a dock at the same time. One boat is headed directly east at a constant speed of 35 knots (nautical miles per hour), and the other is headed directly south at a constant speed of 22 knots. Express the distance d between the boats as a function of the time t .

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

- 21) A rocket is shot straight up in the air from the ground at a rate of 55 feet per second. The rocket is tracked by a range finder that is 454 feet from the launch pad. Let d represent the distance from the rocket to the range finder and t represent the time, in seconds, since "blastoff". Express d as a function of t .

A) $d(t) = \sqrt{454^2 + (55t)^2}$

B) $d(t) = 454^2 + (55t)^2$

C) $d(t) = \sqrt{55^2 + (454t)^2}$

D) $d(t) = 454 + 55t^2$

Ch. 2 Functions and Their Graphs

Answer Key

2.1 Functions

1 Determine Whether a Relation Represents a Function

- 1) A
- 2) C
- 3) A
- 4) A
- 5) C
- 6) A
- 7) A
- 8) A
- 9) A
- 10) A
- 11) B
- 12) B
- 13) B
- 14) B
- 15) A
- 16) A
- 17) B
- 18) A
- 19) A

2 Find the Value of a Function

- 1) A
- 2) A
- 3) A
- 4) A
- 5) A
- 6) A
- 7) A
- 8) A
- 9) A
- 10) A
- 11) A
- 12) A
- 13) A
- 14) A
- 15) A
- 16) A
- 17) A
- 18) A
- 19) A
- 20) A
- 21) A
- 22) A

3 Find the Domain of a Function

- 1) A
- 2) A
- 3) A
- 4) A
- 5) A
- 6) A

7) A

4 Form the Sum, Difference, Product, and Quotient of Two Functions

1) A

2) A

3) A

4) A

5) A

6) A

7) A

8) A

9) A

10) A

11) A

12) A

13) A

14) A

15) A

16) A

17) A

18) A

19) A

20) A

21) A

22) A

23) A

2.2 The Graph of a Function

1 Identify the Graph of a Function

1) D

2) A

3) A

4) A

5) D

6) A

7) D

2 Obtain Information from or about the Graph of a Function

1) A

2) A

3) B

4) A

5) A

6) A

7) A

8) A

9) A

10) D

11) C

12) A

13) A

14) B

15) A


16) A

17) A

18) A

19) A

- 20) B
- 21) A
- 22) A
- 23) D
- 24) A
- 25) A
- 26) A
- 27) A



28)

- 29) A
- 30) A

2.3 Properties of Functions

1 Determine Even and Odd Functions from a Graph

- 1) A
- 2) A
- 3) C
- 4) C
- 5) B
- 6) B
- 7) B
- 8) A

2 Identify Even and Odd Functions from the Equation

- 1) B
- 2) A
- 3) A
- 4) C
- 5) B
- 6) C
- 7) A
- 8) A
- 9) B
- 10) B
- 11) B

3 Use a Graph to Determine Where a Function Is Increasing, Decreasing, or Constant

- 1) A
- 2) A
- 3) A
- 4) A
- 5) A
- 6) A
- 7) A
- 8) A

- 9) A
- 10) A
- 11) A
- 12) A
- 13) A
- 14) A
- 15) A
- 16) A

4 Use a Graph to Locate Local Maxima and Local Minima

- 1) A
- 2) A
- 3) A
- 4) A
- 5) A

5 Use Graphing Utility to Approx Local Maxima and Local Minima and Determine Where a Func Is Incrs or Decrs

- 1) A
- 2) A
- 3) A
- 4) A
- 5) A
- 6) A
- 7) A
- 8) A
- 9) A
- 10) A
- 11) A
- 12) A
- 13) A
- 14) A
- 15) A
- 16) A
- 17) A

6 Find the Average Rate of Change of a Function

- 1) A
- 2) A
- 3) A
- 4) A
- 5) A
- 6) A
- 7) A
- 8) A
- 9) A
- 10) A
- 11) A
- 12) A
- 13) A
- 14) A
- 15) A
- 16) A
- 17) A
- 18) A
- 19) A
- 20) A
- 21) A

- 22) A
- 23) A
- 24) A
- 25) A

2.4 Library of Functions; Piecewise-defined Functions

1 Graph the Functions Listed in the Library of Functions

- 1) A
- 2) A
- 3) A
- 4) A
- 5) A
- 6) A
- 7) A
- 8) A
- 9) A
- 10) A
- 11) A
- 12) A
- 13) A
- 14) A
- 15) A
- 16) A

2 Graph Piecewise-defined Functions

- 1) A
- 2) A
- 3) A

2.5 Graphing Techniques: Transformations

1 Graph Functions Using Vertical and Horizontal Shifts

- 1) A
- 2) A
- 3) A
- 4) A
- 5) A
- 6) A
- 7) A
- 8) A
- 9) A
- 10) A
- 11) A
- 12) A
- 13) A
- 14) A
- 15) A
- 16) A
- 17) A
- 18) A
- 19) A
- 20) A
- 21) A
- 22) A
- 23) A
- 24) A
- 25) A
- 26) A

- 27) A
- 28) A
- 29) A
- 30) A
- 31) A
- 32) A
- 33) A
- 34) A
- 35) A
- 36) A

2 Graph Functions Using Compressions and Stretches

- 1) A
- 2) A
- 3) A
- 4) A
- 5) A
- 6) A
- 7) A
- 8) A
- 9) A
- 10) A
- 11) A
- 12) A
- 13) A
- 14) A
- 15) A
- 16) A
- 17) A
- 18) A

3 Graph Functions Using Reflections about the x-Axis and the y-Axis

- 1) A
- 2) A
- 3) A
- 4) A
- 5) A
- 6) A
- 7) A
- 8) A
- 9) A
- 10) A
- 11) A
- 12) A
- 13) A
- 14) A
- 15) A
- 16) A
- 17) A
- 18) A
- 19) A

2.6 Mathematical Models: Building Functions

1 Build and Analyze Functions

- 1) A
- 2) A
- 3) A

4) A

5) A

6) A

7) A

8) $A(x) = \frac{1}{2}x^3$

9) A

10) $A(x) = \frac{4\sqrt{3} + 9}{16}x^2 - \frac{15}{2}x + 25; \{x | 0 \leq x \leq \frac{20}{3}\}$

11) A

12) $V(s) = \frac{1}{6}s^3$

13) A

14) A

15) A

16) A


17) A

18) D

19) a. $R(x) = -\frac{1}{10}x^2 + 100x$

b. $R(450) = \$24,750.00$

c.



d. 500; \$25,000.00

e. \$50.00

20) $d(t) = \sqrt{1709t}$

21) A